

TRUSTED SINCE 1926

BONIDE®



BONIDE EIGHT DUST, EPA Reg. No. 4-409

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use only in and around the house.

For Dusting Plants: Be sure to lightly cover all plant surfaces. Do not apply when bees are active in the area. Do not apply if rain is expected within 24 hours. Do not apply if wind causes drift.

Protects Your Roses, Flowers & Vegetables From: Alfalfa Loopers, Ants, Aphids, Armyworms, Bagworms, Beetles, Borers, Brown Dog Ticks, Budworms, Cabbageworms, Chinch Bugs, Cicadas, Citrus Black Flies, Codling Moths, Corn Earworms, Crickets, Cutworms, Earwigs, Elm Spanworms, Fall Cankerworms, Fire Ants, Fleas, Green Fruitworms, Gnats, Grasshoppers, Inchworms, Japanese Beetles, Leafhoppers, Leafminers, Leafrollers, Lone Star Ticks, Lygus Bugs, Mealybugs, Millipedes, Mole Crickets, Moths, Navel Orange Worms, Scales, Sowbugs (Pillbugs), Spider Mites, Spring Cankerworms, Tent Caterpillars, Thrips, Tobacco Budworms, Weevils, Whiteflies and Many Others.

VEGETABLES
For Home Vegetable Gardens: Dust upper and lower leaf surfaces. Use sufficient amount to obtain full coverage of all foliage. Apply no more than 13 ounces per 100 square feet.

Sites	Pests	How to Use
Asparagus (pre-harvest)	Asparagus beetles, Cutworms	Apply as needed to ensure uniform coverage. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
(post -harvest)	Larval stages of asparagus beetles, tarnished plant bugs; lygus bugs and adult Japanese beetles	Treat ferns or bush growth after spear harvest when insects are present
Celery - See Leafy Vegetables		
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet armyworms, Cabbage aphids, Cabbage loopers, Diamond back moths, imported cabbageworms	Apply every 5 days or as needed. Do not make more than 8 applications on Cauliflower. Brussels Sprouts and Broccoli and 10 applications on Cabbage. Can be applied up to 2 days prior to harvest.
Corn - See Sweet Corn		
Cucurbits; Balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon	Aphids, cabbage loopers, cucumber beetles (adults), cutworms, leafhoppers, leafminers, melonworms, pickleworms, plant bugs, squash bugs.	Apply as needed to ensure uniform coverage. Do not make more than 8 applications per crop. Can be applied up to the day of harvest.
Eggplant	Colorado potato beetles, Vegetable leaf miners	Apply as needed to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.
Horseradish	Imported crucifer weevils	Apply as needed. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.
Leafy Vegetables: Amaranth, Arugula, Celery, Celtuce, Chervil, Corn salad, Chrysanthemum (edible-leaved and garland), Cress (garden and upland), Dandelion, Dock, Endive, Fennel, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Rhubarb, Spinach, Swiss chard	Alfalfa loopers, aphids, beet armyworms, cabbage loopers, corn earworms, cutworms, European corn borers, fall armyworms, green cloverworms, southern armyworms, tobacco budworms, vegetable leafminers.	Apply every 5 to 10 days as needed. Do not apply more than 7 applications per crop. Can be applied up to 1 day prior to harvest.
Melons - See Cucurbits		
Potatoes	Beet armyworms, Colorado potato beetles, Potato leafhoppers, Potato tuberworms, cabbage loopers, cutworms, European corn borers, Potato psyllids, tarnished plant bugs.	Apply as needed. Do not apply more than 12 applications. Can be applied up to 7 days prior to harvest.
Peppers	Vegetable leafminers, cabbage loopers, Flea beetles, Corn earworms, Pepper weevils, European corn borers.	Apply as needed. Do not apply more than 8 applications per crop. Can be applied up to 3 days prior to harvest.
Sweet Corn	Corn earworms, European corn borers, Fall armyworms, Southern armyworms	Apply every 5 days and when needed. Do not apply more than 6 applications per season. Can be applied up to 1 day prior to harvest.
Tomatoes: (Do not apply on cherry tomatoes and varieties yielding tomatoes less than one inch in diameter.)	Beet and southern armyworms, Tomato pinworms, Tomato fruitworms, Hornworms, Granulate cutworms, Colorado potato beetles, Cabbage loopers.	Apply as needed for uniform coverage. Do not apply more than 6 applications per season. Can be applied up to day of harvest.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

DIRECTIONS FOR APPLYING ROSES & FLOWERS

Sites	Pests	How to Use
Ageratum, Aster, Ardisia, Azalea, Baby's Breath, Begonia, Birds-nest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lily, Marigold, Mock-orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	Aphids, Armyworms, Bagworms, Cabbage Worms, Cicadas, Corn Earworms, Cutworms, Exposed Thrips, Fall Cankerworms, Fall Webworms, Fungus Gnats, Inchworms, Japanese Beetles, Leafminers, Leafrollers, Mealybugs, Scale Crawlers, Spring Cankerworms, Spider Mites, Whiteflies, Rose Chafers, Psyllids, Scale Insects	Apply at the first sign of insects. Lightly cover all plant surfaces. Repeat as necessary; use at intervals of 4-8 days. Applications can be made up to the day of collection.

Use with care on ferns, palms and fuschia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, dust a few plants as described above, wait 24 hours, and check for possible injury.

Repeat applications may be necessary to control citrus blackflies on non-bearing citrus and adult Japanese beetles on roses. For Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. A few caterpillars maybe seen escaping from the tent; however, they should die within several hours.

OTHER OUTDOOR AREAS

Use **Bonide Eight Insect Control Garden Dust** on **Ants, Crickets, Earwigs, Fleas, Lone Star Ticks** for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, dust infested areas thoroughly. **For Ants,** thoroughly treat hills and runways. Repeat application as reinfestations occur.

PRECAUTIONARY STATEMENTS-HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION: May be harmful if swallowed or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water before smoking or eating after handling. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before dusting. Do not use on humans. Do not allow children or pets to contact treated areas.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

If Swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on Skin or Clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If Inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

If in Eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Disposal: *If empty* - Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

**BONIDE PRODUCTS, INC.
 ORISKANY, NY 13424**