

Herbicide

Precautionary Statements

Hazardous to Humans and Domestic Animals

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes Eye Irritation • Prolonged or frequently repeated contact may cause allergic reactions in some individuals. Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Mixers and loaders must wear a chemical-resistant apron in addition to other PPE.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Shake Well Before Using
Net Contents 1 Pint

Table of Contents

Page

Precautionary Statements.....	Front
Hazards to Humans and Domestic Animals.....	Front
First Aid.....	3
Environmental Hazards.....	3
Directions for Use.....	3
Agricultural Use Requirements.....	4
Storage and Disposal.....	5
General Information.....	5
Soil Preparation.....	6
Mixing Directions.....	7
Surflan Pro Alone.....	7
Surflan Pro Tank Mix Combinations.....	7
Mixing Order.....	7
Premixing.....	8
Equipment Cleaning.....	8
Activation and Cultivation.....	8
Weeds Controlled by Surflan Pro.....	8
Weeds Suppressed by Surflan Pro.....	10
Crop-Specific Use Directions.....	11
Ornamental Plantings.....	11
Broadcast Application Rates.....	11
Tank Mix Combinations.....	12
Special Use Precautions.....	12
Recommended Species.....	13
Field and Container Grown Species.....	13
Non-bearing Trees and Vines.....	29
Ornamental Bulbs.....	30
Broadcast Application Rates.....	30
Special Use Precautions.....	30
Shadehouse Areas.....	31
Christmas Tree Plantations.....	31
Surflan Pro Alone.....	31
Broadcast Application Rates.....	31

Table of Contents (continued)	Page
Tank Mix Combinations	31
Special Use Precautions	32
Noncropland Areas and Industrial Sites	32
Noncropland Areas—Tank Mix Combinations	32
Broadcast Application Rates	32
Industrial Sites—Tank Mix Combinations	32
Warm Season Turfgrasses	32
Annual Grasses Controlled by Surflan Pro	33
Annual Broadleaf Weeds Controlled by Surflan Pro	34
Broadleaf Weeds Suppressed by Surflan Pro	34
Application Rates, Frequency and Timing of Application	35
Summer Annual Grasses and Broadleaf Weeds	35
Split Application Program	35
Annual Bluegrass (<i>Poa annua</i>) and Winter Annual Broadleaf Weeds	35
Weed Control in Florida	36
Application Equipment	36
Reseeding	36
Special Use Precautions	36
Terms and Conditions of Use	37
Warranty Disclaimer	37
Inherent Risks of Use	37
Limitation of Remedies	37

Engineering Controls Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center 1-800-858-7378 for emergency medical treatment.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spills.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. **Exception:** If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Workers may enter treated areas without required PPE during the reentry interval following 1/2 to 1 inch of rainfall or irrigation, if they are performing tasks that do not involve contact with the soil subsurface; otherwise, PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Keep all persons, children and pets out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Surflan Pro herbicide is a preemergence surface-applied product for the control of many annual grasses and broadleaf weeds in ornamental plantings, bulbs, ground covers/perennials, established warm-season turfgrass, Christmas tree plantations, non-bearing trees and vines, and noncropland and industrial sites.

Surflan Pro is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Blazon or removed by spraying surface with water or washing with an industrial cleaner immediately after application. Surflan Pro may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

Treatment of Plant Species Not Listed on the Label for Surflan Pro

Users who wish to use Surflan Pro on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury during 30-60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Surflan Pro on plant species not recommended on this label.

Aerial Application: Do not aerially apply this product.

Chemigation: Do not apply this product through any type of irrigation system.

For orchard crops, including citrus, pome fruits, stone fruits, and tree nuts, apply product only as a strip treatment in the tree rows; do not apply to row middles or drive rows.

Do not graze or feed forage from treated areas to livestock.

Precaution: Avoid drift to non-target areas when applying Surflan Pro. Spray drift may result in reduced emergence of non-target plants adjacent to the treated area. Poor weed control may result if directions are not followed. Over-application may result in crop injury or excessive soil residue.

Application

Soil Preparation

Surflan Pro controls weeds growing from seed. Surflan Pro will not control emerged weeds. Surflan Pro does not control established weeds growing from stolens, rhizomes, or root pieces. Therefore, areas to be treated should be free of emerged weeds. Weed residues, prunings, and trash should be thoroughly mixed into the soil or removed prior to treatment. In field applications, the soil should be in good tilth and free of clods at the time of application.

Ground Application: Apply Surflan Pro as a directed spray to the soil surface or over the top of plants. Use only a properly calibrated, low-pressure, herbicide sprayer that will apply the spray uniformly. Use screens no finer than 50 mesh in nozzles and in-line strainers. Apply the appropriate rate of Surflan Pro, as outlined in "Approved Uses" section of this label. In all cases, use sufficient water volume to obtain uniform coverage and deliver the desired rate of Surflan Pro to the treated area. The volume of water used is not critical, as long as the desired rate Surflan Pro is delivered uniformly across the area treated. When calibrating, determine the volume of water delivered by the sprayer to a given area (1,000 sq ft, 1 acre, etc.). Then mix the desired rate of Surflan Pro in the amount of water required to cover the entire area to be treated. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to ensure proper calibration and uniform application. Maintain continuous agitation from mixing through application. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application.

Hand Held or Backpack Sprayer Application: The amount of water used to apply Surflan Pro herbicide is not critical, but should be sufficient for uniform coverage of the target area. Calibrate by determining the volume of water required to treat 1,000 square feet. Use this calibration volume to determine the amount of water and Surflan Pro herbicide needed to treat

Steps in Calibration:

1. Mark an area of 1,000 square feet (i.e. 20 by 50 feet, or 25 by 40 feet).
2. Place the sprayer on a level surface and add water noting the final level of water in the spray tank.
3. Spray the marked area with a sufficient volume of water to provide uniform coverage. Refill the sprayer to the same level as before measuring the amount of water added. The measured water added to the sprayer is the volume needed to cover 1,000 square feet.
4. Determine the application rate (fl oz/1,000 sq ft) for Surfian Pro from the "Approved Uses" section of this label.
5. To each volume of water used, as measured in step 3, add the amount of Surfian Pro as determined in step 4.

Example: If the sprayer used 2 gallons of water to cover 1,000 square feet and the desired application rate of Surfian Pro is 3 fluid oz/1,000 square feet, then you would add 3 fluid ounces of Surfian Pro to every 2 gallons of water to be used.

Mixing Directions

Shake Well Before Using

Precaution: Do not allow the spray mixture to siphon back into water source.

Surfian Pro - Alone

Make sure spray tank is clean and use only clean water. Fill spray tank $\frac{1}{2}$ - $\frac{3}{4}$ full. Start agitation and add the required amount of Surfian Pro. Continue agitation and finish filling the spray tank. Maintain continuous agitation until application is completed.

Surfian Pro - Tank Mix Combinations

Prior to mixing, read and carefully follow all label instructions and precautions for each product added to the tank mixture. Vigorous, continuous agitation is required for all tank mixes of Surfian Pro. Sparger pipe agitators generally provide the best agitation in spray tanks.

Mixing Order: Fill the tank $\frac{3}{4}$ full with clean water. Start agitation and add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: dry flowables (DF); wettable powders (WP); Surfian Pro and other aqueous suspensions (AS), flowables (F), and liquids (L); solutions (S); and emulsifiable concentrates (EC).

Continue agitation and finish filling the spray tank with clean water. Maintain agitation until application is completed. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be completely resuspended before spraying is continued. A sparger agitator is particularly useful for this purpose.

Premixing: When tank mixing, initial mixing and dispersion of certain dry flowable or wettable powder products may be improved by premixing with water (slurrying). Adding the slurried material to the spray tank through a wetting screen of 20 or 35 mesh will help assure good initial dispersion.

Equipment Cleaning

If a buildup of material occurs on the walls of the spray tank, it should be removed between fillings by washing with soap and water and rinsing thoroughly. Tanks, lines, screens, and nozzles should be cleaned thoroughly after each use.

Activation and Cultivation

Surflan Pro will remain stable on the soil surface up to 21 days following application. In the absence of timely rainfall, irrigation can be used to activate Surflan Pro. A minimum of one-half (1/2) inch of rain or its equivalent in sprinkler irrigation is necessary to activate Surflan Pro. If weeds begin to emerge due to lack of rainfall or irrigation, shallow cultivate 1-2 inches deep to destroy existing weeds, or remove them by hand. Shallow cultivation to a depth of 1-2 inches will enhance herbicidal effectiveness. Erratic weed control may result if Surflan Pro is not activated by rainfall, irrigation, or cultivation within 21 days of application, or existing weeds have not been removed.

Weeds Controlled by Surflan Pro

Annual Grasses:

Common Name

barley, little
barnyardgrass
(watergrass)
bluegrass, annual
crabgrass, large
crabgrass, smooth
crowfootgrass
cupgrass, southwestern

Scientific Name

Hordeum pusillum
Echinochloa crus-galli

Poa annua
Digitaria sanguinalis
Digitaria ischaemum
Dactyloctenium aegyptium
Eriochloa gracilis

Annual Grasses: (continued)**Common Name**

foxtail, bristlegress
foxtail, giant
foxtail, green
(pigeongrass)
foxtail, robust
foxtail, yellow
goosegrass
(silver crabgrass)
Johnsongrass
(seedling only)
junglerice
lovegrass, Mexican
lovegrass, orcutt
oat, wild
panicum, browntop
panicum, fall
(spreading panicgrass)
panicum, Texas
(buffalograss)
(Coloradograss)
ryegrass, Italian
signalgrass (Brachiaria)
sprangletop, red
witchgrass

Broadleaf Weeds:**Common Name**

bittercress
carpetweed
chickweed, common
fiddleneck, coast
filaree, redstem
filaree, whitestem

Scientific Name

Setaria magna
Setaria faberi
Setaria viridis

Setaria robusta
Setaria glauca
Eleusine indica

Sorghum halepense

Echinochloa colonum
Eragrostis mexicana
Eragrostis orcuttiana
Avena fatua
Panicum fasciculatum
Panicum dichotomiflorum

Panicum texanum

Cenchrus incertus
Brachiaria spp.
Leptochloa filiformis
Panicum capillare

Scientific Name

Cardamine oligosperma
Mollugo verticillata
Stellaria media
Amsinckia intermedia
Erodium cicutarium
Erodium moschatum

Broadleaf Weeds: (continued)

Common Name

groundsel, common
henbit
knotweed, prostrate
lambquarters
pigweed, prostrate
pigweed, redroot
pigweed, spring
pigweed, tumble
puncturevine
purslane, common
pusley, Florida
(Florida purslane)
(Mexican clover)
(pusley)
rocket, London
rockpurslane, desert
shepherdspurse
spurge, prostrate
woodsorrel, yellow

Scientific Name

Senecio vulgaris
Lamium amplexicaule
Polygonum aviculare
Chenopodium album
Amaranthus blitoides
Amaranthus retroflexus
Amaranthus hybridus
Amaranthus albus
Tribulus terrestris
Portulaca oleracea
Richardia scabra

Sisymbrium irio
Calandrinia ciliata
Capsella bursa-pastoris
Euphorbia humistrata
Oxalis stricta

Weeds Suppressed by Surflan Pro

Control of the following weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, and amount and timing of soil moisture:

Common Name

horseweed
ladysthumb
lettuce, prickly
mallow, common
milkweed, climbing
morningglory
mustard, black

Scientific Name

Coryza canadensis
Polygonum persicaria
Lactuca serriola
Malva neglecta
Sarcostemma cynanchoides
Ipomoea spp.
Brassica nigra

Common Name

mustard, wild
 nightshade, black
 ragweed, common
 smartweed
 sowthistle, annual
 spurge, spotted
 teaweed (prickly sida)
 velvetleaf
 wheat, volunteer

Scientific Name

Brassica kaber
Solanum nigrum
Ambrosia artemisiifolia
Polygonum pensylvanicum
Sonchus oleraceus
Euphorbia maculata
Sida spinosa
Abutilon theophrasti
Triticum spp.

Crop-Specific Use Directions

Ornamental Plantings

Surflan Pro is recommended for use on certain landscape container- and field-grown established ornamental plants including: trees, shrubs, ground covers/perennials, flowers, non-bearing fruit and nut trees, non-bearing vineyards; and in the production of ornamental bulbs (See "Ornamental Bulbs" section for special use directions).

Broadcast Application Rates

Labeled Use Site	Length of Control	Surflan Pro		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
		(qt/acre)	(fl oz/1000 sq ft)		
Landscape Ornamentals	2 - 4 months	2	1.5	2	8
	3 - 6 months	3	2.2	4	12
	4 - 8 months	4	3	4	12
Field-grown and container-grown ornamentals	2 - 4 months	2	1.5	3	8
	3 - 6 months	3	2.2	3	9
	4 - 8 months	4	3	3	12

Tank Mix Combinations

Tank mix combinations of Surflan Pro plus glyphosate, and many other labeled herbicides may be used to control undesirable vegetation in ornamental areas. Surflan Pro may also be tank mixed with Gallery herbicide and applied preemergence to broaden the spectrum of broadleaf weed control in ornamental areas. Applied as directed, these tank mixes of Surflan Pro will provide control of susceptible weed species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Surflan Pro Plus Glyphosate: Tank mix combinations of Surflan Pro plus glyphosate are recommended to control existing undesirable vegetation. Applied as directed, Surflan Pro plus glyphosate will provide postemergence control of susceptible weed species listed on the label for glyphosate and residual preemergence control of susceptible weed species listed on the label for Surflan Pro. Refer to the label for glyphosate for specific use directions, precautions, and limitations before use.

Precautions: Do not apply sprays containing glyphosate over the top of ornamental plants. Extreme care must be exercised to prevent sprays containing glyphosate from coming in contact with foliage and stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying with glyphosate in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

Special Use Precautions:

Apply only to established plants that have been transplanted into their growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Routed liners should be removed from their original growing containers and placed in new containers at least two weeks prior to treatment or injury may occur.

To avoid possible injury, do not apply Surflan Pro to:

- Nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds.
- Unrooted liners or cuttings that have been planted in pots for the first time.
- Pots less than four inches wide.
- Ground covers until they are established and well rooted.
- Ornamental plantings where there is likelihood of runoff onto lawn areas.
- Areas containing dichondra or cool season turfgrass species.

On container grown ornamentals where weed seed germination continues for extended periods of time, do not make repeat applications of Surflan Pro for at least 90 days or crop injury may occur.

Applications of Surflan Pro over the top of plants with newly forming buds may cause injury. In this situation a directed spray is recommended.

For soils treated with Surflan Pro during the previous season, plant only the ornamental species listed on this label or injury may occur.

Ice Plant: When establishing unrooted ice plant on coarse-textured soils in landscape plantings, do not exceed the 2 quart per acre rate of Surflan Pro or crop injury may occur.

Note: Injury on the following plant species has been observed following applications of Surflan Pro and use is not recommended:

Deutzia gracilis (slender deutzia)

Pseudotsuga menziesii (Douglas-fir)

Thuja occidentalis 'Techny' (Techny arborvitae)

Tsuga canadensis (eastern hemlock)

Begonia spp. (begonia)

Coleus hybridus (coleus)

Surflan Pro May be Used on the Following Established Plant Species:
(Note: Limitations on recommended treatment methods).

Trees

Recommended Treatment Method

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Abies balsamea</i>	Fir, balsam	F
<i>Abies concolor</i>	Fir, white	F
<i>Abies fraseri</i>	Fir, fraser	F
<i>Abies grandis</i>	Fir, grand	F
<i>Abies veitchi</i>	Fir, Vietch	F
<i>Abies lasiocarpa</i>	Fir, alpine	F

(continued)

Trees (cont.)

Scientific Name	Common Name	Recommended Treatment Method
		F = Field Grown C = Container Grown
<i>Abutilon hybridum</i>	Albus-flowering maple	F
	Luteus-flowering maple	F
	Roseus-flowering maple	F
	Tangerine-flowering maple	F
	Vesuvius red-flowering maple	F
<i>Acer ginnala</i>	Flame maple	F
<i>Acer rubrum</i>	Red sunset maple	F
<i>Acer saccharinum</i>	Silver maple	F
<i>Acer</i> spp.	Maple	F
<i>Alsophila australis</i>	Australian tree fern	C,F
<i>Areacastrum romanzoffianum</i>	Queen palm	F
<i>Betula nigra</i>	Birch, river	F
<i>Betula papyrifera</i>	Paper birch	F
<i>Betula pendula</i>	Birch, white	F
<i>Bucida buceras</i>	Black olive	F
<i>Carya</i> spp.	Pecan, ornamental	C,F
<i>Cedrus, atlantica</i>	Atlas cedar	C,F
<i>Cedrus deodara</i>	Deodar cedar	C,F
<i>Ceratonia siliqua</i>	Carob	F
<i>Cercidium floridum</i>	Palo Verde, blue	F
<i>Cercis canadensis</i>	Redbud	C,F
<i>Chamaecyparis lawsoniana</i>	Falsecypress, Lawson	F
<i>Chamaecyparis obtusa</i>	Filicoides-fernspray cypress	F
	Gracilis-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i>	Sawara-false cypress	F
	Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	Cat Palm	F
<i>Chamaedorea costaricana</i>	Palm	F
<i>Chamaedorea elegans</i>	Parlor palm	F
<i>Citrus</i> spp.	Citrus, ornamental	C,F

Trees (cont.)

	Recommended Treatment Method
	F = Field Grown
	C = Container Grown
Scientific Name	Common Name
<i>Cornus florida</i>	Dogwood, flowering F
<i>Cryptomeria japonica</i>	Cryptomeria, Japanese C,F
<i>Cupaniopsis anacardioides</i>	Carrot wood F
<i>Cupressus arizonica</i> (<i>glabra</i>)	Cypress, Arizona C,F
<i>Cupressus glabra</i>	Arizona cypress C,F
<i>Cupressocyparis leylandii</i>	Leyland cypress C,F
<i>Cupressus sempervirens</i>	Cypress, Italian C,F
<i>Dicksonia antarctica</i>	Tasmanian tree fern C,F
<i>Elaeagnus angustifolia</i>	Russian olive C,F
<i>Eucalyptus camaldulensis</i>	Red gum eucalyptus F
<i>Eucalyptus cinerea</i>	Eucalyptus, mealy F
	Silver dollar eucalyptus F
<i>Eucalyptus nicholii</i>	Eucalyptus, narrow-leaved F
<i>Eucalyptus sideroxylon</i>	Eucalyptus, red ironbark F
<i>Ficus benjamina</i>	Ficus F
<i>Fraxinus</i> spp.	Ash F
<i>Ginkgo biloba</i>	Ginkgo (Maidenhair tree) C,F
<i>Gleditsia triacanthos</i>	Honey locust F
<i>Heteromeles arbutiflora</i>	Toyon F
<i>Juniperus virginiana</i>	Redcedar, Eastern F
<i>Koelreuteria paniculata</i>	Goldenrain tree F
<i>Liquidambar styraciflua</i>	Sweetgum, American C,F
<i>Magnolia</i> spp.	Magnolia F
<i>Malus</i> spp.	Crabapple F
<i>Morus alba</i>	White mulberry F
<i>Picea abies</i>	Pendula-weeping Norway spruce F
	Repens-spreading Norway spruce F
	Spruce, Norway F
<i>Picea englemanni</i>	Spruce, Englemann F
<i>Picea glauca</i>	Spruce, white F
	Conica-dwarf Alberta spruce F
<i>Picea glauca conica</i>	Dwarf Alberta spruce F

(continued)

Trees (cont.)

		Recommended Treatment Method
		F = Field Grown
		C = Container Grown
Scientific Name	Common Name	
<i>Picea mariana</i>	Spruce, black	F
<i>Picea pungens</i>	Glauca-Colorado blue spruce	F
	Hoopsii-Hoop's blue spruce	F
	Koster-Koster blue spruce	F
	Spruce, Colorado	C,F
<i>Pinus aristata</i>	Bristlecone pine	F
<i>Pinus canariensis</i>	Canary Island pine	F
<i>Pinus contorta</i>	Shore pine, beach pine	F
<i>Pinus eldarica</i>	Eldarica pine	F
<i>Pinus halepensis</i>	Aleppo pine	C,F
<i>Pinus radiata</i>	Monterey pine	F
<i>Pinus</i> spp.	Pine	C,F
<i>Pinus strobus</i>	Eastern white pine	F
<i>Pinus sylvestris</i>	Scotch pine	F
<i>Pinus thunbergiana</i>	Japanese black pine	F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	California sycamore	F
<i>Podocarpus</i> spp.	Podocarpus	F
<i>Populus deltoides</i>	Cottonwood	F
	Cottonwood (grown for pulp)	F
<i>Prunus caroliniana</i>	Laurelcherry, Carolina	F
<i>Prunus glandulosa</i>	Dwarf flowering almond	C,F
<i>Prunus laurocerasus</i>	Laurelcherry, English	F
<i>Prunus mahaleb</i>	Cherry, Mahaleb	F
<i>Prunus yedoensis</i>	Yoshino flowering cherry	F
<i>Pyrus communis</i>	Pear	F
<i>Quercus palustris</i>	Pin oak	F
<i>Quercus phellos</i>	Willow oak	F
<i>Quercus rubra</i>	Red oak	C,F
<i>Quercus</i> spp.	Oak	C,F
<i>Salix babylonica</i>	Babylon weeping willow	F
	Corkscrew willow	F

(continued)

Trees (cont.)

Scientific Name	Common Name	Recommended Treatment Method	
		F = Field Grown	C = Container Grown
<i>Schinus molle</i>	California pepper tree		F
<i>Sequoia sempervirens</i>	Redwood, coast		F
<i>Sequoiadendron giganteum</i>	Giant sequoia		F
<i>Swietenia mahogani</i>	Mahogany		F
<i>Tabebuia caraiba</i>	Yellow tab		F
<i>Tilia cordata</i>	Linden, little leaf		C,F
<i>Ulmus parvifolia</i>	Chinese elm		F
<i>Umbellularia californica</i>	California laurel		F
<i>Washingtonia robusta</i>	Mexican fan palm		F

Shrubs

Scientific Name	Common Name	Recommended Treatment Method	
		F = Field Grown	C = Container Grown
<i>Abelia grandiflora</i>	Glossy abelia		F
<i>Acacia redolens</i>	Acacia, prostrate		F
<i>Agave americana</i>	Century plant		F
<i>Agave macroculmis</i>	Agave		F
<i>Anisodonteia hypomandarum</i>	Cape mallow		C,F
<i>Arctostaphylos stanfordiana</i>	Manzanita, Stanford		F
<i>Astilbe chinensis</i>	Astilbe/false spirea		C,F
<i>Baccharis pilularis</i>	Coyotebush		F
<i>Berberis thunbergii</i>	Aurea-golden Japanese barberry		C,F
	Crimson pygmy barberry		C,F
	Atropurea-redleaf Japanese barberry		C,F
	Barberry, Japanese		C,F
	Barbara Karst		F
	California gold		F
	Scarlet O'Hara		F
	Texas dawn		F

(continued)

Shrubs (cont.)

Scientific Name	Common Name	Recommended Treatment Method
		F = Field Grown C = Container Grown
<i>Buddleia davidii</i>	Butterfly bush	C,F
<i>Buxus microphylla</i>	Littleleaf boxwood	F
<i>Buxus microphylla japonica</i>	Boxwood, Japanese	C,F
<i>Buxus sempervirens</i>	Boxwood, common	C,F
<i>Callistemon citrinus</i>	Bottlebrush, lemon	C,F
<i>Cassia artemisioides</i>	Cassia, feathery	F
<i>Ceanothus americanus</i>	Jerseytea, redroot	C,F
<i>Ceanothus</i> spp.	Wild lilac	C,F
<i>Chaenomeles japonica</i>	Flowering quince	C,F
<i>Chamaecyparis obtusa</i>	Kosteri cypress	F
	Nana-dwarf Hinoki cypress	F
	Torulosa cypress	F
<i>Chamaecyparis pisifera</i>	Squarrosa Minima cypress	F
<i>Chamaecyparis pisifera</i> spp.	Filifera-thread cypress	F
<i>Chrysalidocarpus lutescens</i>	Areca palm	F
<i>Clethra</i>	Summersweet	C,F
<i>Cleyera japonica</i>	Cleyera, Japanese	C,F
<i>Coleonema pulchrum</i>	Pink breath of heaven	C,F
<i>Cornus alba</i>	Sibirica-Siberian dogwood	F
<i>Cornus kousa</i>	Dogwood, kousa	C,F
<i>Cornus stolonifera</i>	Flaviramea-yellowtwig dogwood	F
<i>Cotoneaster adpressus</i>	Prae-cox-early cotoneaster	F
<i>Cotoneaster apiculatus</i>	Cotoneaster, cranberry	C,F
<i>Cotoneaster buxifolius</i>	Cotoneaster, brightbead	F
<i>Cotoneaster congestus</i>	Cotoneaster, Pyrenees	F
<i>Cotoneaster dammeri</i>	Cotoneaster, bearberry	C,F
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster	F
<i>Cotoneaster horizontalis</i>	Cotoneaster, rock	C,F
<i>Cotoneaster lacteus</i>	Cotoneaster, parney	C,F
<i>Cotoneaster microphyllus</i>	Cotoneaster, rockspray	F

Shrubs (cont.)

	Recommended Treatment Method	
	F = Field Grown	C = Container Grown
Scientific Name	Common Name	
<i>Cotoneaster salicifolia</i>	Willowleaf cotoneaster	C,F
<i>Cytisus praecox</i>	Hollandia-warmminster broom	F
<i>Cytisus scoparius</i>	Lena-Scotch broom	F
<i>Dasyliirion wheeleri</i>	Sotol, desert spoon	F
<i>Deutzia crenata</i>	Nakiana-dwarf deutzia	F
<i>Dodonaea viscosa</i>	Hopseedbush, clammy	F
	Hopseed bush	F
<i>Escallonia exoniensis</i>	Escallonia	C,F
<i>Euonymus alata</i>	Euonymus, winged	F
<i>Euonymus fortunei</i>	Canadale gold euonymus	C,F
	Emerald'n gold euonymus	C,F
	Euonymus, stringybark	C,F
	Wintercreeper	C,F
<i>Euonymus japonica</i>	Euonymus, evergreen	C,F
	Silver king euonymus	F
<i>Euonymus kiatschovica</i>	Spreading euonymus	F
<i>Euonymus vegetus</i>	Bigleaf wintercreeper	C,F
<i>Fatsyhedera lizei</i>	Fatsyhedera	C,F
<i>Fatsia japonica</i>	Japanese aralia	C,F
<i>Felicia amelloides</i>	Blue marguerite	C,F
<i>Forsythia intermedia</i>	Forsythia, border	F
<i>Gardenia jasminoides</i>	Gardenia	C,F
<i>Genista pilosa</i>	Woadwaxen	F
<i>Hibiscus rosa-sinesis</i>	Ross Estey-hibiscus	F
	Hibiscus, Chinese	F
<i>Hibiscus syriacus</i>	Rose of Sharon, Red Bird	F
	Rose of Sharon, Red Heart	F
	Rose of Sharon, Woodbridge	F
	Rose-of-Sharon (Shrubalthea)	F
<i>Hydrangea macrophylla</i>	Hydrangea, French	C,F

(continued)

Shrubs (cont.)

Scientific Name	Common Name	Recommended Treatment Method	
		F = Field Grown	C = Container Grown
<i>Hydrangea quercifolia</i>	Hydrangea, Oakleaf		C,F
<i>Ilex aquifolium</i>	Balkans holly		F
	Gold coast holly		F
	Holly, English		F
<i>Ilex aquipernyi</i>	San Jose holly		C,F
<i>Ilex cornuta</i>	Dwarf Burford holly		C,F
	Holly, Chinese		C,F
<i>Ilex crenata</i>	Compacta-dwarf Japanese holly		C,F
	Convexa holly		C,F
	Helleri-Heller's Japanese holly		C,F
	Holly, Japanese		C,F
<i>Ilex glabra</i>	Nordica-inkberry holly		F
<i>Ilex meserveae</i>	Blue boy holly		F
	Blue girl holly		F
	Ebony magic holly		F
<i>Ilex vomitoria</i>	Nana-dwarf yaupon holly		C,F
	Pendula-weeping yaupon holly		C,F
	yaupon holly		C,F
<i>Juniperus chinensis</i>	Media-old gold juniper		C,F
<i>Juniperus conferta</i>	Emerald sea shore juniper		F
<i>Juniperus horizontalis</i>	Huntington blue juniper		C,F
	Wiltonii-blue carpet juniper		C,F
<i>Juniperus procumbens</i>	Nana-dwarf Japaneses garden juniper		C,F
<i>Juniperus prostrata</i>	Prostrate juniper		C,F
<i>Juniperus sabina</i>	Broadmoor juniper		F
	Foemina-Hicks juniper		F
	Tamariscifolia-Tam juniper		F
<i>Juniperus scopulorum</i>	Emerald green juniper		F
<i>Juniperus</i> spp.	Juniper		C,F

(continued)

Shrubs (cont.)

Scientific Name	Common Name	Recommended Treatment Method	
		F = Field Grown	C = Container Grown
<i>Juniperus squamata</i>	Blue juniper		F
	Blue star juniper		F
	Parsonii juniper		F
<i>Justicia brandegeana</i>	Shrimp plant	C,F	
<i>Justicia spicigera</i>	Honeysuckle, Mexican		F
<i>Kalmia latifolia</i>	Laurel, mountain		F
<i>Lagerstroemia indica</i>	Crape myrtle	C,F	
<i>Lavandula angustifolia</i>	English lavender	C,F	
<i>Leucothoe axillaris</i>	Leucothoe, coast		F
<i>Leucothoe fontanesiana</i>	Leucothoe, drooping		F
<i>Ligustrum amurense</i>	Privet, amur	C,F	
<i>Ligustrum japonicum</i>	Privet, Japanese	C,F	
	yellow tip ligustrum	C,F	
<i>Ligustrum lucidum</i>	Privet, glossy	C,F	
<i>Ligustrum ovalifolium</i>	California privet		F
<i>Ligustrum texanum</i>	Howardi privet		F
	Wax leaf privet		F
<i>Ligustrum vicaryi</i>	Privet, golden	C,F	
	Vicary golden privet	C,F	
<i>Livistona chinensis</i>	Chinese fountain palm		F
<i>Lonicera fragrantissima</i>	Winter honeysuckle		F
<i>Lonicera periclymenum</i>	Flowering woodbine		F
	Serotina woodbine		F
<i>Lonicera sempervirens</i>	Trumpet honeysuckle		F
<i>Lorpetalum chinense</i>	(No common name)	C,F	
<i>Mahonia aquifolium</i>	Oregon grape		F
<i>Myoporum parvifolium</i>	Myoporum, prostrate		F
<i>Myrtus communis</i>	Myrtle, true		C,F

(continued)

Shrubs (cont.)

Scientific Name	Common Name	Recommended Treatment Method
		F = Field Grown
		C = Container Grown
<i>Nandina domestica</i>	Compacta-dwarf heavenly bamboo	C,F
	Harbour dwarf-heavenly bamboo	C,F
	Heavenly bamboo (Nandina)	C,F
	Nana compacta-heavenly bamboo	C,F
	Nana purpurea-heavenly bamboo	C,F
	Woods dwarf-heavenly bamboo	C,F
<i>Nerium oleander</i>	Hardy red oleander	C,F
	Oleander	C,F
	Ruby lace oleander	C,F
<i>Osmanthus heterophyllus</i>	Osmanthus, holly-leaf	F
<i>Pachysandra terminalis</i>	Japanese spurge	C,F
<i>Philadelphus</i> spp.	Mockorange	C,F
<i>Phoenix roelofenii</i>	Pigmy date palm	F
<i>Photinia fraseri</i>	Fraser's photinia	C,F
	Photinia	C,F
<i>Pieris japonica</i>	Lily-of-the-valley	F
	Snowdrift lily-of-the-valley	F
	Temple bells lily-of-the-valley	F
	Valley rose lily-of-the-valley	F
	Andromeda	C,F
<i>Pittosporum</i> spp.	Pittosporum	C,F
<i>Pittosporum tobira</i>	Green pittosporum	F
	Japanese pittosporum	F
	Tobira	F
	Wheeler's dwarf pittosporum	F
<i>Platyclusidus orientalis</i>	Arborvitae, Oriental	C,F
<i>Plumbago ariculata</i>	Blue cape plumbago	F
<i>Podocarpus macrophyllus</i>	Yewpine	C,F
<i>Potentilla fragiformis</i>	Cinquefoil	F
<i>Potentilla fruticosa</i>	Cinquefoil	C,F

(continued)

Shrubs (cont.)

	Recommended Treatment Method
	F = Field Grown
	C = Container Grown
Scientific Name	Common Name
<i>Protea neriifolia</i>	Protea F
<i>Pyracantha coccinea</i>	Firethorn, scarlet C,F
<i>Pyracantha fortuneana</i>	Lolendei Monrovia pyracantha C,F
<i>Pyracantha fortuneana</i>	Monon pyracantha C,F
	Red elf hybrid pyracantha C,F
	Rutgers hybrid pyracantha C,F
	Santa Cruz pyracantha C,F
	Victory pyracantha C,F
<i>Pyracantha skoidzumi</i>	Firethorn, formosa C,F
<i>Pyracantha, fortuneana</i>	Firethorn C,F
<i>Rhaphiolepis indica</i>	Enchantress-Moness rhaphiolepis F
	Rhaphiolepis (India hawthorn) C,F
	Springtime-Monme rhaphiolepis F
<i>Rhaphiolepis ovata</i>	Roundleaf rhaphiolepis F
<i>Rhipsalidopsis gaertneri</i>	Eastercactus C,F
<i>Rhododendron calendulaceum</i>	Flame azalea F
<i>Rhododendron campylocarpum</i>	Butterfly rhododendron F
<i>Rhododendron carolinianum x daurium</i>	PJM rhododendron F
<i>Rhododendron catawbiense</i>	Catawba album rhododendron C,F
	Catawba rhododendron C,F
	Lord Roberts rhododendron C,F
	Rocket rhododendron C,F
<i>Rhododendron forrestii x griersonianum</i>	Elizabeth rhododendron F
<i>Rhododendron</i>	America rhododendron F
<i>hybrid spp.</i>	English Roseum rhododendron F
	Nova Zembla rhododendron F
	Scintillation rhododendron F
<i>Rhododendron impeditum</i>	Rhododendron F
<i>Rhododendron indica</i>	Formosa azalea C,F
	Waucabusa azalea C,F

(continued)

Shrubs (cont.)

Scientific Name

Rhododendron kerume

Rhododendron maximum

Rhododendron mucronulatum

Rhododendron satuski

Rhododendron spp.

Rhododendron spp.
hybrids

Rhus lancea

Rosa rugosa

Rosmarinus officinalis

Senecio cineraria

Spiraea vanhouttei

Syringa vulgaris

Syzygium paniculata

Taxus cuspidata

Taxus media

Thuja occidentalis

Recommended Treatment Method

F = Field Grown

C = Container Grown

Common Name

Coral bells azalea

Hino crimson azalea

Hino pink azalea

Snow azalea

Rhodie max (rosebay)

Rhododendron

Gumpo pink azalea

Higasa azalea

Azalea

Rhododendron

Carror azalea

Girard Roberta azalea

Golden flare exbury azalea

Sumac, African

Ramanas rose

Rosemary

Dusty miller

Bridal wreath

Lilac, common

Brush cherry

Yew, Japanese

Yew

Arborvitae, American

Emerald arborvitae

Globosa-globe arborvitae

Little giant-dwarf arborvitae

Nigra-dark American arborvitae

Pyramidalis arborvitae

Rheingold arborvitae

Woodwardii arborvitae

C,F

C,F

C,F

C,F

C,F

F

F

F

C,F

C,F

C,F

F

F

C,F

F

F

C,F

F

C,F

C,F

F

F

C,F

F

F

F

F

F

F

F

Shrubs (cont.)

	Recommended Treatment Method
	F = Field Grown
	C = Container Grown
Scientific Name	Common Name
<i>Thuja orientalis</i>	Aureus nana-dwarf golden arborvitae F
<i>Thuja plicata</i>	Minima glauca-dwarf arborvitae F
<i>Trachelospermum jasminoides</i>	Red Cedar, Western F
<i>Veitchia merrillii</i>	Star jasmine, Chinese F
<i>Viburnum carlesii</i>	Christmas palm F
<i>Viburnum davidii</i>	Koreanspice viburnum C,F
<i>Viburnum japonicum</i>	David viburnum F
<i>Viburnum judd</i> (V X Judi)	Viburnum F
<i>Viburnum opulus sterile</i>	Viburnum C,F
<i>Viburnum plicatum tomentosum</i>	Common snowball viburnum F
<i>Viburnum setigerum</i>	Doublefile viburnum F
<i>Viburnum suspensum</i>	Tea viburnum F
<i>Viburnum tinus</i>	Virburnum, Sandankwa F
<i>Viburnum tinus compactum</i>	Viburnum, Laurustinus C,F
<i>Viburnum trilobum compactum</i>	Compactum-spring bouquet viburnum F
<i>Viburnum x pragense</i>	Spring bouquet viburnum F
<i>Weigela florida</i>	Dwarf cranberry bush F
<i>Xylosma congestum</i>	Viburnum F
<i>Yucca elata</i>	Bristol ruby weigela F
<i>Yucca recurvifolia</i>	Java red weigela F
	Minuet weigela F
	Weigela, oldfashioned F
	Xylosma F
	Yucca, soaptree C,F
	Yucca, pendulous F

Groundcovers/Perennials

Recommended Treatment Method

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Agapanthus africanus</i>	Lily-of-the-Nile	C,F
<i>Ajuga</i> spp.	Carpet bugle	F
<i>Arctotheca calendula</i>	Cape weed	F
<i>Asparagus retrofractus</i>	(No common name)	C,F
<i>Asparagus varieegata</i>	Tree fern	C,F
<i>Aster novae-angliae</i>	New England aster	C,F
<i>Aster novi-belgii</i>	New York aster	C,F
<i>Athyrium nipponicum</i>	Japanese painter fern	C,F
<i>Brassica oleracea</i>	Wild cabbage	C,F
<i>Callistepheus chinensis</i>	China aster	C,F
<i>Campanula elatines</i>	Bellflower	C,F
<i>Carpobrotus edulis</i>	Ice plant, largeleaf (see label)	F
<i>Clytostoma callistegioides</i>	Trumpet vine, violet	C,F
<i>Cortaderia selloana</i>	Pampas grass	F
<i>Cuphea hyssopifolia</i>	False Mexican heather	C,F
<i>Delosperma alba</i>	White iceplant	F
<i>Dietes vegeta</i>	Fortnight lily	C,F
<i>Digitalis mertonensis</i>	Foxglove	C,F
<i>Doronicum cordatum</i>	Leopard's bane	C,F
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant	F
<i>Erianthus ravennae</i>	Hardy pampus grass	C,F
<i>Festuca ovina glauca</i>	Blue fescue	F
<i>Gaillardia grandiflora</i>	Blanket flower	C,F
<i>Gazania rigens leucolaena</i>	Gazania, trailing	C,F
<i>Gazania</i> spp.	Gazania	F
<i>Hedera canariensis</i>	Ivy, Algerian	F
<i>Hedera helix</i>	Ivy, English	F
<i>Heliotropium fragrans</i>	Common heliotrope	C,F
<i>Hemerocallis</i> spp.	Daylily	C,F
<i>Hosta lancifolia</i>	Albo-marginata hosta	C,F

(continued)

Groundcovers/Perennials (cont.)

Recommended Treatment Method

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Hosta</i> spp.	Lily, plantain	C,F
<i>Heuchera micrantha</i>	Coral bells	C,F
<i>Hypericum</i> spp.	St. Johnswort	C,F
<i>Iberis sempervirens</i>	Evergreen candytuft	C,F
<i>Lampranthus spectabilis</i>	Trailing iceplant	F
<i>Leptospermum scaparium</i>	New Zealand teatree/Manuka	C,F
<i>Limonium perezii</i>	Statice/Sea lavender	C,F
<i>Liriope gigantea</i>	White lily turf	F
<i>Liriope muscari</i>	Lilac beauty lily turf	C,F
	Majestic lily turf	C,F
	Monroe white lily turf	C,F
	Silvery sunproof lily turf	C,F
	Variiegated liriope lily turf	C,F
	Big blue lily turf	C,F
<i>Lobelia erinus</i>	Edging lobelia	C,F
<i>Lonicera japonica</i>	Honeysuckle, Japanese	F
<i>Mesembryanthemum crystallinum</i>	Ice plant (see label)	F
<i>Monarda didyma</i>	Bee Balm	C,F
<i>Ophiopogon japonicus</i>	Mondo grass	F
<i>Osteospermum fruticosum</i>	Daisy, trailing African	F
<i>Pachysandra terminalis</i>	Japanese spurge	F
<i>Pennisetum setaceum</i>	Fountaingrass	C,F
<i>Polystichum polyblepharum</i>	Tassel fern	C,F
<i>Sedum brevifolium</i>	Stonecrop	C,F
<i>Sedum kamtschaticum</i>	Stonecrop	C,F
<i>Sedum spurium</i>	Stonecrop, tworow	C,F
<i>Tulbaghia violacea</i>	Society garlic	C,F
<i>Verbena rigida</i>	Veined verbena	C,F
<i>Veronica</i> spp.	Speedwell	C,F
<i>Vinca major</i>	Periwinkle, bigleaf	F
<i>Vinca minor</i>	Periwinkle, dwarf	F

Flowers

Recommended Treatment Method

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Achillea</i> spp.	Yarrow	C,F
<i>Antirrhinum majus</i>	Snapdragon	F
<i>Caladium bicolor</i>	Caladium, fancy leafed	F
<i>Chrysanthemum</i> spp.	Chrysanthemum	C,F
Mixed Hybrid	Dahlia	C,F
<i>Cladium bicolor</i>	Fancy-leafed caladium	F
<i>Coreopsis lanceolata</i>	Coreopsis	F
<i>Coreopsis verticulata</i>	Threadleaf coreopsis	C,F
<i>Dianthus barbatus</i>	Sweet William	F
<i>Dianthus gratianopolitanus</i>	Cheddar pink	C,F
<i>Dicentra spectabilis</i>	Bleeding heart	C,F
<i>Dimorphotheca</i> spp.	Marigold, cape	F
<i>Echinacea purpurea</i>	Coneflower, purple	C,F
<i>Evolvulus nuttallianus</i>	Blue daze	C,F
<i>Geum quellyon</i>	Geum	F
<i>Gladiolus hortulanus</i>	Gladiolus	F
<i>Gypsophila paniculata</i>	Baby's breath	F
<i>Impatiens wallerana</i>	Impatiens (Busy lizzie)	F
<i>Iris</i> spp.	Iris, bearded	F
<i>Liatis spicata</i>	Blazing star	C,F
<i>Pelargonium hortorum</i>	Geranium	F
<i>Petunia</i> spp.	Petunia	C,F
<i>Portulaca grandiflora</i>	Moss, rose	F
<i>Ranunculus asiaticus</i>	Ranunculus, Persian	F
<i>Rosa</i> spp.	Rose	F
<i>Rudbeckia fulgida</i>	Blackeyed susan	C,F
<i>Rudbeckia hirta</i>	Daisy, gloriosa (black-eyed Susan)	F
<i>Salvia</i> spp.	Salvia (Sage)	F
<i>Stokesia laevis</i>	Aster, stokes	F
<i>Strelitzia reginae</i>	Bird of paradise	F
<i>Tagetes</i> spp.	Marigold	F
<i>Viola wittrockiana</i>	Pansy	F
<i>Zinnia elegans</i>	Zinnia, common	F

Non-bearing Trees and Vines

Recommended Treatment Method

F = Field Grown

C = Container Grown

almond	F
apple	F
apricot	F
avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
kumquat	C,F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	C,F
peach	F
pear	F
pecan	C,F
pistachio	F
plum	F

(continued)

Non-bearing Trees and Vines (cont.)

Recommended Treatment Method

F = Field Grown

C = Container Grown

pomegranate
prune
raspberry
walnut, black
walnut, English

F
F
F
F
F

† Non-bearing plants are defined as those that will not bear fruit for at least one year after treatment.

Ornamental Bulbs

Surflan Pro may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth, and tulip. Apply Surflan Pro to the soil surface 2-4 weeks after planting, but prior to the emergence of annual weeds. For fall planted bulbs, apply Surflan Pro again in late winter or early spring to weed-free soil surfaces.

Broadcast Application Rates

Time of Application	Soil Texture	Surflan Pro		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
		(qt/acre)	(fl oz/1000 sq ft)		
Fall	Coarse	0.75	0.5	3	1.5
Fall	Medium and Fine	1.5	1.0	3	2.25
Feb. - March	All Soil Textures	0.75	0.5	3	2.25

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than $\frac{3}{4}$ inch.

Do not apply to gladioli corms prior to emergence or less than one (1) inch in diameter.

Shadehouse Areas

Surflan Pro may be applied to drainage areas under benches in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within 3 weeks prior to enclosure of greenhouse or poly-type structures.

Christmas Tree Plantations

Surflan Pro Alone

Apply Surflan Pro as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species, including fir (*Abies* spp.), pine (*Pinus* spp.), and spruce (*Picea* spp.). Follow all instructions provided in the "General Information" section of this label.

Broadcast Application Rates

Length of Control	Surflan Pro		Minimum Time Between Appls. (months)	Total Amount Allowed Per Year (qt/acre)
	(qt/acre)	(fl oz/ 1000 sq ft)		
2 - 4 months	2	1.5	2	8
4 - 8 months	4	3	2	8

Tank Mix Combinations

Tank mix combinations of Surflan Pro plus other labeled herbicides may be used as directed or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Surflan Pro Plus Glyphosate: Apply tank mix combinations of Surflan Pro plus glyphosate only as directed sprays in Christmas tree plantings. When applied according to use directions, Surflan Pro plus glyphosate will provide postemergence control of susceptible weed species listed on the label for glyphosate and residual preemergence control of susceptible weed species listed on the label for Surflan Pro. Refer to the label for glyphosate for specific use directions, precautions, and limitations before use.

Special Use Precautions:

Do not apply to Douglas-fir (*Pseudotsuga menziesii*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plants that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Noncropland Areas and Industrial Sites

Noncropland Areas - Tank Mix Combinations

Tank mix combinations of Surflan Pro plus glyphosate and many other labeled herbicides may be used to control undesirable vegetation in noncropland areas such as roadsides, rights-of-way, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Broadcast Application Rates

Length of Control	Surflan Pro		Minimum Time Between Appls. (months)	Total Amount Allowed Per Year (qt/acre)
	(qt/acre)	(fl oz/ 1000 sq ft)		
2 - 4 months	2	1.5	2	6
4 - 8 months	4	3	4	12
8 - 12 months	6	4.5	8	12

Industrial Sites - Tank Mix Combinations

Tank mix combinations of Surflan Pro plus glyphosate, Spike herbicide, and many other labeled herbicides may be used as overtop sprays to control existing vegetation on industrial sites such as utility substations, highway guard rails, sign posts, and delineators. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitation before use.

Warm Season Turfgrasses

Surflan Pro may be applied as a preemergence treatment for control of annual grasses and certain broadleaf weeds in established warm season turf including bahiagrass, bermuda-

grass, buffalograss, centipedegrass, St. Augustinegrass, zoysiagrass, and established tall fescue growing in warm season areas. Established turf is defined as a dense turf having a well-anchored root system and healthy, vigorous top growth. Use Surflan Pro only as a part of a total turf management program that includes good fertilization practices.

Surflan Pro may be tank mixed with Gallery herbicide (California registration pending) and applied preemergence to broaden the spectrum of broadleaf weed control in warm season turf. Refer to the label for Gallery for specific use directions, precautions, and limitations before use.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Surflan Pro.

Surflan Pro will not control emerged weeds. Successful preemergence control of weeds listed on this label requires that Surflan Pro be applied prior to weed germination and be activated by at least one-half (1/2) inch of rainfall or irrigation within 21 days of application.

Surflan Pro may injure turf that is not well established or is stressed or weakened due to unfavorable winter climatic conditions, drought, nematodes, or other factors which damage or weaken turf root systems. Apply Surflan Pro only to healthy, well-established turf that has a well-anchored root system.

Do not apply Surflan Pro in the spring or early summer to tall fescue turfgrass reseeded the previous fall. In such cases, apply Balan 2.5G granular herbicide at 60-80 pounds per acre in early summer (Round 1) and Surflan Pro at 1.5 quarts per acre approximately eight weeks later (Round 2). Do not apply Surflan Pro at the single application rate (2 quarts per acre) to established tall fescue; in such cases, apply 1.5 quarts per acre of Surflan Pro in an initial application, followed by a second application of 1.5 quarts per acre 8-10 weeks later.

In bermudagrass areas that have been overseeded with winter grasses, a spring application of Surflan Pro will thin the overseeded grasses.

Annual Grasses Controlled by Surflan Pro

Common Name	Summer Annuals:	Scientific Name
barnyardgrass (watergrass)		<i>Echinochloa crus-galli</i>
crabgrass, large		<i>Digitaria sanguinalis</i>
crabgrass, smooth		<i>Digitaria ischaemum</i>
crabgrass		<i>Digitaria spp.</i>

(continued)

Summer Annuals: (cont.)**Common Name**

crowfootgrass
 foxtail, bristleglass
 foxtail, giant
 foxtail, green
 (pigeongrass)
 foxtail, robust
 foxtail, yellow
 goosegrass
 (silver crabgrass)
 Johnsongrass
 (seedling only)
 ryegrass, Italian
 sandbur, field

Scientific Name

Dactyloctenium aegyptium
Setaria magna
Setaria faberi
Setaria viridis

Setaria robusta
Setaria glauca
Eleusine indica

Sorghum halepense

Lolium multiflorum
Cenchrus incertus

Common Name

bluegrass, annual

Winter Annuals:**Scientific Name**

Poa annua

Annual Broadleaf Weeds Controlled by Surflan Pro**Common Name**

carpetweed
 knotweed, prostrate
 purslane, common

Summer Annuals:**Scientific Name**

Mollugo verticillata
Polygonum aviculare
Portulaca oleracea

Common Name

chickweed, common
 henbit

Winter Annuals:**Scientific Name**

Stellaria media
Lamium amplexicaule

Broadleaf Weeds Suppressed by Surflan Pro**Common Name**

groundsel, common
 spurge, prostrate
 woodsorrel, yellow

Scientific Name

Senecio vulgaris
Euphorbia humistrata
Oxalis stricta

Application Rates, Frequency, and Timing of Application

Surflan Pro can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for annual bluegrass (*Poa annua*) and winter annual broadleaf weed control.

Broadcast Application Rates (Warm Season Turfgrasses)

Use Area	Surflan Pro		Minimum Time Between Appls. (months)	Total Amount Allowed Per Year (qt/acre)
	(qt/acre)	(fl oz/ 1000 sq ft)		
All, except	1.5	1	3	6
Florida	2	1.5	3	6
Florida	1.5	1	3	4.5

1. Summer Annual Grasses and Broadleaf Weeds

Single Application Program: Apply 2 quarts per acre of Surflan Pro in late winter or early spring, prior to the onset of conditions favorable for annual weed germination.

Split Application Program: As an alternative to a single application program, Surflan Pro may be applied in a split application. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. Apply 1.5 quarts per acre of Surflan Pro in an initial application, followed by a second application of 1.5 quarts per acre 8-10 weeks later.

The second treatment of the split application may follow application of a different pre-emergence grass herbicide in place of the initial application of Surflan Pro.

2. Annual Bluegrass (*Poa annua*) and Winter Annual Broadleaf Weeds

In areas of heavy annual bluegrass infestation, its elimination will result in temporary thinning of turfgrass cover. Proper fertilization, irrigation, and soil incorporated reseeded should be employed to speed the restoration of desirable turfgrass cover in areas previously occupied by annual bluegrass (see section on reseeding).

Apply Surflan Pro as a pre-emergence treatment in late summer or early fall, prior to the expected germination period for annual bluegrass and winter annual broadleaf weeds. If annual bluegrass infestation is severe and its elimination will result in thinning of turfgrass cover, apply Surflan Pro at 1.5 quarts per acre. If thinning of turfgrass cover is not a potential problem, Surflan Pro may be applied at 2 quarts per acre.

Weed Control in Florida

In Florida, apply 1.5 quarts per acre of Surflan Pro three times per year, or every 90-100 days, in the fall, early spring, and early summer. Do not apply more than 1.5 quarts per acre of Surflan Pro in any single application.

Application Equipment

Apply Surflan Pro evenly over the turfgrass area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. For best results, use application equipment designed to uniformly broadcast liquid herbicides. Calibrate application equipment prior to use, according to manufacturer's directions. Check equipment frequently to make sure it is working properly and distributing spray uniformly.

Reseeding

Herbicides that control annual weeds may also affect establishment of desirable turfgrass seedlings. Reseeding should be delayed for at least 90-120 days following application of Surflan Pro. When reseeding, it is essential that proper cultural practices such as soil cultivation and seedbed preparation, irrigation, and fertilization be followed. For satisfactory reseeding results following use of Surflan Pro, the seeding rate should be increased and equipment designed to place seed in full contact with soil (such as the Rogers Aero Seeder) should be employed.

Special Use Precautions:

To avoid possible injury, do not apply Surflan Pro to:

- Cool season turfgrass species.
- Golf course putting greens and tees or lawns containing dichondra or cool season turfgrass species.
- Newly sprigged or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfgrasses are well established and have well-anchored root systems.
- Newly hydromulched areas of bermudagrass until such areas are well established.
- Bermudagrass variety "Sun Turf" when tank mixed with atrazine.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer

Agrisel USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. AGRISEL USA, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Agrisel USA, Inc. or the seller. All such risks shall be assumed by buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Agrisel USA, Inc.'s election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Agrisel USA, Inc. shall not be liable for losses or damages resulting from handling or use of this product unless Agrisel USA, Inc. is promptly notified of such loss or damage in writing. In no case shall Agrisel USA, Inc. be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Agrisel USA, Inc. or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Surflan is a registered trademark of United Phosphorus, Inc.

EPA-Accepted 06-11-03

Rev. 11/04