



Armadillo Scram Professional repels armadillos and stops digging damage in two ways: Food tastes bad, and surrounding soil and armadillo burrows become bad smelling.

Assess and Quote


- Measure and calculate total square feet of application (width x length). Labor/application time is approx. 10 min. per 1000 sq. feet.
- 22 lb. container will provide protection of approx.
 1,160 linear feet at 1lb per 600 sq. feet.




Apply using a spreader at a rate of 1 pound per 600 square feet.

DAY 1: Apply with spreader. Cover the entire area where armadillo damage has occurred and up to boundaries where armadillos have entered. Proper spreader settings are in Application Instructions within each bucket. Extend the coverage to about thirty (30) feet around & beyond the damaged area within large properties. **Hand sprinkle,** using the scoop provided, within damaged garden plots and beds. Look behind shrubs and along buildings for excavated nesting areas and apply liberally within them. If burrow entrances can be located, drop a half scoop into each hole and flush down thoroughly with water.

- DAY 14: Reapply Armadillo Scram in 14 days after the initial application. Reapply every 30 days thereafter. Heavy or extended periods of rainfall will shorten the effective period.
- Armadillos do not hibernate and are active all year throughout their range. Use Armadillo Scram Proyear round where damage or an infestation occurs.





Recommended Spreader





