

Specimen Label

| | | | |
|-----------|-------|---|-----------|
| TRICLOPYR | GROUP | 4 | HERBICIDE |
|-----------|-------|---|-----------|



Garlon® 4 Ultra

HERBICIDE

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

For the control of woody plants and vines, and annual and perennial broadleaf weeds on:

- forest sites;
- non cropland areas including: electrical power and utility rights-of-way, industrial sites, non-irrigation ditch banks, pipelines, railroads, roadsides; and
- natural areas and wildlife habitat and management areas;
- including grazed areas on all of these listed sites.

Active Ingredient:

| | |
|--|---------|
| triclopyr: 2-[(3,5,6-trichloro-2-pyridinyl)oxy] acetic acid, butoxyethyl ester | 60.45% |
| Other Ingredients..... | 39.55% |
| Total..... | 100.00% |

Acid equivalent: triclopyr – 43.46% - 4 lb/gal

Precautionary Statements

Hazards to Humans and Domestic Animals

EPA Reg. No. 62719-527

Keep Out of Reach of Children

CAUTION

Causes Moderate Eye Irritation • Harmful If Swallowed • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers who handle this pesticide must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, or viton ≥14 mils
- Shoes plus socks

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, or viton ≥14 mils
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter or allow others to enter the treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store above 28°F or agitate before use.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

Storage and Disposal (Cont.)

Nonrefillable containers 5 gallons or less:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Refillable containers 5 gallons or larger:

Container Handling: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Nonrefillable containers 5 gallons or larger:

Container Handling: Nonrefillable container. Do not reuse or refill this container.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Product Information

Garlon 4 Ultra herbicide is recommended for the control of woody plants and vines, and herbaceous broadleaf weeds on forest sites, conifer plantations, non-cropland areas, including airports, barrow ditches, communication transmission lines, electrical power and utility rights-of-way, fencerows, gravel pits, industrial sites, military lands, mining and drilling areas, non-irrigation ditch banks, oil and gas pads, parking lots, petroleum tank farms, pipelines, railroads, roadsides, storage areas, storm water retention areas, substations, unimproved rough turf grasses, vacant lots and other non-crop residential areas; and natural areas (open space) for example campgrounds, parks, prairie management, trials and trialheads, recreation areas, wildlife openings, and wildlife habitat and management areas including grazed area on all these listed sites.

Use Precautions

When applying this product in tank mix combination, follow all applicable use directions and precautions on each manufacturer's label. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Sprays applied directly to Christmas trees may result in conifer injury. When treating unwanted vegetation in Christmas tree plantations, care should be taken to direct sprays away from conifers.

Garlon 4 Ultra is formulated as a low volatile ester. However, the combination of spray contact with impervious surfaces, such as roads and rocks, and increasing ambient air temperatures, may result in an increase in the volatility potential for this herbicide, increasing a risk for off-target injury to sensitive crops such as grapes and tomatoes.

Use Restrictions

Chemigation: Do not apply this product through any type of irrigation system.

Do not apply Garlon 4 Ultra directly to, or otherwise permit it to come into direct contact with cotton, grapes, peanuts, soybeans, tobacco, vegetable crops, flowers, citrus, or other desirable broadleaf plants. Do not permit spray mists containing it to drift onto such plants.

It is permissible to treat non-irrigation ditch banks, seasonally dry wetlands (such as flood plains, deltas, marshes, swamps, or bogs) and transitional areas between upland and lowland sites where surface water is not present except in isolated pockets due to uneven or unlevel conditions. Do not apply to open water (such as lakes, reservoirs, rivers, streams, creeks, salt water bays, or estuaries).

Do not apply on ditches that are used to transport irrigation water. Do not apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result.

Do not apply this product using mist blowers.

Maximum Use Rates

- Apply no more than 2 lb ae of triclopyr (2 quarts of Garlon 4 Ultra) per acre per growing season on range and pasture sites, or any area where grazing or harvesting hay is allowed.
- On forestry sites, apply no more than 6 lb ae of triclopyr (6 quarts of Garlon 4 Ultra) per acre per year.
- For all use sites other than range, pasture, forestry sites, and grazed /hayed areas, apply no more than 8 lb ae triclopyr (8 quarts of Garlon 4 Ultra) per acre per year.
- See Table 1 below for relationship between mixing rate, spray volume, and maximum application rate.

Grazing

- There are no grazing restrictions for livestock or dairy animals on treated areas
- Portions of grazed areas that intersect treated non-cropland and rights-of-way sites may be treated at up to 8 lb ae per acre if the area to be treated on the day of application comprises no more than 10% of the total grazable area.

Haying (harvesting of dried forage)

- Do not harvest hay for 14 days after application.

Slaughter Restriction: During the season of application, withdraw livestock from grazing treated grass at least 3 days before slaughter.

Avoid Injurious Spray Drift

Make applications only when there is little or no hazard from spray drift. Small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants that are near enough to be injured. It is suggested that a continuous smoke column at or near the spray site or a smoke generator on the spray equipment be used to detect air movement, lapse conditions, or temperature inversions (stable air). If the smoke layers or indicates a potential of hazardous spray drift, do not spray.

Aerial Application (Helicopter Only): For aerial application on rights-of-way or other areas near susceptible crops, apply through a Microfoil¹ or Thru-Valve¹ boom, or other drift control application equipment and/or use an agriculturally labeled drift control additive. If a spray thickening agent is used, follow all use recommendations and precautions on the product label. Spray only when the wind velocity is low (follow state regulations). Avoid application during air inversions.

¹ Reference within this label to a particular piece of equipment produced by or available from other parties is provided without consideration for use by the reader at its discretion and subject to the reader's independent circumstances, evaluation, and expertise. Such reference by Dow AgroSciences is not intended as an endorsement of such equipment, shall not constitute a warranty (express or implied) of such equipment, and is not intended to imply that other equipment is not available and equally suitable. Any discussion of methods of use of such equipment does not imply that the reader should use the equipment other than as advised in directions available from the equipment's manufacturer. The reader is responsible for exercising its own judgment and expertise, or consulting with sources other than Dow AgroSciences, in selecting and determining how to use its equipment.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications:

- The distance of the outer most operating nozzles on the boom must not exceed 3/4 the length of the rotor.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory. [This information is advisory in nature and does not supersede mandatory label requirements.]

Aerial Drift Reduction Advisory

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than 3/4 of the rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a local, low level temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of the smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Ground Equipment: To aid in reducing spray drift potential when making ground applications near susceptible crops or other desirable broadleaf plants, Garlon 4 Ultra should be used in thickened (high viscosity) spray mixtures using an agriculturally labeled drift control additive, high viscosity invert system, or equivalent as directed by the manufacturer. When using a spray thickening or inverting additive, follow all use directions and precautions on the product label. With ground equipment, spray drift can be reduced by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by keeping the operating spray pressures at the lower end of the manufacturer's recommended pressures for the specific nozzle type used (low pressure nozzles are available from spray equipment manufacturers); and by spraying when wind velocity is low. Do not apply with nozzles that produce a fine droplet spray. Select nozzles and pressures which provide adequate plant coverage, but minimize the production of fine spray particles.

High Volume Leaf-Stem Treatment: To minimize spray drift, keep sprays no higher than brush tops and keep spray pressures low enough to provide coarse spray droplets. A agriculturally labeled thickening agent may be used to reduce spray drift.

Mixing Directions for all use sites

Garlon 4 Ultra may be foliarly applied by diluting with water or by preparing an oil-water emulsion. For woody plant control, an oil-water emulsion performs more dependably under a broader range of conditions than a straight water dilution and is recommended for aerial applications.

Oil-Water Mixture Sprays

Prepare a premix of oil, surfactant and Garlon 4 Ultra in a separate container using diesel fuel, fuel oil, or kerosene plus an emulsifier such as Sponto 712 or Triton X-100. Use a jar test to check spray mix compatibility before preparing oil-water emulsion sprays in the mixing tank. Do not allow any water or mixtures containing water to get into the premix or Garlon 4 Ultra since a thick "invert" (water in oil) emulsion may form that will be difficult to break. Such an emulsion may also be formed if the premix of Garlon 4 Ultra is put into the mixing tank before the addition of water. Fill the spray tank about one-half full with water, then slowly add the premix with continuous agitation and complete filling the tank with water. Continue moderate agitation.

Oil Mixture Sprays for Basal Treatment

Prepare oil-based spray mixtures using either a commercially available basal oil, kerosene diesel fuel, or No. 1 or No. 2 fuel oil. Substitute other oils or diluents only as recommended by the oil or diluent's manufacturer. When mixing an oil mixture, read and follow the use directions and precautions on the manufacturer's product label. Add Garlon 4 Ultra to the required amount of oil in the spray tank or mixing tank and mix thoroughly. If the mixture stands over four hours, reagitation is required.

Oil Mixtures of Garlon 4 Ultra and Tordon 22K: Tordon 22K and Garlon 4 Ultra may be used in tank mix combination for basal bark treatment of woody plants. These herbicides are incompatible and will not form a stable mixture when mixed together directly in oil. Make a stable tank mixture for basal bark application by first combining each product with a compatibility agent prior to final mixing in the desired ratio. Tordon 22K is not registered for use in the states of California and Florida.

Herbicide Resistance Management

Triclopyr, the active ingredient in this product, is a Group 4 herbicide based on the mode of action classification system of the Weed Science Society of America.

Any weed population may contain or develop plants resistant to Group 4 herbicides. Resistant weeds may dominate the weed population if these herbicides are used repeatedly in the same field. Such resistant weed plants may not be effectively managed using Group 4 herbicides but may be effectively managed utilizing other herbicides alone or in mixtures from a different herbicide Groups that are labeled for control of these weeds and/or by using cultural or mechanical practices. However, a herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides.

Rotate the use of Garlon 4 Ultra or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use less the resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is less prone to resistance.

Adopt an integrated weed management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation that considers tillage (or other mechanical control methods), cultural, biological, and other management practices.

Scout after a herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide

from a different group or by mechanical method such as hoeing, mowing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

If a weed population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.

Contact your local extension specialist or certified crop advisor for additional pesticide resistance management or integrated weed management recommendations for specific use sites.

Plants Controlled by Garlon 4 Ultra

Woody Plants Controlled

| Common Name | Scientific Name | Life Cycle | Plant Family |
|------------------------|------------------------------------|------------|----------------|
| acacia, twisted | <i>Acacia tortuosa</i> | perennial | Fabaceae |
| alder | <i>Aldus spp.</i> | perennial | Betulaceae |
| arrowwood | <i>Viburnum ventanum</i> | perennial | Caprifoliaceae |
| ash | <i>Fraxinus spp.</i> | perennial | Oleaceae |
| aspen | <i>Populus tremuloides</i> | perennial | Salicaceae |
| bear clover (bearmat) | <i>Chamaebatia foliolosa</i> | perennial | Fabaceae |
| beech | <i>Fagus spp.</i> | perennial | Fagaceae |
| birch | <i>Betula spp.</i> | perennial | Betulaceae |
| blackberry | <i>Rubus spp.</i> | perennial | Rosaceae |
| blackbrush | <i>Acacia rigidula</i> | perennial | Fabaceae |
| blackgum | <i>Nyssa salvatica</i> | perennial | Cornaceae |
| boxelder (1) | <i>Acer negundo</i> | perennial | Aceraceae |
| Brazilian pepper | <i>Schinus terebinthifolius</i> | perennial | Anacardiaceae |
| buckthorn | <i>Rhamnus spp.</i> | perennial | Rhamnaceae |
| casacara | <i>Rhamnus pushiana</i> | perennial | Rhamnaceae |
| ceanothus | <i>Ceanothus spp.</i> | perennial | Rhamnaceae |
| cherry | <i>Prunus spp.</i> | perennial | Rosaceae |
| cherry, choke | <i>Prunus virginiana</i> | perennial | Rosaceae |
| chinquapin | <i>Quercus muhlenbergii</i> | perennial | Fagaceae |
| cottonwood | <i>Populus deltoides</i> | perennial | Salicaceae |
| crataegus (hawthorn) | <i>Crataegus spp.</i> | perennial | Rosaceae |
| creeper, Virginia (1) | <i>Parthenocissus quinquefolia</i> | perennial | Vitaceae |
| dogwood | <i>Cornus spp.</i> | perennial | Cornaceae |
| Douglas-fir | <i>Psuedotsuga menziesii</i> | perennial | Pinaceae |
| elderberry | <i>Sambucus Canadensis</i> | perennial | Caprifoliaceae |
| elm | <i>Ulmus, spp</i> | perennial | Ulmaceae |
| elm, winged | <i>Ulmus alata</i> | perennial | Ulmaceae |
| gallberry | <i>Ilex coriacea</i> | perennial | Aquifoliaceae |
| granjeno | <i>Celtis ehrenbergiana</i> | perennial | Ulmaceae |
| guajillo | <i>Acacia berlandieri</i> | perennial | Fabaceae |
| guava | <i>Psidium guajava</i> | perennial | Myrtaceae |
| gorse | <i>Ulex europaeus</i> | perennial | Fabaceae |
| hazel | <i>Corylus americana</i> | perennial | Betulaceae |
| hickory | <i>Carya spp.</i> | perennial | Juglandaceae |
| hornbeam | <i>Carpinus spp.</i> | perennial | Betulaceae |
| huisache (suppression) | <i>Acacia farnesiana</i> | perennial | Fabaceae |
| ivy, poison | <i>Toxicodendron radicans</i> | perennial | Anacardiaceae |
| kudzu | <i>Pueraria lobata</i> | perennial | Fabaceae |
| locust | <i>Robinia spp.</i> | perennial | Fabaceae |
| madrone | <i>Arbutus spp.</i> | perennial | Ericaceae |
| magnolia, sweetbay | <i>Magnolia virginiana</i> | perennial | Magnoliaceae |
| maples | <i>Acer spp.</i> | perennial | Aceraceae |
| maple, bigleaf (1) | <i>Acer macrophyllum</i> | perennial | Aceraceae |
| milkweed vine (1) | <i>Asclepias spp.</i> | perennial | Asclepiaceae |
| mulberry | <i>Morus spp.</i> | perennial | Moraceae |
| myrtle, wax | <i>Morella cerifera</i> | perennial | Myricaceae |
| oaks | <i>Quercus spp.</i> | perennial | Fagaceae |

Plants Controlled by Garlon 4 Ultra (Cont.)

Woody Plants Controlled

| Common Name | Scientific Name | Life Cycle | Plant Family |
|--------------------------|-------------------------------------|------------|----------------|
| oak, poison | <i>Toxicodendron diversilobum</i> | perennial | Anacardiaceae |
| Osage orange | <i>Maclura pomifera</i> | perennial | Moraceae |
| peppervine | <i>Ampelopsis arborea</i> | perennial | Vitaceae |
| persimmon | <i>Disospyros spp.</i> | perennial | Ebenaceae |
| pine | <i>Pinus spp.</i> | perennial | Pinaceae |
| poplar | <i>Populus spp.</i> | perennial | Salicaceae |
| poplar, tulip | <i>Liriodendron tulipifera</i> | perennial | Magnoliaceae |
| primrose, willow | <i>Ludwigia peruviana</i> | perennial | Onagraceae |
| rose, wild | <i>Rosa spp.</i> | perennial | Rosaceae |
| salmonberry | <i>Rubus spectabilis</i> | perennial | Rosaceae |
| saltbush (silver myrtle) | <i>Baccharis spp</i> | perennial | Asteraceae |
| saltcedar | <i>Tamarix spp.</i> | perennial | Tamariaceae |
| sassafras | <i>Sassafras spp.</i> | perennial | Lauraceae |
| scotchbroom | <i>Cytisus scoparius</i> | perennial | Fabaceae |
| sumac | <i>Rhus spp.</i> | perennial | Anacardiaceae |
| sweetgum | <i>Liquidamber styraciflura</i> | perennial | Hamamelidaceae |
| sycamore | <i>Platanus occidentalis</i> | perennial | Plantanaceae |
| tanoak | <i>Notholithocarpus densiflorus</i> | perennial | Fagaceae |
| tree of heaven | <i>Ailanthus altissima</i> | perennial | Simaroubaceae |
| trumpet creeper (1) | <i>Campsis radicans</i> | perennial | Bignoniaceae |
| willow | <i>Salix spp.</i> | perennial | Saliciaceae |

¹ For best control, use either a basal bark or cut stump treatment.

² For complete control, re-treatment may be necessary.

Annual and Perennial Broadleaf Weeds

| Common Name | Scientific Name | Life Cycle | Plant Family |
|------------------------------|---------------------------------|------------|----------------|
| beggarweed, creeping | <i>Desmodium incanum</i> | perennial | Fabaceae |
| bindweed, field (top growth) | <i>Convolvulus arvensis</i> | perennial | Convolvulaceae |
| burdock, common | <i>Arctium minus</i> | biennial | Asteraceae |
| carrot, wild | <i>Daucus carota</i> | biennial | Apiaceae |
| chicory | <i>Cichorium intybus</i> | perennial | Asteraceae |
| cinquefoil, sulfur (2) | <i>Potentilla recta</i> | perennial | Rosaceae |
| clover | <i>Trifolium spp.</i> | perennial | Fabaceae |
| dandelion (top growth) | <i>Taraxacum officinale</i> | perennial | Asteraceae |
| dock, curly | <i>Rumex crispus</i> | perennial | Polygonaceae |
| dogfennel | <i>Eupatorium capillifolium</i> | perennial | Asteraceae |
| goldenrod | <i>Solidago spp.</i> | perennial | Asteraceae |
| ivy, ground | <i>Glechoma hederacea</i> | perennial | Lamiaceae |
| kudzu | <i>Pueraria montana</i> | perennial | Fabaceae |
| lambsquarters | <i>Chenopodium spp.</i> | annual | Chenopodiaceae |
| lespedeza, annual | <i>Lespedeza striata</i> | annual | Fabaceae |
| lespedeza, Sericea (1) | <i>Lespedeza cuneata</i> | perennial | Fabaceae |
| lettuce, prickly | <i>Lactuca serriola</i> | annual | Asteraceae |
| matchweed | <i>Lippia nodiflora</i> | perennial | Verbanaceae |
| medic, black | <i>Medicago lupulina</i> | perennial | Fabaceae |
| mustard | <i>Brassica spp.</i> | annual | Brassicaceae |
| mustard, garlic (4) | <i>Alliaria petiolata</i> | biennial | Brassicaceae |
| plantain | <i>Plantago spp.</i> | annual | Plantaginaceae |
| ragweed, common | <i>Ambrosia artemisiifolia</i> | annual | Asteraceae |
| ragweed, western | <i>Ambrosia psilostachya</i> | perennial | Asteraceae |
| smartweed, Pennsylvania | <i>Polygonum pennsylvanicum</i> | annual | Polygonaceae |
| soda apple, tropical (3) | <i>Solanum viarum</i> | perennial | Solanaceae |
| thistle, bull | <i>Cirsium vulgare</i> | biennial | Asteraceae |
| thistle, Canada | <i>Cirsium arvense</i> | perennial | Asteraceae |
| Vetch | <i>Vicia spp.</i> | perennial | Fabaceae |
| violet, wild | <i>Viola papilionacea</i> | perennial | Violaceae |
| yarrow, common | <i>Achillea millefolium</i> | perennial | Asteraceae |

- Sericea lespedeza:** Apply 1 to 2 pints of Garlon 4 Ultra per acre. For best results, apply after maximum foliage development in the late spring to early summer, but prior to bloom.
- Sulfur cinquefoil:** Apply 1 to 2 pints of Garlon 4 Ultra per acre. For best results, apply to plants in the rosette stage.
- Tropical soda apple:** Apply 2 pints of Garlon 4 Ultra per acre when tropical soda apple plants reach the first flower stage. For best results, apply in a total spray volume of 40 gallons per acre using ground equipment. An agricultural surfactant may be added at the manufacturer's recommended rate to provide more complete wetting and coverage of the foliage. Spot treatments may be used to control sparse plant stands. For spot treatment use a 1 to 1.5% solution of Garlon 4 Ultra in water (1 to 1 1/2 gallons of Garlon 4 Ultra in 100 gallons total spray mixture) and spray the entire plant to completely wet the foliage. **In Florida**, control of tropical soda apple may be improved by using the following management practices:
 - Mow plants to a height of 3 inches every 50 to 60 days or whenever they reach flowering. Continue the mowing operation through April.
 - In late May to June (50 to 60 days after the April mowing), apply Garlon 4 Ultra as a broadcast treatment.
 - Use spot treatment to control any remaining plants or thin stands of plants that germinate following a broadcast treatment
- Garlic mustard:** apply as a 1.25 to 2.5% v/v foliar spray-to-wet application

Use Information

Use Garlon 4 Ultra at rates of 1 to 8 quarts per acre to control broadleaf weeds and woody plants. It is suggested that rates higher in this rate range be used to control woody plants. In all cases, use the amount specified in enough water to give uniform and complete coverage of the plants to be controlled. The order of addition to the spray tank is water, spray thickening agent (if used), surfactant (if used), additional herbicide (if used), and Garlon 4 Ultra. If a standard agricultural surfactant is used, use at a rate of 1 to 2 quarts per acre. Use continuous adequate agitation.

Before using any recommended tank mixtures, read the directions and all precautions on both labels.

For best results apply when woody plants and weeds are actively growing. When hard to control species such as ash, blackgum, choke cherry, elm, maples (other than vine or big leaf), oaks, pines, or winged elm are prevalent, during applications made during late summer when the plants are mature, or during drought conditions, use the higher rates of Garlon 4 Ultra alone or in combination with Graslan L or Tordon 22K herbicide. Graslan L and Tordon 22K are restricted use pesticides. Graslan L and Tordon 22K are not registered for use in the states of California and Florida.

When using Garlon 4 Ultra in combination with Freelexx or a 2,4-D low volatile ester herbicide, generally the higher rates of Garlon 4 Ultra should be used for satisfactory brush control.

Use the higher dosage rates when brush approaches an average of 15 feet in height or when the brush covers more than 60% of the area to be treated. If lower rates are used on hard to control species, resprouting may occur the year following treatment.

On sites where easy to control brush species dominate, rates less than those listed may be effective. Consult state or local extension personnel for such information.

Foliage Treatment With Ground Equipment

High Volume Foliage Treatment

For control of woody plants, use Garlon 4 Ultra at the rate of 2 to 6 quarts per 100 gallons of spray mixture, or Garlon 4 Ultra at 2 to 4 quarts may be tank mixed with labeled rates of Freelexx or a 2,4-D low volatile ester herbicide, Graslan L, or Tordon 22K and diluted to make 100 gallons of spray. Do not apply more than 2 gallons of Garlon 4 Ultra per acre. Apply at a volume of 100 to 400 gallons of total spray per acre depending upon size and density of woody plants. Graslan L and Tordon 22K are not registered for use in the states of California and Florida. When tank mixing, follow applicable use directions and precautions on each manufacturer's label.

Depending upon the size and density of the woody plants, apply sufficient spray volume to thoroughly wet all leaves, stems, and root collars. To minimize spray drift, select the minimum spray pressure that provides adequate plant coverage without forming a mist and direct sprays no higher than the top of the target plants. Use a drift control additive cleared for application to growing crops to reduce spray drift. Before using any tank mixture, read the directions and use precautions on both labels. For best results, apply when woody plants and weeds are actively growing.

Table 1: The following table is provided as a guide to the user to achieve the proper rate of Garlon 4 Ultra on forestry and non-cropland sites.

| Total Spray Volume (gallons/acre) | Rate of Garlon 4 Ultra | |
|-----------------------------------|---|---|
| | Forestry Sites (qt/100 gallons of spray) ¹ | Non-Cropland Sites (qt/100 gallons of spray) ² |
| 400 | 1.5 | 2 |
| 300 | 2 | 2.7 |
| 200 | 3 | 4 |
| 100 | 6 | 8 |
| 50 | 12 | 16 |
| 40 | 15 | 20 |
| 30 | 20 | 26.7 |
| 20 | 30 | 40 |
| 10 | 60 | 80 |

¹ Do not exceed the maximum use rate of 6 quarts of Garlon 4 Ultra (6 lb ae of triclopyr) per acre per year.

² Do not exceed the maximum use rate of 8 quarts of Garlon 4 Ultra (8 lb ae of triclopyr) per acre per year for non-grazable areas, or 2 quarts (2 lb ae of triclopyr) per acre per year for grazed areas, except on portions of grazed areas that meet the following requirement. Portions of grazed areas that intersect treated non-cropland, rights-of-way and forestry sites may be treated at up to 8 lb ae per acre if the area to be treated on the day of application comprises no more than 10% of the total grazable area.

Low Volume Foliar Treatment

To control susceptible woody plants, mix up to 5% v/v of Garlon 4 Ultra in water and apply 10 to 100 gallons of finished spray. The spray concentration of Garlon 4 Ultra and total spray volume per acre should be adjusted according to the size and density of target woody plants and kind of spray equipment used. With low volume sprays, use sufficient spray volume to obtain uniform coverage of target plants including the surfaces of all foliage, stems, and root collars (see Use Precautions and Restrictions). For best results, a surfactant should be added to all spray mixtures. Match equipment and delivery rate of spray nozzles to height and density of woody plants. When treating tall, dense brush, a truck mounted spray gun with spray tips that deliver up to 2 gallons per minute at 40 to 60 psi may be required. Backpack or other types of specialized spray equipment with spray tips that deliver less than 1 gallon of spray per minute may be appropriate for short, low to moderate density brush.

See Table 1 for relationship between mixing rate, spray volume and maximum application rate.

Tank Mixing: As a low volume foliar spray, up to 9 quarts of Garlon 4 Ultra may be applied in tank mix combination with labeled rates of Tordon 22K or Graslan L in 10 to 100 gallons of finished spray. Graslan L and Tordon 22K are not registered for use in the states of California and Florida.

Broadcast Applications With Ground Equipment

Apply Garlon 4 Ultra using equipment that will assure thorough and uniform coverage at spray volumes applied. See Table 1 for relationship between mixing rate, spray volume and maximum application rate.

Woody Plant Control

Foliage Treatment: Use 4 to 8 quarts of Garlon 4 Ultra in enough water to make 5 gallons or more per acre of total spray, or 1 1/2 to 3 quarts of Garlon 4 Ultra may be combined with labeled rates of Freelexx or a 2,4-D low volatile ester, Graslan L, or Tordon 22K in sufficient water to make 5 gallons or more per acre of total spray. Graslan L and Tordon 22K are not registered for use in the states of California and Florida.

Broadleaf Weed Control

Use Garlon 4 Ultra at rates of 1 to 4 quarts in a total volume of 5 gallons or more per acre as a water spray mixture. Apply anytime weeds are actively growing. Garlon 4 Ultra at 0.25 to 3 quarts may be tank mixed with labeled rates of Freelexx or a 2,4-D amine or low volatile ester, Tordon 22K, or Graslan L to improve the spectrum of activity. For thickened (high viscosity) spray mixtures, Garlon 4 Ultra can be mixed with diesel oil or other inverting agent. When using an inverting agent, read and follow the use directions and precautions on the product label. Graslan L and Tordon 22K are not registered for use in the states of California and Florida.

Aerial Application (Helicopter Only)

Aerial sprays should be applied using suitable drift control (see Use Precautions and Restrictions).

Foliage Treatment (Utility and Pipeline Rights-of-Way)

Use 4 to 8 quarts of Garlon 4 Ultra alone, or 3 to 4 quarts of Garlon 4 Ultra in a tank mix combination with labeled rates of Freelexx or a 2,4-D low volatile ester, Graslan L or Tordon 22K and apply in a total spray volume of 10 to 30 gallons per acre. Use the higher rates and volumes when plants are dense or under drought conditions. Graslan L and Tordon 22K are not registered for use in the states of California and Florida.

Portions of grazed areas that intersect treated non-cropland, rights-of-way and forestry sites may be treated at up to 8 lb ae per acre if the area to be treated on the day of application comprises no more than 10% of the total grazable area.

Basal Bark, Dormant Stem and Cut Surface Treatments for use on all sites

Individual plant treatments such as basal bark and cut surface applications may be used on any use site listed on this label at a maximum use rate of 8 quarts of Garlon 4 Ultra (8 lb ae of triclopyr) per acre. These types of applications are made directly to ungrazed parts of plants and, therefore, are not restricted by the grazing maximum rate of 2 quarts of Garlon 4 Ultra (2 lb ae of triclopyr) per acre.

Conventional Basal Bark Treatment

To control susceptible woody plants with stems less than 6 inches in basal diameter, mix 1 to 5 gallons of Garlon 4 Ultra in enough oil to make 100 gallons of spray mixture. Apply with backpack sprayer or power spraying equipment using low pressure (20 to 40 psi). Spray the basal parts of brush and tree trunks to a height of 12 to 15 inches from the ground, thoroughly wetting the indicated area. Spray until runoff at the ground line is noticeable. Old or rough bark requires more spray than smooth young bark. Apply anytime, including the winter months, except when snow or water prevent spraying to the ground line. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Low Volume Basal Bark Treatment

To control susceptible woody plants with stems less than 6 inches in basal diameter, mix 20 to 30 gallons of Garlon 4 Ultra in enough oil to make 100 gallons of spray mixture. Apply with a backpack or sprayer using low pressure and a solid cone or flat fan nozzle. Spray the basal parts of brush and tree trunks to a height of 12 to 15 inches from the ground in a manner that thoroughly wets the lower stems, including the root collar area, but not to the point of runoff. Herbicide concentration should vary with size and susceptibility of species treated. Treatments may be applied throughout the year including when snow is present. Efficacy may be reduced when stem surfaces are saturated with water. See Table 1 for relationship between mixing rate, spray volume and maximum application rate. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Garlon 4 Ultra Plus Milestone for basal bark applications

Mix Garlon 4 Ultra with Milestone in a commercially available basal diluent (or other oils or basal diluents as recommended by the manufacturer); the basal oil should be compatible with a water soluble herbicide such as Milestone. Make a stable tank mixture for basal bark application by first combining each product with a compatibility agent prior to final mixing in the desired ratio. If using a tank mix, mix the oil-based products such as Garlon 4 Ultra thoroughly with basal oil and add any other oil-based products before adding the water based products. If the mixture stands for more than 30 minutes, reagitation may be required. Oil and water based mixtures can separate over time. Long-term storage is not recommended without vigorous agitation prior to use or without a recommended compatibility agent.

Garlon 4 Ultra Plus Tordon 22K in Oil Tank Mix: Garlon 4 Ultra and Tordon 22K may be used in tank mix combination as a low volume basal bark treatment to improve control of certain woody species such as ash, elm, maple, poplar, aspen, hackberry, oak, oceanspray, birch, hickory, pine, tanoak, cherry, locust, sassafras, and multiflora rose. (See product bulletin for mixing instructions.) Tordon 22K is not registered for use in the states of California and Florida.

Streamline Basal Bark Treatment (Southern States)

To control or suppress susceptible woody plants for conifer release, mix 20 to 30 gallons of Garlon 4 Ultra in enough oil to make 100 gallons of spray mixture. Streamline basal bark treatments are most effective on stems less than 4 inches in basal diameter. Apply with a backpack sprayer or using equipment that provides a directed straight stream spray. Apply the spray in a 2- to 3-inch wide band to one side of stems less than 3 inches in basal diameter. When the optimum amount of spray mixture is applied, the treated zone should widen to encircle the stem within approximately 30 minutes. Treat both sides of stems which are 3 to 4 inches in basal diameter. Direct the spray at bark that is approximately 12 to 24 inches above ground. Pines (loblolly, slash, shortleaf, and Virginia) up to 2 inches in diameter breast height (dbh) can be controlled by directing the spray at a point approximately 4 feet above ground. Vary spray mixture concentration with size and susceptibility of the species being treated. Better control is achieved when spray is applied to thin juvenile bark and above rough thickened mature bark. This technique is not recommended for scrub and live oak species, including blackjack, turkey, post, live, bluejack and laurel oaks, or bigleaf maple. Apply anytime, including winter months, except when snow or water prevents spraying at the desired height above ground level. **Note:** Best

results with some hardwood species occur when applications are made from approximately 6 weeks prior to leaf expansion in the spring until approximately 2 months after leaf expansion is completed. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Low Volume Stem Bark Band Treatment (North Central and Lake States)

The treatment band may be positioned at any height up to the first major branch. For best results apply the band as low as possible. Spray mixture concentration should vary with size and susceptibility of species to be treated. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Thinline Basal Bark Treatment

To control susceptible woody plants with stems less than 6 inches in diameter, apply Garlon 4 Ultra, either undiluted or mixed at 50 to 75% v/v with oil, in a thin stream to all sides of the lower stems. The stream should be directed horizontally to apply a narrow band of Garlon 4 Ultra around each stem or clump. Use a minimum of 2 to 15 milliliters of Garlon 4 Ultra or oil mixture with Garlon 4 Ultra to treat single stems and from 25 to 100 milliliters to treat clumps of stems. Use an applicator metered or calibrated to deliver the small amounts required. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Dormant Stem Treatment

Dormant stem treatments will control susceptible woody plants and vines with stems less than 2 inches in diameter. Plants with stems greater than 2 inches in diameter may not be controlled and resprouting may occur. This treatment method is best suited for sites with dense, small diameter brush. Dormant stem treatments of Garlon 4 Ultra can also be used as a chemical side-trim for controlling lateral branches of larger trees that encroach onto roadside, utility, or other rights-of-way.

High volume and low volume applications using backpacks deliver approximately the same amount of herbicide per acre but differ in delivery volumes to achieve that rate.

High Volume Applications

Mix 4 to 8 quarts of Garlon 4 Ultra in 2 to 3 gallons of crop oil concentrate or other recommended oil and add this mixture to enough water to make 100 gallons of spray solution. Use continuous adequate agitation. Apply using low pressure (20 to 40 psi). In western states, apply anytime after woody plants are dormant and most of the foliage has dropped. In other areas apply anytime within 10 weeks of budbreak, generally February through April. Garlon 4 Ultra may be mixed with 4 quarts of Weedone 170 herbicide to improve the control of black cherry and broaden the spectrum of herbicidal activity. Do not apply to wet or saturated bark as poor control may result.

Low Volume Applications

Mix Garlon 4 Ultra at 4 to 6 gallons and 2 to 3 gallons of crop oil concentrate or other recommended oil and add this mixture to enough water to make 100 gallons of spray solution. Use continuous adequate agitation. Apply with backpack or other low volume spraying equipment, using low pressure (20 to 40 psi). Garlon 4 Ultra may be mixed with other herbicides to broaden the spectrum of herbicidal activity. Do not apply to wet or saturated bark as poor control may result.

Cut Surface

Cut surface applications with Garlon 4 Ultra can be made anytime after cutting up to re-sprouting. After re-sprouting basal bark or foliar applications are more suitable.

Basal Cut Stump Treatment

To control resprouting, mix 20 to 30 gallons of Garlon 4 Ultra in enough oil to make 100 gallons of spray mixture. Apply with a backpack or sprayer using low pressures and a solid cone or flat fan nozzle. Spray the root collar area and any exposed roots of root suckering species, sides of the stump, and the outer portion of the cut surface, including the cambium, until thoroughly wet, but not to the point of runoff. Spray mixture concentration should vary with size and susceptibility of species treated, using the higher rate for larger stumps, stumps with thicker bark or harder to control plants. Apply anytime, including in winter months, except when snow or water prevent spraying to the ground line. **Mixing with oil requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Cut Stump Treatment

To control resprouting of difficult to control species like saltcedar and other *Tamarix* species, bigleaf maple, tanoak, Oregon myrtle, and other susceptible species, apply Garlon 4 Ultra as a 50% dilution v/v in water by spraying all the exposed cambium layer on the freshly cut surface, or use undiluted Garlon 4 Ultra immediately after cutting. Use of undiluted Garlon 4 Ultra is most effective for hard-to-control species. Treatments may be applied throughout the year; however, control may be reduced with treatment during periods of moisture stress as in late summer or

early spring sap flow. Cut stumps so that they are approximately level to facilitate uniform coverage of Garlon 4 Ultra. Use an applicator that can be calibrated to deliver the small amounts of material required.

Forest Management Applications

All application methods described on this label may be used on forest management sites.

For broadcast applications, apply 1 to 6 quarts of Garlon 4 Ultra per acre in a total spray volume of 5 to 25 gallons per acre by air or 10 to 100 gallons per acre by ground. Use spray volumes sufficient to provide thorough coverage of treated foliage. Nozzles or additives that produce larger droplets of spray may require higher spray volumes to provide adequate coverage.

Plant Back Interval for Conifers: Conifers planted sooner than one month after treatment with Garlon 4 Ultra at less than 4 quarts per acre or sooner than two months after treatment at 4 to 6 quarts per acre may be injured. When tank mixtures of herbicides are used for forest site preparation, labels for all products in the mixture should be consulted and the longest recommended waiting period observed.

Forest Site Preparation (Not For Conifer Release)

Southern States Including Alabama, Arkansas, Delaware, Florida, Georgia, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia: To control susceptible woody plants and broadleaf weeds, apply Garlon 4 Ultra at a rate of 4 to 6 quarts per acre. To broaden the spectrum of woody plants and broadleaf weeds controlled, apply 2 to 4 quarts of Garlon 4 Ultra per acre in tank mix combination with labeled rates of Graslan L or Tordon 22K. Graslan L and Tordon 22K are not registered for use in the state of Florida. Where grass control is also desired, Garlon 4 Ultra, alone or in combination with Tordon 22K or Graslan L, may be applied with labeled rates of other herbicides registered for grass control in forests. Use of tank mix products must be in accordance with the most restrictive of label limitations and precautions. Do not exceed labeled application rates. Garlon 4 Ultra cannot be tank mixed with any product containing a label prohibition against such mixing.

In Western, Northeastern, North Central, and Lake States (States Not Listed Above as Southern States): To control susceptible woody plants and broadleaf weeds, apply Garlon 4 Ultra at a rate of 3 to 6 quarts per acre. To broaden the spectrum of woody plants and broadleaf weeds controlled, apply 1.5 to 3 quarts per acre of Garlon 4 Ultra in tank mix combination with labeled rates of Graslan L, Tordon 22K, or Freelexx or a 2,4-D low volatile ester or Freelexx. Graslan L and Tordon 22K are not registered for use in the state of California. Where grass control is also desired, Garlon 4 Ultra, alone or in tank mix combination with Graslan L or Tordon 22K, may be applied with labeled rates of other herbicides registered for grass control in forests. When applying tank mixes, follow applicable use directions and precautions on each product label.

Southern Coastal Flatwoods: To control susceptible broadleaf weeds and woody species such as gallberry and wax-myrtle, and for partial control of saw-palmetto, apply 2 to 4 quarts of Garlon 4 Ultra per acre. To broaden the spectrum of species controlled to include fetterbush, staggerbush, titi, and grasses, apply 2 to 3 quarts per acre of Garlon 4 Ultra in tank mix combination with labeled rates of Arsenal Applicator's Concentrate herbicide. Where control of gallberry, wax-myrtle, broadleaf weeds, and grasses is desired, apply 2 to 3 quarts of Garlon 4 Ultra per acre in tank mix combination with labeled rates of Accord Concentrate or Accord SP herbicide.

These treatments may be broadcast during site preparation of flat planted or bedded sites or, on bedded sites, applied in bands over the top of beds. For best results, apply in late summer or fall. Efficacy may not be satisfactory when applications are made in early season prior to August.

Note: Do not apply after planting pines.

Directed Sprays Applications for Conifer Release

To release conifers from competing hardwoods and brush such as red maple, sugar maple, striped maple, sweetgum, red and white oaks, ash, hickory, alder, birch, aspen, pin cherry, *Ceanothus* spp., blackberry, chinquapin, and poison oak, mix 4 to 20 quarts of Garlon 4 Ultra in enough water to make 100 gallons of spray mixture. This spray mixture should be directed onto foliage of competitive hardwoods using backpack sprayers with flat fan nozzles or equivalent anytime after the hardwoods and brush have reached full leaf size, but before autumn coloration. The majority of treated hardwoods and brush should be less than 6 feet in height to ensure adequate spray coverage. Care should be taken to direct spray solutions away from contact with conifer foliage, particularly foliage of desirable pines. See Table 1 for relationship between mixing rate, spray volume and maximum application rate.

Note: Spray may cause temporary damage and growth suppression where contact with conifers occurs; however, injured conifers should recover and grow normally. Over-the-top spray applications can kill pines.

Broadcast Applications for Mid-Rotation Understory Brush Control in Southern Coastal Flatwoods Pine Stands (Ground Equipment Only)

For control of susceptible species, such as gallberry and wax-myrtle, and broadleaf weeds, apply 2 to 4 quarts of Garlon 4 Ultra per acre. To broaden the spectrum of woody plants controlled to include fetterbush, staggerbush, and titi, apply 2 to 3 quarts of Garlon 4 Ultra per acre in tank mix combination with labeled rates of Arsenal Applicator's Concentrate. Saw-palmetto will be partially controlled by use of Garlon 4 Ultra at 4 quarts per acre or by mixtures of Garlon 4 Ultra at 2 to 3 quarts per acre in tank mix combination with either Arsenal Applicator's Concentrate or Escort herbicide. These mixtures should be broadcast applied over target understory brush species, **but to prevent injury to pines, make applications underneath the foliage of pines.** Apply sprays in 30 gallons or more per acre of total volume. For best results, apply in late summer or fall. Efficacy may not be satisfactory when applications are made in early season prior to August.

Broadcast Applications for Conifer Release in the Pacific Northwest and California

Dormant Conifers Before Bud Swell (Excluding Pines): To control or suppress deciduous hardwoods such as vine maple, bigleaf maple, alder, scotch broom, or willow **before leaf-out**, or evergreen hardwoods such as madrone, chinquapin, and *Ceanothus* spp., use Garlon 4 Ultra at 1 to 2 quarts per acre. Use diesel or fuel oil as a diluent, or use water plus 1 to 2 gallons per acre of diesel oil or a suitable surfactant or oil substitute at manufacturer's recommended rates. **Mixing with oil as the only diluent requires vigorous agitation to form an oil solution.** Once a solution is formed it will stay stable.

Conifer Plantations (Excluding Pines) After Hardwoods Begin Growth and Before Conifer Bud Break ("Early Foliar" Hardwood Stage): Use Garlon 4 Ultra at 1 to 1.5 quarts alone or with Freelexx or 2,4-D low volatile ester herbicide in water carrier to provide no more than 3 lb ae per acre from both products. After conifer bud break, these sprays may cause more serious injury to the crop trees. Use of a surfactant may cause unacceptable injury to conifers especially after bud break.

Conifer Plantations (Excluding Pines) After Conifers Harden Off In Late Summer and While Hardwoods Are Still Growing Actively: Use Garlon 4 Ultra at rates of 1 to 1.5 quarts per acre alone or with Freelexx or a 2,4-D low volatile ester. Treat as soon after conifer bud hardening as possible so that hardwoods and brush are actively growing. Use of oil, oil substitute, or surfactant may cause unacceptable injury to the conifers.

Broadcast Applications for Conifer Release in the Eastern United States

To release spruce, fir, red pine, and white pine from competing hardwoods such as red maple, sugar maple, striped maple, alder, birch (white, yellow, and grey), aspen, ash, pin cherry, and *Rubus* spp. and perennial and annual broadleaf weeds, use Garlon 4 Ultra at rates of 1.5 to 3 quarts per acre alone or with Freelexx or a 2,4-D amine or low volatile ester. Apply in late summer or early fall after conifers have formed their overwintering buds and hardwoods are in full leaf and prior to autumn coloration.

Broadcast Applications for Conifer Release in the Lake States Region

To release spruce, fir, and red pine from competing hardwoods such as aspen, birch, maple, cherry, willow, oak, hazel, and *Rubus* spp. and perennial and annual broadleaf weeds, use Garlon 4 Ultra at rates of 1.5 to 3 quarts per acre. Apply in late summer or early fall after conifers have formed their overwintering buds and hardwoods are in full leaf and prior to autumn coloration.

Basal Bark and Dormant Stem Treatments

Individual plant treatments such as basal bark and cut surface applications may be used on any use site listed on this label at a maximum use rate of 8 lb ae of triclopyr per acre. See above in the section **Basal Bark, Dormant Stem and Cut Surface Treatments for use on all site** for more use information.

Low Volume Basal Bark Treatment

To control susceptible woody plants such as mesquite, huisache, red maple, red and white oak, birches and aspen with stems less than 6 inches in basal diameter.

Streamline Basal Bark Treatment

To control or suppress susceptible woody plants such as mesquite, huisache, red maple, white and red oak, elbowbush, greenbriar, hackberry, pricklyash, yaupon and wild grape

Cut Stump, Basal Cut Stump, Dormant Stem, Thinline Basal Bark Treatments

To control resprouting, apply undiluted Garlon 4 Ultra to wet the cambium and adjacent wood around the entire circumference of cut stumps. Treatments may be applied throughout the year; however, control may be reduced with treatment during periods of moisture stress as in late summer. Cut stumps so that they are approximately level to facilitate uniform coverage of Garlon 4 Ultra. Use an applicator which can be calibrated to deliver the small amounts of material required.

Growing Point and Leaf Base (Crown) Treatment of Yucca

Prepare a 2% v/v solution of Garlon 4 Ultra in basal oil, diesel or fuel oil (13 fl oz of Garlon 4 Ultra in 5 gallons of spray mixture). Thoroughly wet the center of the plant including growing point and leaf bases to the soil surface. Complete coverage of leaves is not necessary.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, Dow AgroSciences **MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the extent permitted by law, all such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

To the extent permitted by law, Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

[®][™] Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

**Produced for
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268**

Label Code: CD02-329-020
Replaced Label: D02-329-005
EPA accepted 02/05/18