

**KILLS
RATS, MICE
& VOLES**

Kaput[®]



MOUSE BLOCKS

This product may only be used inside and within 100 feet of buildings or inside of transport vehicles (ships, trains, or aircraft).



KILLS RATS, MICE & VOLES

**ABOUT
350
BLOCKS**

NET WT. 22 LBS (9.98 kg)

ACTIVE INGREDIENT:

Warfarin (CAS No. 81-81-2)	0.025%
OTHER INGREDIENTS	99.975%
TOTAL	100.000%

EPA Reg. No. 72500-7

EPA Est. 72500-CO-1

Keep Out of Reach of Children

CAUTION

Harmful if swallowed.

See back panel for First Aid and additional Precautionary Statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL:

Read this entire label and follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets or other non-target wildlife animals to rodenticides. To help prevent exposure:

1. Store product not in use in locations out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, OR in tamper-resistant bait stations. These stations must be resistant to destruction by children under six years of age and dogs, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals or areas prone to vandalism.
3. Dispose of product container and unused, spoiled and unconsumed bait as specified on this label.

NOTE: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

RATS AND MICE

Use Restrictions: This product may only be used to control Norway rats, roof rats, house mice, deer mice, and white-footed mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not broadcast bait. Burrow baiting with this product is prohibited.

Do not sell this product in individual containers holding less than four (4) pounds of bait.

Selection of Treatment Areas: Determine where target rodents will most likely find and consume the bait. Generally, these areas will be along walls and building foundations, in cellars and crawl spaces, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of target rodents have been seen. When controlling white-footed mice, the bases of trees located in permitted use sites (See **Use Restrictions**) may provide suitable baiting points. Remove as much alternative food as possible.

Application Directions:

Mice: Apply 1 block per placement. Use 2 blocks only at points of extremely high mouse activity. Space placements at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for 15 days or until there are no longer fresh signs of mouse activity or feeding on bait.

Rats: Apply 4 to 16 bait blocks per placement. Space placements at 15 to 30 foot intervals. Maintain an uninterrupted supply of fresh bait for 10 days or until there are no longer fresh signs of rat activity or feeding on bait.

Follow-up: While wearing gloves, collect and properly dispose of all visible dead animals and leftover bait. To discourage

reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.

VOLES

Use Restrictions: This product may only be used to control pine voles (*Microtus pinetorum*), meadow voles (*M. pennsylvanicus*), mountain voles (*M. montanus*), and California voles (*M. californicus*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not broadcast bait. Burrow baiting with this product is prohibited.

Do not sell this product in individual containers holding less than four (4) pounds of bait.

Selection of Treatment Areas: Determine areas within 100 feet of man-made structures where voles or signs of vole activity (girdling of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

Application Directions: Place 1 bait block in each bait station. Situate bait stations near runways, burrow openings or plants that voles have been damaging. Maintain an uninterrupted supply of fresh bait for 15 days or until there are no longer fresh signs of vole activity or feeding on bait.

Follow-up: While wearing gloves, collect and properly dispose of all visible dead animals and leftover bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Non-refillable container; do not reuse or refill this container. Offer for recycling, if available, or reconditioning, if appropriate; otherwise, dispose of empty container in a sanitary landfill; or by incineration.

Lot Number: See Container Lid

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets.

All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

FIRST AID HAVE LABEL WHEN OBTAINING TREATMENT ADVICE

If
Swallowed

- Call a poison control center, doctor, or National Pesticide Information Network (1-800-858-7378) immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian or 1-800-222-1222 at once.

NOTE TO PHYSICIAN OR VETERINARIAN

Contains the anticoagulant Warfarin. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. For humans or animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1 orally or intramuscularly.

Environmental Hazards

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinseate.

WARRANTY

To the extent consistent with applicable law, Manufacturer and Seller make no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

Notice: This product contains a chemical known to the state of California to cause birth defects or other reproductive harm.

Scimetrics
LTD. CORP.

Pest Management Solutions

P.O. Box 1045, Wellington, CO 80549-1045

(970) 482-1330

customerservice@kaputproducts.com

Made in U.S.A.

Label Version 1/16/13

