

GROUP U15 FUNGICIDE

PULL HERE TO OPEN ►



syngenta.

## Fungicide

For control of diseases of ornamental plants grown in greenhouses, outdoor growing structures (including shade houses, lath houses and other outdoor growing structures), and nurseries and outdoor ornamental plants grown in commercial landscapes

*Active Ingredient:*

Oxathiapiprolin\* . . . . . 18.7%

*Other Ingredients* . . . . . 81.3%

*Total:* . . . . . 100.0%

\*CAS No. 1003318-67-9

Segovis® is a water-based suspension concentrate that contains 18.7% oxathiapiprolin, equivalent to 1.67 pounds per gallon of active ingredient.

### KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1533      Product of Germany  
EPA Est. 072344-MO-004      Formulated in the USA

SCP 1533A-L1 0815      4056107

**1 pint (16 fluid ounces)** Net Contents



### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

### HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident),  
Call  
**1-800-888-8372**

## PRECAUTIONARY STATEMENTS

### Personal Protective Equipment

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes and socks

### User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

*continued...*

## PRECAUTIONARY STATEMENTS *(continued)*

### Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Segovis must be used only in accordance with instructions on this label, in separately issued labeling or exemptions under FIFRA (Supplemental Labels, Special Local Need Registration, FIFRA Section 18 exemptions), or as otherwise permitted by FIFRA. Always read the entire label, including the Conditions of Sale and Limitation of Warranty and Liability.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes and socks

**FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.**

## PRODUCT INFORMATION

Apply Segovis as a foliar protectant fungicide for control of diseases caused by downy mildews and *Phytophthora* spp. Segovis can also be applied as a drench to plants grown in containers and in-ground for control of root and stem diseases caused by *Phytophthora* spp. Apply Segovis prior to disease development, rotating with other effective fungicides having a different mode of action.

### Application

Mix only the amount of spray solution needed for immediate application. Thorough coverage is necessary to provide good foliar disease control. Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip).

### Adjuvants

For some ornamental uses on this label, a spreading/penetrating type adjuvant such as a non-ionic surfactant, crop oil concentrate, silicone based, or blend may be added at the manufacturer's recommended rates to improve coverage on waxy or hard-to-wet plant surfaces. When an adjuvant is to be used with this product, the use of an adjuvant that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification program is recommended.

### Efficacy

Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if maximum amount of Segovis has been used. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure, highly susceptible varieties, or when environmental conditions conducive to disease exist.

### Integrated Pest Management (IPM)

Segovis should be integrated into an overall disease and pest management strategy whenever the use of a fungicide is required. Cultural practices known to reduce disease development should be followed. Consult your local agricultural authorities for additional IPM strategies established for your area. Segovis may be used in State Agricultural Extension advisory (disease forecasting) programs which recommend application timing based on environmental factors favorable for disease development.

## Resistance Management

**GROUP U15 FUNGICIDE**

Fungal pathogens can develop resistance to products with the same mode of action when used repeatedly. Because resistance development cannot be predicted, use of this product should conform to resistance management strategies established for the crop and use area. Consult your local or state agricultural authorities for resistance management strategies that are complementary to those in this label. Resistance management strategies may include rotating and/or tank-mixing with products having different modes of action or limiting the total number of applications per season. Syngenta encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label. Segovis must not be alternated or tank mixed with any fungicide to which resistance has already developed.

Segovis contains the active ingredient oxathiapiprolin, which inhibits the oxysterol-binding protein (OSBP). Repeated use of products for control of specific plant pathogens may lead to selection of resistant strains of fungi and result in a reduction of disease control. A disease management program that includes rotation and/or tank mixing with fungicides with a different mode of action is essential to reduce the risk of fungicide resistance development. Do not tank mix Segovis with any fungicide for which resistance to the target disease has developed. Unless otherwise specified in the specific use directions, make no more than 2 sequential applications of Segovis before rotating to a fungicide with a different mode of action. Do not use Segovis for more than 33% of the total fungicide applications per season per crop. Do not follow soil applications of Segovis with foliar applications. Different application methods (foliar and soil) must not be combined when protecting a crop during a growing season. For guidance on a particular crop and disease control situation, consult your state extension specialist or official state recommendations.

### Spray Drift Management

To avoid spray drift, do not apply when conditions favor drift beyond the target area. The interaction of many equipment and weather related factors determine the potential for spray drift. AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

## MIXING INSTRUCTIONS

### Mixing Instructions

- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.

- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.
- Do not allow spray mixture to stand overnight or for prolonged periods of time (more than 3 hours) without agitation.

#### **Segovis Alone (no tank mix)**

- Add  $\frac{1}{2}$  -  $\frac{2}{3}$  of the required amount of water to the spray or mixing tank.
- With the agitator running, add Segovis to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after Segovis has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

#### **Segovis + Tank Mixtures**

Segovis is usually compatible with all tank-mix partners. To determine the physical compatibility of Segovis with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

**It is important to mix only the amount of product that can be sprayed immediately. Continuous agitation is recommended. If circumstances cause a delay, the product(s) may settle and be difficult to re-suspend. If this occurs, good agitation is required for a minimum of 15 minutes before and during spray operation.**

#### **Mixing in the Spray Tank**

- Add  $\frac{1}{2}$  -  $\frac{2}{3}$  of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and Segovis to the spray tank.
- Allow Segovis to completely disperse.
- Spray the mixture with the agitator running.

## APPLICATION EQUIPMENT

Segovis may be applied with application equipment commonly used for greenhouse and nursery crop production. Proper adjustments and calibration of spraying equipment to give good canopy penetration and coverage is essential for good disease control.

### APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION)

- Apply this product only through overhead, hand held, micro-irrigation systems (e.g., drip, trickle, spaghetti tubes and micro sprinklers) and motorized calibrated irrigation systems. Do not apply this product through any other type of irrigation system.
- Plant injury and/or poor disease control can result from non-uniform distribution of treated water.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
- Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended.

### **Motorized Calibrated Irrigation Equipment**

- Determine the acreage covered by the sprinklers.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 20 to 30-minute interval. When applying Segovis through irrigation equipment, use the lowest obtainable water volume while maintaining uniform distribution.
- Determine the amount of Segovis required to treat the area covered by the irrigation system.
- Add the required amount of Segovis into the same quantity of water used to calibrate the injection period.
- Operate the system at the same pressure and time interval established during the calibration.
- Stop injection equipment after treatment is completed. Continue to operate the system until the Segovis solution has cleared the last sprinkler head.

### **Specific Instructions for Public Water Systems**

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.



6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

## ORNAMENTAL USE DIRECTIONS

### APPLICATION INSTRUCTIONS

#### Foliar Application

Make foliar applications in an adequate water volume to achieve thorough and uniform coverage without excessive runoff (to drip). Apply Segovis prior to disease development to achieve best disease control.

#### Drench Applications – Ornamentals

- Prepare the Segovis drench solution according to Table 1 below.
- Apply enough drench solution to thoroughly wet the root zone of the plants without leaching through the container. For plants grown in flats or beds apply 1–2 pt of drench solution per sq ft. For plants grown in containers, refer to the suggested drench volumes listed below. For container sizes not listed, adjust volume appropriately.
- Make no more than 2 drench applications of Segovis before switching to another effective non-Group U15 fungicide.

**Table 1. Volume of Segovis Drench Solution by Container Diameter**

Container Size (Diameter – Inches)	Drench Solution Per Container (fl oz)
4	3
5	4
6	6
8	10
10	20
12	30

Segovis may be applied by drench to container, bench, flat, plug, liner, bed-grown ornamentals in nurseries, forest nurseries, greenhouses, lath and shade houses or other ornamental production structures. Apply according to the use directions in **Table 1**.

### Soil Applications – In-ground-grown Plants

- Apply in a manner that ensures the product solution adequately saturates the target crop root/crown zone.
- Apply using drip application, surface band, or directed application or in-furrow application.
- Apply surface band or directed applications in a 4- to 12-inch band.
- If the application method does not move the product to the target root/crown disease zone, the application must be followed with irrigation or cultivation to correctly place the product for disease control.

**Table 2. Soil Application Rates for Segovis Product per 1,000 feet of row (based on plant spacing)**

Segovis Conversion Chart for Drip (Trickle) Chemigation and Directed/Banded/In-Furrow application.							
Corresponding field rate (fl oz/acre)	Rate in fl oz of product/1,000 row ft, based on plant row spacing						
	30"	34"	36"	48"	60"	72"	84"
2.4	0.14	0.16	0.17	0.22	0.28	0.33	0.39
4.8	0.28	0.31	0.33	0.44	0.55	0.66	0.77
9.6	0.55	0.62	0.66	0.88	1.10	1.32	1.54
19.2	1.10	1.25	1.32	1.76	2.20	2.65	3.09

### Restriction

**Do not** apply more than 0.5 lb ai per acre per calendar year for any combination of outdoor-applied foliar and soil uses.

**Table 3. Foliar Diseases Controlled with Segovis**

Ornamental Crops	Diseases/ Pathogens	Rate (fl oz/ 100 gal)	Remarks
Pot and Bedding Plants Breeding Crops Bulb Crops (including Calla Lillies, Easter Lillies, Gladiolas, and Caladiums) Cut Flowers Evergreens (including conifers) Flowers Grown for Seed Production Flowering Plants Foliage Plants Ground Covers Non-bearing Fruit Trees Non-bearing Nut Trees Non-bearing Vines Ornamental Tree and Shrubs Perennial Plants	Downy Mildew Diseases	0.6 – 2.4	Begin foliar applications prior to disease development and continue on a 5- to 14-day interval when conditions are favorable for disease development. Make no more than 2 consecutive applications before switching to another effective non-Group-U15 fungicide. Use the shorter interval and/or higher rates under high disease pressure or when conditions are conducive to disease development. Segovis may be tank mixed with another fungicide labeled for downy mildew that has a different mode of action. Segovis may be tank mixed with Subdue Maxx®, Heritage®, or Micora™ for broader spectrum disease control. <sup>1</sup>
	Phytophthora Diseases	0.6 – 2.4	Apply Segovis as a foliar and stem spray on a 5- to 14-day interval preventatively or at first sign of disease symptoms. Make no more than 2 consecutive applications before switching to another effective non-Group-U15 fungicide.

<sup>1</sup> Observe all precautions and limitations on labeling of all products used in tank mixes.

**Restrictions**

- Do not apply greater than 2.4 fl oz Segovis per acre in a single application.

- For plants grown outdoors, **do not** apply more than 9.6 fl oz of Segovis per acre per calendar year (0.125 lb ai per acre per calendar year).
- For plants grown indoors, **do not** apply more than 9.6 fl oz of Segovis per acre per crop (0.125 lb ai per acre per crop). Do not apply to more than 8 crops grown in the same growing area per year.

**Table 4. Soil Diseases Controlled with Segovis –Container-Grown Plants**

Ornamental Crops	Diseases/ Pathogens	Rate (fl oz/ 100 gal)	Remarks
Pot and Bedding Plants Breeding Crops Bulb Crops (including Calla Lillies, Easter Lillies, Gladiolas, and Caladiums) Cut Flowers Evergreens (including conifers) Flowers Grown for Seed Production Flowering Plants Foliage Plants Ground Covers Non-bearing Fruit Trees Non-bearing Nut Trees Non-bearing Vines Ornamental Tree and Shrubs Perennial Plants	Phytophthora Diseases	0.65 – 3.2	Apply Segovis as a drench application to container grown plants on a 5- to 14-day interval preventatively or at first sign of disease symptoms. Refer to <b>Drench Applications – Ornamentals</b> section and <b>Table 1</b> for more information. Make no more than 2 consecutive applications before switching to another effective non-Group-U15 fungicide.

**Restrictions**

- **Do not** apply greater than 19.3 fl oz Segovis per acre in a single application.
- For plants grown outdoors, **do not** apply more than 38.6 fl oz of Segovis per acre per calendar year (0.5 lb ai per acre per calendar year).
- For plants grown indoors, **do not** apply more than 38.6 fl oz of Segovis per acre per crop (0.5 lb ai per acre per crop). Do not apply to more than 8 crops grown in the same growing area per year.

**Table 5. Soil Diseases Controlled with Segovis – In-Ground-Grown Plants**

Ornamental Crops	Diseases/ Pathogens	Rate (fl oz)	Remarks
Pot and Bedding Plants Breeding Crops Bulb Crops (including Calla Lillies, Easter Lillies, Gladiolas, and Caladiums) Cut Flowers Evergreens (including conifers) Flowers Grown for Seed Production Flowering Plants Foliage Plants Ground Covers Non-bearing Fruit Trees Non-bearing Nut Trees Non-bearing Vines Ornamental Tree and Shrubs Perennial Plants	Phytophthora Diseases	4.8 – 38.6 per 100 gallons [1 gallon pack size or larger] 2.4 – 19.3 per 50 gallons [1 quart pack size or larger] 1.2 – 9.15 per 25 gallons [1 pint pack size or larger]	Apply Segovis as a soil applica- tion to plants preventatively or at first sign of disease symptoms. Refer to <b>Soil Applications –                      In-ground-grown Plants</b> and <b>Table 2</b> for more information. The minimum retreatment interval is 7 days for soil applied applications. Make no more than 2 con- secutive applications before switching to another effective non-Group-U15 fungicide.

**Restrictions**

- **Do not** apply greater than 19.3 fl oz Segovis per acre in a single application.
- **Do not** apply more than 38.6 fl oz of Segovis per acre per calendar year (0.5 lb ai per acre per calendar year).

### **Plant Safety**

Segovis has been shown to be safe when applied at the recommended rates to ornamental plants. However, due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to Segovis. Neither the manufacturer nor the seller has determined whether Segovis can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The user should conduct small scale testing at the recommended rates to ensure plant safety prior to broad scale commercial use on plant genera and species not listed in this label.

When using an adjuvant or tank-mix partner, the user should conduct small scale testing at the recommended rates to ensure plant safety prior to broad scale commercial use.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

### **Pesticide Storage**

Store in original containers only. Store in a cool, dry place. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

### **Pesticide Disposal**

Pesticide wastes may be acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

### **Container Handling [less than or equal to 5 gallons]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

*continued...*

## **STORAGE AND DISPOSAL (continued)**

### **Container Handling [greater than 5 gallons – mini-bulk]**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

### **Container Handling [greater than 5 gallons - bulk]**

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

## **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

---



Segovis®, Daconil Action™, Heritage®,  
Micora™, Subdue Maxx®, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company



©2015 Syngenta

For non-emergency (e.g., current product information), call  
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1533A-L1 0815**  
**4056107**

GROUP U15 FUNGICIDE



**Segovis®**

## Fungicide

For control of diseases of ornamental plants grown in greenhouses, outdoor growing structures (including shade houses, lath houses and other outdoor growing structures), and nurseries and outdoor ornamental plants grown in commercial landscapes

Active Ingredient:

Oxathiapiprolin\* . . . . . 18.7%

Other Ingredients . . . . . 81.3%

Total: . . . . . 100.0%

\*CAS No. 1003318-67-9

Segovis® is a water-based suspension concentrate that contains 18.7% oxathiapiprolin, equivalent to 1.67 pounds per gallon of active ingredient.

## KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1533 EPA Est. 072344-MO-004

Segovis® is a trademark of a Syngenta Group Company.


SCP 1533A-L1 0815 4056107

**1 pint**

(16 fluid ounces)

Net Contents





**FIRST AID:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**HOT LINE NUMBER:** For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call **1-800-888-8372**.

**PRECAUTIONARY STATEMENTS**

**Environmental Hazards:** For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original containers only. Store in a cool, dry place. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

**Pesticide Disposal:** Pesticide wastes may be acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER.**

©2015 Syngenta  
Manufactured for:  
Syngenta Crop Protection, LLC  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1533A-L2 0815 4056108**