

Devito™



For Both Indoor and Outdoor Use.

For use in, on and around buildings and structures for the control of listed pests, including on non-residential lawns, ornamental trees and shrubs around residential, institutional, public, commercial, agricultural and industrial buildings.

ACTIVE INGREDIENT:

(% by weight)

Lambda-cyhalothrin*: [1 α (S*), 3 α (Z)]-(\pm)-cyano-(3-phenoxyphenyl) methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate. 9.7%

OTHER INGREDIENTS: 90.3%

TOTAL 100.0%

*Synthetic pyrethroid

EPA Reg. No.: 91234-250

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	<ul style="list-style-type: none">▪ Call a poison control center or doctor immediately for treatment advice.▪ Have person sip a glass of water if able to swallow.▪ Do not induce vomiting unless told to do so by the poison control center or doctor.▪ Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. To protect the environment, **do not** allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run-off to water bodies or drainage systems. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.

PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard. **Do not** mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

PRODUCT INFORMATION

Devito with EnduraCap Technology is a unique formulation utilizing EnduraCap™ Technology which is a proprietary blend of ingredients for use with the active ingredient Lambda-cyhalothrin.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For both indoor and outdoor use.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on lawns or grounds.

This product is RESTRICTED for use in New York State.

Application Restrictions for Residential Outdoor Surface and Space Sprays:

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

- **Do not** apply or irrigate to the point of runoff.
- **Do not** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- **Do not** apply the product into fish pools, ponds, streams, or lakes. **Do not** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

- **Do not** allow to enter indoor or outdoor drains unless labeled for drain treatments.

No permita la entrada a desagües internos o externos a menos que el etiquetado indique que está permitido el uso del producto para tratamiento de desagües.



- Follow proper disposal procedures on this label.

Siga las indicaciones del etiquetado para el desecho apropiado del producto.

- **Do not** allow the product to enter any drain during or after application except where labelled for this use.
- **Do not** allow children or pets to contact treated surfaces until spray has dried.
- Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).
- **Do not** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

PERIMETER PEST CONTROL

Devito with EnduraCap Technology is for use as a general surface (nonfood/nonfeed areas), crack and crevice, or spot treatment in, on, and around buildings and structures and their immediate surroundings, and on modes of transport. Areas of use: industrial buildings, houses, patios and porches (spot and crack and crevice treatment only), closets, furniture, apartment buildings, mobile homes, laboratories, buses, greenhouses (non-commercial), stores, factories, warehouses, wineries, vessels, railcars, trucks, trailers, aircraft (cargo and other non-cabin areas only), schools, nursing homes, hospitals, mausoleums, restaurants, hotels, correctional facilities, livestock/poultry housing, pet kennels, food granaries, food grain mills and food manufacturing, processing, and servicing establishments.

Nonfood/nonfeed areas of food/feed handling establishments: garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

For indoor applications, retreat at 21-day intervals to maintain control. More frequent treatments are limited to spot treatments. Do not apply to carpets as a pin stream or coarse spray treatment.

Mixing Instructions

Dilute **Devito with EnduraCap Technology** with water for application using hand-held or power-operated application equipment as a coarse spray for crack and crevice or spot and general surface (nonfood/nonfeed areas only) treatments. Application equipment that delivers low volume treatments (such as the Micro-Injector® or Actisol® applicator) may also be used to make crack and crevice or spot and general surface treatments. Fill applicator tank with the desired volume of water and add **Devito with EnduraCap Technology**. Close and shake before use in order to ensure proper mixing. Shake or re-agitate applicator tank before use if application is interrupted. Mix only amount of treatment volume as required. **Devito with EnduraCap Technology** may be applied by using a paintbrush or other porous applicator attached to a handle as a general surface treatment.

Tank Mixing

Devito with EnduraCap Technology may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, perform a small volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Devito with EnduraCap Technology** last. If mixed with EC formulations, use within 24 hours. Fill tank to desired volume and continue to agitate while making applications.

Devito with EnduraCap Technology may be tank mixed with an Insect Growth Regulator (IGR). It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Foam Applications

Devito with EnduraCap Technology may be converted to a foam and the foam used to treat structural voids to control or prevent pests including ants, bees, termites (above ground only), wasps, or other listed arthropods harboring in walls, under slabs, or in other void areas.

RATES FOR PESTS (HAND APPLICATION EQUIPMENT)

PESTS	DILUTION RATE	SPECIFIC USE DIRECTIONS
Ants Bedbugs (adult) Bees Boxelder Bugs Carpenter Bees Carpet Beetles Centipedes Cigarette Beetles Clover Mites Cockroaches ¹ Confused Flour Beetles Crickets Earwigs Firebrats Fleas ² Flies Lesser Grain Borers Millipedes Mosquitoes Red Flour Beetles Rice Weevils Saw-toothed Grain Beetles Silverfish Sow Bugs Spiders Stink Bugs Ticks Wasps	0.2 fl. oz. (6 mL) to 0.4 fl. oz. (12 mL) per gallon of water (0.015 - 0.03% AI)	<p>Ants: Apply to any trails around doors and windows and other places where ants may be found. Locate and treat nests. Where ants are trailing inside, apply as a residual surface treatment to active areas: baseboards, corners, around pipes, in and behind cabinets, behind and under refrigerators, sinks, furnaces and stoves, cracks and crevices. When combining baits and residual surface insecticides, apply surface insecticides in cracks and crevices, along baseboards, and infested surfaces and outside barrier treatments. Use baits in other areas that are untreated by residual insecticides; also see Outdoor Surfaces Use.</p> <p>Bedbugs: Clean floors and surfaces by vacuuming. Apply as a coarse, low-pressure spray to harborage areas: crevices, baseboards, loose plaster, behind bed frames and headboards, beneath beds and furniture, and to bedsprings and bed frames. Do not apply to furniture surfaces or mattresses where people will be laying or sitting. Infested bedding should not be treated; remove bedding, place in sealed plastic bags, and take for laundering and drying at high temperatures.</p> <p>Cockroaches, Crickets, Earwigs, Firebrats, Silverfish, Spiders and Stink Bugs: Apply as a coarse, low pressure spray to areas where these pests hide: baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, cabinets, behind and under refrigerators, sinks, furnaces and stoves, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices; also see Outdoor Surfaces Use.</p> <p>Bees, Flies, Mosquitoes, and Wasps: Apply directly to walls, ceilings, window screens, and other resting areas as a residual surface treatment. May be used inside residential buildings as well as in and around carports, garages, and storage sheds; also see Outdoor Surfaces Use. Use caution when treating nests of stinging insects as Devito with EnduraCap Technology does not provide instant knockdown. Protective equipment for the applicator against stinging pests may be required. Treat bee, wasp and hornet nests late in the day when most insects will be present. Allow 2 - 3 days for the colony to die and repeat after 21 days, if necessary.</p> <p>Yards or other frequented areas enclosed by landscaping can benefit from the creation of a mosquito barrier to reduce invading mosquitoes by the treatment of perimeter vegetation. Apply Devito with EnduraCap Technology at specified rates in 2 - 5 gals. of water per 1,000 sq. ft.</p> <p>Carpenter Bees: Apply coarse spray to thoroughly wet wood surfaces where bees have been previously active or to provide protection against further damage. Apply early in the spring to prevent bees from invading wood. When bees have infested wood, surface applications can help control embedded larvae and bees that emerge from the wood.</p> <p>Pantry Pests (i.e., Carpet beetle, Cigarette beetle, Confused flour beetle, Lesser grain borer, Red flour beetle, Rice weevil, and Saw-toothed grain beetle): Apply to cupboards, shelving, and storage areas. Remove all utensils, uncovered foodstuffs (or any having original package opened), and shelf paper before making application. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff, or other items. Destroy any food-stuff accidentally contaminated with treatment solution.</p> <p>Boxelder Bugs, Centipedes, Millipedes, Pillbugs, and Sow Bugs: Apply around doors and windows and other places where these pests may be found or where they may enter premises.</p> <p>Treat baseboards, storage areas, and other locations listed on this label. Apply treatments to prevent infestation as described below; also see Outdoor Surfaces Use.</p> <p>Fleas and Ticks: To control nuisance fleas and ticks (e.g. dog ticks) apply to kennels, yards, runs, and other areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent.</p> <p>Treat entire area rather than making spot treatments, and retreat as necessary to maintain control, as permissible on this label. Do not apply to pasture or cropland, and do not allow animals and people access to treated areas until the deposit has dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks; also see Outdoor Surfaces Use. Do not apply directly to animals.</p> <p>Cluster Flies: Apply in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors, paying particular attention to south-facing surfaces, but do not allow runoff to occur. Heavy precipitation prior to frost may require retreatments to maintain protection. In winter and spring when flies become active and are emerging, interior crack and crevice and void treatments can help reduce the infestation, along with ULV or general surface application in infested attics or unoccupied lofts.</p> <p>Litter Beetles (Darkling, Hide, and Carrion Beetles) and Flies In Animal Housing (Such As Poultry Houses): To control adult litter beetles, apply Devito with EnduraCap Technology to walls and floors at cleanout, before reintroduction of animals. This will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to areas where beetles frequently occur, walls, supports, cages, stalls, and around feeders. To help control flies, apply a directed application to horizontal surfaces and overhead areas and allow to dry before reintroduction of animals; also see Livestock/ Poultry Housing Structures and Pet Kennels.</p>
Cockroaches ¹ Crickets ⁶ Flies ⁵ Litter Beetles ³ (adults/immature stages such as darkling, hide, and carrion) Mosquitoes ⁴ Pillbugs Scorpions Spiders ⁵ Spider Mites (Two-spotted, Spruce) Stink Bugs ⁶ Ticks ⁶	0.8 fl. oz. (24 mL) per gallon of water (0.06% AI)	<p>Boxelder Bugs, Centipedes, Millipedes, Pillbugs, and Sow Bugs: Apply around doors and windows and other places where these pests may be found or where they may enter premises.</p> <p>Treat baseboards, storage areas, and other locations listed on this label. Apply treatments to prevent infestation as described below; also see Outdoor Surfaces Use.</p> <p>Fleas and Ticks: To control nuisance fleas and ticks (e.g. dog ticks) apply to kennels, yards, runs, and other areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent.</p> <p>Treat entire area rather than making spot treatments, and retreat as necessary to maintain control, as permissible on this label. Do not apply to pasture or cropland, and do not allow animals and people access to treated areas until the deposit has dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks; also see Outdoor Surfaces Use. Do not apply directly to animals.</p> <p>Cluster Flies: Apply in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors, paying particular attention to south-facing surfaces, but do not allow runoff to occur. Heavy precipitation prior to frost may require retreatments to maintain protection. In winter and spring when flies become active and are emerging, interior crack and crevice and void treatments can help reduce the infestation, along with ULV or general surface application in infested attics or unoccupied lofts.</p> <p>Litter Beetles (Darkling, Hide, and Carrion Beetles) and Flies In Animal Housing (Such As Poultry Houses): To control adult litter beetles, apply Devito with EnduraCap Technology to walls and floors at cleanout, before reintroduction of animals. This will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to areas where beetles frequently occur, walls, supports, cages, stalls, and around feeders. To help control flies, apply a directed application to horizontal surfaces and overhead areas and allow to dry before reintroduction of animals; also see Livestock/ Poultry Housing Structures and Pet Kennels.</p>

PESTS	DILUTION RATE	SPECIFIC USE DIRECTIONS
Termites (above ground only)	0.2 fl. oz. (6 mL) to 0.4 fl. oz. (12 mL) per gallon of water (0.015 - 0.03% AI)	Devito with EnduraCap Technology may be used on termites to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. DO NOT use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.
¹ For cockroaches, the rate for maintenance treatments is 0.015% and for clean-out treatments is 0.03%. For control of severe infestations, use 0.06% rate. ² For outdoor use only and use 0.03% rate. ³ For control of LIGHT beetle infestations, use 0.03% rate. ⁴ For residual control, use 0.06% rate. ⁵ In all states except California, rates for flies may be increased to 0.06% when environmental conditions are severe and/or populations are high. ⁶ For clean-out/severe infestations, use 0.06% rate.		

Application within Food Handling Establishments

For places other than private residences in which exposed food is held, processed, prepared, or served: areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing foods, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups), and serving areas.

Use as a crack and crevice or spot treatment in and around both food and nonfood areas. Apply in small amounts directly into cracks and crevices, using equipment capable of delivering a pin stream of insecticide, in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment and bases. Clean food contact surfaces and equipment with an effective cleaning compound and rinse with potable water before using.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.). Take extreme care that the product is not introduced into the air. **Do not** contaminate food and food processing surfaces.

Application within Food Serving Areas

For facilities where foods are served, such as dining rooms. Apply as a crack and crevice or spot treatment to selective surfaces: baseboards, under elements of construction, and into cracks and crevices. **Do not** treat surfaces likely to be contacted by food. (**Do not** apply when facility is in operation or foods are exposed.) Food must be covered or removed in area being treated. **Do not** apply directly to food or allow applications to contaminate food.

NOTE: Application of this product in the Food Areas and/or Food Serving Areas of Food Handling Establishments other than as a spot and/or crack and crevice treatment is not permitted. Limit the use of application equipment (such as the Micro-Injector® or Actisol® applicator) in food areas to crack and crevice treatment only.

Livestock/Poultry Housing Structures and Pet Kennels

Apply as a general surface (including directed sprays) and/or crack and crevice treatment. Make interior and exterior perimeter applications in and around the livestock, poultry, and pet housing structures. Normal cleaning of the structure must also be done along with applications of **Devito with EnduraCap Technology** to effectively control the crawling and flying insect pests listed in the table.

With the exception of cattle and calves that may be present, only apply **Devito with EnduraCap Technology** to livestock barns or housing structures when unoccupied by animals. Apply to floors, vertical, and overhead surfaces where crawling or listed flying insect pests are or may be present. Cover feeders, waterers, and feed carts before application to prevent contamination. **Do not** apply to milk rooms or feed rooms. Pay attention to animal areas: stanchions, pipes, windows and doors, and areas where insect pests hide or congregate. Exterior applications to south facing walls and foundation perimeters can help prevent interior infestations of listed flying and crawling insect pests.

For poultry houses, apply to floor area (birds grown on litter) or to walls, posts, and cage framing (birds grown in cages). Application may also be made into cracks and crevices around insulation. Reapply after each growout or sanitization procedure. Indoor control can

be enhanced by making perimeter treatments around the outside of building foundations to prevent immigrating adult beetles. Apply in a uniform band 1 - 2 ft. up and 2 - 6 ft. out from foundation. Maintaining a year-round treatment program will prevent background populations from reaching problem levels.

RESTRICTIONS:

- **DO NOT** make interior applications of **Devito with EnduraCap Technology** in areas where animals, except cattle and calves, are present in the facility. Allow treated surfaces to completely dry before restocking the facility.
- **DO NOT** make applications to any animal feedstuffs, water, or watering equipment.
- **DO NOT** contaminate any animal food, feed, or water in and around livestock, poultry, or pet housing when making applications.
- **DO NOT** apply directly to animals.

Outdoor Surfaces Use

For control of ants, bees, centipedes, cockroaches, crickets, fleas, flies, millipedes, mosquitoes, scorpions, sowbugs, stink bugs, pillbugs, spiders, termites (above ground only), ticks, wasps, and other listed perimeter arthropod pests.

Apply with either hand or power application equipment as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches (crack and crevice only), screens, window frames, eaves, patios (crack and crevice only), garages, refuse dumps, and other similar areas where these insect pests are active.

For termites, **DO NOT** use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.

Perimeter Barrier Treatments

Apply a continual band of insecticide around building foundations and around windows, doors, service line entrances, eaves, vents, and other areas to reduce the potential for entry by crawling pests. Before application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation. Apply the band up to 7 ft. wide around the structure (or according to state regulations governing commercial pest control) and upwards along the foundation to 2 ft. and around windows, doors, and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active.

Amount of concentrate is dependent upon pest species (see pest table and comments), infestation levels, and service interval desired.

Rate Table for Perimeter Barrier Applications

Fl. oz. (mL) of Devito with EnduraCap Technology	Gals. of Water ¹	Area of Coverage (sq. ft.)
0.2 fl. oz. (6 mL)	1 - 5 gals.	800 - 1,600 sq. ft.
0.4 fl. oz. (12 mL)	1 - 5 gals.	800 - 1,600 sq. ft.
0.8 fl. oz. (24 mL)	1 - 5 gals.	800 - 1,600 sq. ft.

¹ Application volume may be greater than 5 gals./800 - 1,600 sq. ft. if required under heavy vegetative or landscaping materials in order to obtain desired coverage.

Examples of Dilutions for Perimeter Barrier Applications

Application Volume Gals. of Solution per 1,000 sq. ft.	Application Rate fl. oz. of Devito with EnduraCap Technology per 1,000 sq. ft.	Fl. oz. (mL) of Devito with EnduraCap Technology to Dilute in Water According to Spray Tank Volumes		
		5 gallons	10 gallons	50 gallons
1 gal. per 1,000 sq. ft.	0.2 fl. oz. (6 mL)	1 fl. oz. (30 mL)	2 fl. oz. (60 mL)	10 fl. oz. (300 mL)
	0.4 fl. oz. (12 mL)	2 fl. oz. (60 mL)	4 fl. oz. (120 mL)	20 fl. oz. (600 mL)
	0.8 fl. oz. (24 mL)	4 fl. oz. (120 mL)	8 fl. oz. (240 mL)	40 fl. oz. (1200 mL)
2 gal. per 1,000 sq. ft.	0.2 fl. oz. (6 mL)	0.5 fl. oz. (15 mL)	1 fl. oz. (30 mL)	5 fl. oz. (150 mL)
	0.4 fl. oz. (12 mL)	1 fl. oz. (30 mL)	2 fl. oz. (60 mL)	10 fl. oz. (300 mL)
	0.8 fl. oz. (24 mL)	2 fl. oz. (60 mL)	4 fl. oz. (120 mL)	20 fl. oz. (600 mL)
5 gal. per 1,000 sq. ft.	0.2 fl. oz. (6 mL)	0.2 fl. oz. (6 mL)	0.4 fl. oz. (12 mL)	2 fl. oz. (60 mL)
	0.4 fl. oz. (12 mL)	0.4 fl. oz. (12 mL)	0.8 fl. oz. (24 mL)	4 fl. oz. (120 mL)
	0.8 fl. oz. (24 mL)	0.8 fl. oz. (24 mL)	1.6 fl. oz. (48 mL)	8 fl. oz. (240 mL)

Example Calculation: To apply the mid-rate of **Devito with EnduraCap Technology** at a volume of 5 gal./1,000 sq. ft., mix 4 oz of concentrate in 50 gallons of water. The percent active ingredient in the finished **Devito with EnduraCap Technology** dilution can be calculated with the following formula:

$$\text{mL needed to add} = \frac{\text{times } 9.7\% \text{ active in concentrate, divided by gal. finished dilution times } 3,785 \text{ mL/gal.} = \% \text{ active in dilution. (Example: } 4 \text{ oz. in } 50 \text{ gal. is } 120 \text{ mL times } 9.7 \text{ equals } 1,164, \text{ and } 50 \text{ gal. times } 3,785 \text{ is } 189,250. \text{ Dividing } 1,164 \text{ by } 189,250 \text{ equals } 0.006\% \text{ active in the tank dilution.)}}$$

NOTE: Do not use water based sprays of **Devito with EnduraCap Technology** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard. Thoroughly wash out sprayer and screen with water and detergent before using **Devito with EnduraCap Technology**. **Devito with EnduraCap Technology** has not stained or caused damage to painted or varnished surfaces, plastics, fabrics, or other surfaces where water applied alone causes no damage. However, treat a small area and allow to dry, to determine whether staining will occur.

LET TREATED SURFACES DRY BEFORE ALLOWING HUMANS AND ANIMALS TO CONTACT SURFACES.

RESTRICTIONS:

- **DO NOT** use this product with oil.
- **DO NOT** apply this product in any room being used as living, eating, sleeping, or recovery area by patients, the elderly, or infirm when they are in the room.

- **DO NOT** apply to institutions (including libraries, sports facilities, etc.,) in the immediate area when occupants are present.
- **DO NOT** apply to classrooms when in use.
- **DO NOT** apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- **DO NOT** allow applications to contact water inhabited by fish, such as in aquariums and ornamental fish ponds that are located in/around structures being treated.
- **DO NOT** apply to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields or any other area frequented by the general public).

LAWNS/TURFGRASS

Devito with EnduraCap Technology may be used for applications to maintain non-residential indoor or outdoor areas where turf is grown. Treat non-residential landscapes around institutional, public, commercial and industrial buildings. Application rates for turf are lower than structural pest control rates, reflecting that treatment intervals are more frequent.

Applicators must ensure that they are certified in the necessary pesticide certification categories to allow application of **Devito with EnduraCap Technology** away from structures, such as to turf. Structural pest control certification categories may limit the distance away from structures for pesticide application. Consult your state extension office or pesticide regulatory officials for further information.

Mixing Instructions

Devito with EnduraCap Technology is to be mixed with water and may be used in all types of standard application equipment. Fill applicator tank with the desired volume of water and add **Devito with EnduraCap Technology**. It is suggested that the water be 5 - 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add **Devito with EnduraCap Technology** to applicator tank water with maximum agitation. Close and shake or re-agitate applicator tank before use if application is interrupted. Make up only amount of treatment volume as required.

Tank Mixing

Devito with EnduraCap Technology may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, perform a small volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Devito with EnduraCap Technology** last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Follow all restrictions and precautions which appear on the labels of these products.

RESTRICTIONS:

- **KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL SPRAY HAS DRIED FOLLOWING APPLICATION.**
- **DO NOT** apply when turfgrass is waterlogged or when soils are saturated with water (i.e. will not accept irrigation).
- **DO NOT** apply more than 0.36 lbs. of a.i. (52.4 fl. oz. of concentrate) per acre per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. For soil or foliar applications, **DO NOT** apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- **DO NOT** make outdoor broadcast applications to turf when wind speed is 15 mph or greater.
- In the state of New York, **DO NOT** apply within 100 ft. of coastal marshes or streams that drain into coastal marshes.
- **DO NOT** apply to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields, or any other area frequented by the general public).

RATES FOR LAWN AND TURFGRASS PESTS

PEST	AMOUNT OF DEVITO WITH ENDURACAP TECHNOLOGY*	APPLICATION INSTRUCTIONS
Ants (including Imported Fire Ants) Armyworms Centipedes Chinch Bugs ¹ Crickets Cutworms Earwigs Fleas (adult) Grasshoppers Japanese Beetles (adult) Millipedes Mites Mosquitoes (adult) Pillbugs Sod Webworms Sow Bugs Stink Bugs Ticks (including species which transmit Lyme disease)	3.4 - 7 mL per 1,000 sq. ft. or 5 - 10 fl. oz. per acre	<p>Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher labeled rates as pest pressure increases.</p> <p>Apply at 7-day intervals if retreatment is necessary. More frequent treatments should be limited to spot treatments.</p> <p>For spot treatments, use 0.5 fl. oz. of Devito with EnduraCap Technology per 1 - 2.5 gals. of water.</p> <p>Surface Insect Control (armyworm, cutworms, fleas, etc): Apply Devito with EnduraCap Technology at labeled rates in 2 - 5 gals. of water per 1,000 sq. ft. The use of a spreader-sticker may be useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12 - 24 hours to control surface-feeding insect pests.</p> <p>Thatch Inhabiting Insect Control (chinch bugs, billbugs, etc): Apply Devito with EnduraCap Technology at labeled rates in 2 - 10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant or similar adjuvant is permitted at label rates. Lightly irrigate after application with up to 1/2 inch of water to move the Devito with EnduraCap Technology into the thatch layer. If irrigation is not available, then use high water application rates.</p> <p>Subsurface Insect Control (mole crickets, grubs, etc): Apply Devito with EnduraCap Technology at labeled rates in 4 - 10 gals. of water per 1,000 sq. ft. the use of a nonionic wetting agent, penetrant or similar adjuvant is permitted following label rates. Use the highest water application rates possible with your sprayer. Apply Devito with EnduraCap Technology to turf wet with dew, rain or irrigation. Water-in immediately after application with 1/4 - 1/2 inch of water.</p> <p>Fire Ant Control: Treat individual mounds with a drench application using a watering can. Use 0.5 fl. oz. of Devito with EnduraCap Technology per 2.5 gals. of water. Thoroughly soak each mound and a 3 ft. diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. Apply in early morning or late evening hours. You may make applications on a monthly basis to maintain protection of treated areas.</p> <p>Mosquito Control: Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. Apply Devito with EnduraCap Technology at labeled rates in 2 - 5 gals. of water per 1,000 sq. ft.</p>
Bluegrass Billbugs (adult) Black Turfgrass Ataenius (adult) Chiggers Fleas (adult) Grubs (suppression) Hyperodes Weevils (adult) Mole Crickets (nymphs and young adults)	7 mL per 1,000 sq. ft. or 10 fl. oz. per acre	
¹ Not for use on chinch bugs in New York State.		
* Example: To treat the listed turf pests at the 7 mL per 1,000 sq. ft. rate, determine the gallons of dilution per 1,000 sq. ft. needed to adequately cover the turf. At a 5 gallon per 1,000 sq. ft. dilution, take the 7 mL and divide it by the 5 (gallon) to get 1.4 mL to be added per gallon. For a 50 gallon tank, therefore, this would be 1.4 mL x 50 = 70 mL (or 2.5 oz.) in 50 gallons of water.		

Devito with EnduraCap Technology Mixing Chart for Turf Insect Pest Control

Rate of Devito with EnduraCap Technology	Application Rate Per 1,000 Square Feet of Turf				
	2 Gallons	4 Gallons	6 Gallons	8 Gallons	10 Gallons
5 oz. per acre	5.7 oz.	2.9 oz.	1.9 oz.	1.4 oz.	1.2 oz.
10 oz. per acre	11.5 oz.	5.7 oz.	3.8 oz.	2.9 oz.	2.3 oz.
Conversion Rate: 1 fluid ounce (fl. oz.) equals 29 milliliters (mL).					

ORNAMENTALS

Devito with EnduraCap Technology may be used for applications to maintain indoor or outdoor areas where ornamentals are grown. Treat non-turf areas of residential and non-residential landscaped areas around institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields (trees, shrubs, flowers, evergreens, foliage plants and groundcovers). Application rates for ornamentals are lower than structural pest control rates, reflecting that treatment intervals are more frequent.

Applicators must ensure that they are certified in the necessary pesticide certification categories to allow application of **Devito with EnduraCap Technology** away from structures, such as to ornamental plantings. Structural pest control certification categories may limit the distance away from structures for pesticide application. Consult your state extension office or pesticide regulatory officials for further information.

IMPORTANT: Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

NOTE: While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution.

It is advised to pre-spray a selection of ornamental plants and observe them for 7 - 10 days prior to treating large areas if local use experience is unavailable.

Mixing Instructions

Devito with EnduraCap Technology is to be mixed with water and may be used in all types of standard application equipment. Fill applicator tank with the desired volume of water and add **Devito with EnduraCap Technology**. It is suggested that the water be 5 - 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add **Devito with EnduraCap Technology** to applicator tank water with maximum agitation. Close and shake or re-agitate applicator tank before use if application is interrupted. Make up only amount of treatment volume as required.

Tank Mixing

Devito with EnduraCap Technology may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, perform a small volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Devito with EnduraCap Technology** last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Follow all restrictions and precautions which appear on the labels of these products.

RESTRICTIONS:

- **DO NOT** apply more than 0.36 lbs. of the a.i. (52.4 fl. ozs. of concentrate) per acre per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. For soil or foliar applications, **DO NOT** apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- **DO NOT** make outdoor broadcast applications to ornamentals when wind speed is 15 mph or greater.
- **DO NOT** apply to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields, or any other area frequented by the general public).

TANK DILUTION RATES FOR ORNAMENTAL PESTS

PEST	AMOUNT OF DEVITO WITH ENDURACAP TECHNOLOGY	APPLICATION INSTRUCTIONS
Ants (including Imported Fire Ants) Armyworms Azalea Caterpillars Aphids Bagworms Black Vine Weevils (adult) Boxelder Bugs Budworms California Oakworms Cankerworms Cockroaches Crickets Cutworms Eastern Tent Caterpillars Elm Leaf Beetles European Sawflies Fall Webworms Flea Beetles Forest Tent Caterpillars Gypsy Moth Larvae (continued)	1.5 - 5 fl. oz. per 100 gals. or 44 - 148 mL per 100 gals.	Application to ornamentals should be started prior to the establishment of high insect pest populations. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at labeled rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. For spot treatments use 0.5 fl. oz. Devito with EnduraCap Technology per 1 - 2.5 gals. of water. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. Scale: Thoroughly cover the plant with Devito with EnduraCap Technology spray, including trunks, stems, twigs, and foliage for control of scale insects (crawler stage). Bagworm: Apply Devito with EnduraCap Technology when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.

PEST	AMOUNT OF DEVITO WITH ENDURACAP TECHNOLOGY	APPLICATION INSTRUCTIONS
Japanese Beetles (adults) June Beetles (adults) Lace Bugs Leaf-feeding Caterpillars Leafhoppers Leafminers (adults) Leaf Rollers Leaf Skeletonizers Midges Mosquitoes Oleander Moth Larvae Pillbugs Pine Sawflies Pine Shoot Beetles Pinetip Moths Plant Bugs Root Weevils Sawflies Scale Insects (crawlers) Spiders Spittlebugs Stink Bugs Striped Beetles Striped Oakworms Thrips Tip Moths Tussock Moth Larvae Wasps	1.5 - 5 fl. oz. per 100 gals. or 44 - 148 mL per 100 gals.	<p>Application to ornamentals should be started prior to the establishment of high insect pest populations. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at labeled rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. For spot treatments use 0.5 fl. oz. Devito with EnduraCap Technology per 1 - 2.5 gals. of water. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.</p> <p>Scale: Thoroughly cover the plant with Devito with EnduraCap Technology spray, including trunks, stems, twigs, and foliage for control of scale insects (crawler stage).</p> <p>Bagworm: Apply Devito with EnduraCap Technology when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.</p>
Broadmites Brown Softscales California Redscale (crawlers) Clover Mites Mealybugs Pineneedlescales (crawlers) Spider Mites Whiteflies	3 - 5 fl. oz. per 100 gals. or 88 - 148 mL per 100 gals.	

Devito with EnduraCap Technology Mixing Chart for Ornamental Insect Pest Control
(Devito with EnduraCap Technology to add per spray tank)

Rate of Devito with EnduraCap Technology	25 Gallons	50 Gallons	100 Gallons	200 Gallons	300 Gallons
1.5 oz. ¹	0.4 oz.	0.8 oz.	1.5 oz.	3.0 oz.	4.5 oz.
3 oz. ²	0.8 oz.	1.5 oz.	3.0 oz.	6.0 oz.	9.0 oz.
5 oz. ³	1.3 oz.	2.5 oz.	5.0 oz.	10.0 oz.	15.0 oz.

¹ Equivalent to 3.5 mL per 1,000 sq. ft. (5 oz. per acre) when applied at 8 gal. per 1,000 sq. ft.

² Equivalent to 7 mL per 1,000 sq. ft. (10 oz. per acre) when applied at 8 gal. per 1,000 sq. ft.

³ Equivalent to 9.5 mL per 1,000 sq. ft. (14 oz. per acre) when applied at 8 gal. per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used, must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Devito™ with EnduraCap Technology is a trademark of Atticus, LLC.

20230816a

Treats up to 1000 sq ft per Pound

InTice¹⁰™

perimeter bait

Kills Roaches, Crickets, Mole Crickets, Sowbugs, Pillbugs, Earwigs, Silverfish, Millipedes, Snails, Slugs and Ants*

*Except Pharaoh Ants, Fire Ants, Harvester Ants

For Indoor, Outdoor, & Turf Uses
Mold & Moisture Resistant
Ready-to-Use Bait

Active Ingredient: Orthoboric Acid10%

Other Ingredients:.....90%

Total:.....100%

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions: In food areas of restaurants, residential, health care, educational, daycare, agricultural buildings, animal rearing and handling establishments, food handling and food processing establishments, this product may only be applied as a crack and crevice treatment. Any product remaining outside of cracks and crevices must be cleaned up and removed.

Use Sites: In and around residential, multi-family, commercial, industrial, municipal, institutional, research, recreational, health care, educational, daycare, hospitality and agricultural buildings and other man-made structures, garages, home greenhouses, landscape beds, transport vehicles, sewers, animal rearing and handling establishments and food handling and food processing establishments. Turf sites include lawns, ornamental turf, parks, playing fields, right-of-ways, and golf courses. Turf use not allowed in the State of California.

Outdoor Application

Apply bait at a rate of 1 lb (about 3 cups) per 1000 square feet. Apply in a band from two up to four feet wide around the perimeter of buildings. Also apply around exterior light fixtures and along the edges of sidewalks, patios and driveways and under decks. Apply bait early in the season in areas likely to be infested, such as mulch beds, flowerbeds, fruit and vegetable gardens, compost heaps, wood piles, and around trees, stumps and trash areas. The bait may also be sprinkled in and around rodent bait stations to control insects that harbor in these stations. When treating for ants, apply ½ to 1 oz of bait around ant mounds and sprinkle along foraging trails. For ants, pay particular attention to harborage areas and apply bait around trees and stumps, in tree cavities, in and around firewood piles, around landscaping stones and in mulch beds. Bait may also be applied in insect bait stations. Reapply bait after one week if infestation is still present. In home greenhouses, apply bait at a rate of 3 oz (2/3 cups) per 100 sq ft on and under benches. Do not apply bait directly in flats or pots. Reapply as needed.

For Carpenter Ants: Evenly apply bait at a rate of 14 oz by weight (3 cups) per 100 sq ft. Apply in a band not to exceed one foot in width around trees and stumps, firewood piles, structure foundations and other potential nest sites. The bait may also be applied in tree cavities and firewood piles, and under porches and decks. Do not apply more than 2 lbs per 1000 sq ft of total site area, or 90 lbs per acre.

Sewer Application: Apply bait along ledges inside sewers and around manhole covers.

Indoor Application

Apply bait only in areas inaccessible to children and pets, or in tamper-resistant bait stations. Apply bait at a rate of 3 oz (2/3 cup) per 100 square feet. Using a hand shaker or duster, apply bait in attics, crawl spaces, trash closets, utility closets, dropped ceilings, on dirt or gravel cellar floors, under and behind appliances and vending machines, in wall voids and voids under and behind cabinets, sinks and tubs, in drawer wells, garbage chutes, pipe chases, elevator pits, around drains, around electrical conduits and in cracks and crevices where insects may harbor. For crack and crevice treatments, use a duster and apply 1 to 3 puffs per linear foot based on level of infestation. Focus application in areas where insects or their signs are seen. Reapply bait after one week if infestation is still present. A power duster may be used to apply bait in appropriate areas, such as attics and crawlspaces. The duster manufacturer's recommendations for settings should be followed.

Application tips for certain sites follow:

1. In attics, be sure to get bait near the eaves and vent pipes where insects often first enter, as well as around any pipes or potential access points between the attic and the main structure. Attics

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals:

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. In the State of California: Personal Protective Equipment (PPE) Applicators and handlers must wear: shoes plus socks.

First Aid

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product or label with you when calling a Poison Control Center or doctor or going for treatment. In case of emergency, for additional information call 800 858 7378.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

EPA Reg. No.: 73079-6

EPA Est. No.: 73079-MO-1

Net Contents: 1, 4, 10, 40 lbs



should have proper ventilation to insure that no food odor from the bait seeps into the living area, particularly in hot weather. If in doubt, apply the product in a manner such that it can be easily removed.

2. When treating cabinetry areas, if there is no gap between the lower cabinets and the kickplate, then small holes can be drilled to access the voids. Escutcheon plates around pipes can also be pulled back to allow access to voids.

3. The bottom drawers in kitchen and bathroom cabinets can be removed and bait dusted into drawer wells. Do not leave any bait exposed when drawers are replaced.

4. Electrical switch plate covers can be removed to allow access to voids, but bait must not be applied directly in electrical boxes.

5. If there are cracks between baseboards, cabinets, doorframes, hinges, counters or trim, and walls, these cracks can be lightly dusted with bait. If practical, these areas can be sealed once insect problems have been eliminated, to help prevent future problems.

6. Insects need moisture, so kitchen and bath areas and areas immediately surrounding them, and any other damp areas should be carefully inspected and treated if necessary. As a separate matter from the use of this product, it is advisable to fix any leaks to minimize structural damage and odors, and to prevent potential chronic insect problems.

To help prevent insect problems from developing in homes and buildings, a preventative program is recommended. Once per quarter, apply bait in a band around the outside building perimeter. Once per year, apply bait in attics and crawlspaces. Follow Indoor/Outdoor Application Directions above.

Food Processing and Handling Establishment Application, including Federally-

Inspected Meat and Poultry Plants: Follow instructions for other sites as appropriate. In food areas, only apply bait in cracks, crevices, and other inaccessible areas. Do not contaminate food and feedstuffs. Never apply bait directly to a surface where food is stored, prepared or served. Any bait that is left visible after treatment must be swept into cracks and crevices or removed.

Turf Application (Except in the State of California)

Apply bait evenly over the affected area at a rate of 1 lb per 1000 sq ft or 45 lbs per acre. Do not water the bait in. For mole cricket treatments, results will be best if the area is irrigated prior to bait treatment. Mole crickets typically come to the surface to feed in the evening.

Storage and Disposal

For 10 lb and 40 lb containers:

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place. Do not store where children or animals may gain access.

Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP which is managed as a public-private partnership.

Or, for 1 lb and 4 lb containers:

This storage/application container may be refilled by the end-user of the product. Refill only with InTice 10 Perimeter Bait.

Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP which is managed as a public-private partnership.

Green Zone is not a third party certification.

doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas. **BED BUGS:** Apply SPECTRE PS Insecticide to cracks, crevices, voids and/or shelters where evidence of bed bugs occurs, including bed frames, box springs, inside empty dressers, furniture and clothes closets, carpet edges, high and low wall moldings, behind headboards, pictures and wallpaper edges. Do not make surface applications to mattresses or bedding. Mattress treatments must be limited to seams, folds and edges only. Do not treat pillows, bedding or clothing. Remove linens and wash before reuse. Allow treated mattress to dry thoroughly before remaking the bed. Remove all clothes and other articles from dressers or clothes closets before application. Allow treated areas to dry thoroughly before use. **BOOKLICE** and **SILVERFISH:** Apply SPECTRE PS Insecticide as an Injection treatment into cracks and crevices of baseboards, door frames, bookcases and other harborage sites. Also make spot treatments to areas such as storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves and drawers. **CRAWLING PESTS** [including Centipedes, Cockroaches, Crickets, Dermestid Beetles, Earwigs, Sowbugs/Pillbugs and Spiders (excluding Brown Recluse)]: Apply SPECTRE PS Insecticide as an Injection treatment using the prescribed amount into cracks, crevices and void spaces where pests may rest, nest, trail, run or hide, including openings around pipes and sinks, under refrigerators, behind baseboards, washing machines, stoves, cabinets, sewers, floors drains and meter boxes. Also make spot treatments to areas such as storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves and drawers. **CHOCOLATE MOTHS, DRUGSTORE BEETLES, FLOUR BEETLES, GRAIN WEEVILS, INDIANMEAL MOTHS and TROGODERMAS** (in exposed adult and larval stages): Apply SPECTRE PS Insecticide into cracks, crevices and/or voids where these pests may harbor such as cracks and crevices in cabinets and pantries, wall voids, around window and door frames, and in machinery. Also make spot treatments to areas such as storage areas; closets; around water pipes, doors and windows; around refrigerators, cabinets, sinks, stoves and other equipment; shelves, drawers and similar areas. **DRAIN FLIES:** Release approximately 1 to 2 sec of SPECTRE PS Insecticide into the drain basin. **INTERIOR/LIMITED EXTERIOR APPLICATION OF STRUCTURES: FOR ACTIVE AND HIBERNATING STAGES OF ASIAN LADY BEETLES, BOXELDER BUGS, CLUSTER FLIES, ELM LEAF BEETLES, HONEY BEES, STINK BUGS, WASPS AND YELLOW JACKETS:** Apply SPECTRE PS Insecticide as an Injection or spot treatment into cracks, crevices and/or voids where these insects may harbor, congregate or hibernate such as cracks and crevices, behind siding, wall voids, around window and door frames. **CARPENTER BEES:** Locate entrance holes. Treat galleries through entrance holes with SPECTRE PS Insecticide. Seal bee entrances after application. **CRAWLING INSECTS** (including Centipedes, Crickets, Earwigs, Pillbugs/Sowbugs and Springtails): Apply as an Injection or spot treatment into cracks and crevices where insects may enter premises or harbor such as weep holes, under siding, wall voids, soffits, around attic vents, cracks, crevices and openings around windows doors and pipes, and exterior areas where utilities may enter the structure. **FLIES (House and Stable):** Apply SPECTRE PS Insecticide as an Injection or spot treatment to areas where flies rest, congregate or enter the structure on both the exterior and interior. Apply around doors, windows, garages, garbage storage areas, upper corners of walls, ceiling areas, around sky lights, light fixtures and similar areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store this product in a cool, dry (preferably locked) place away from heat or open flame, in an area that is inaccessible to children and animals. **PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Do not puncture or incinerate! If Empty: Place in trash or offer for recycling is available. If partly filled: Call your local solid waste agency for disposal instructions.

LIMITATION OF WARRANTY AND LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc. election, the replacement of product.

SPECTRE PS Insecticide is not manufactured or distributed by BASF Corporation, seller of PT® Phantom® II. PT® and Phantom® II are trademarks of BASF Corporation.



SPECTRE® PS INSECTICIDE

For sale to, use and storage only by individuals/firms licensed or registered by the state to apply termiticide and/or general pest control products.

SPECTRE PS Insecticide is a pressurized product for localized use in and around apartments, food storage area, homes, hospitals, hotels, meat packing and food processing plants, motels, nursing homes, restaurants and other food handling establishments, schools, supermarkets, warehouses and other commercial and industrial buildings. Kills many nuisance pests, stored product pests and occasional pests that infest structures, including: Bed bugs, Ants, Beetles/Weevils, Bees/Wasps, Flies, Drain Flies, Cockroaches, Crickets, Earwigs, Indian meal moths, Silverfish, Springtails and Spiders (excluding Brown Recluse).

Active Ingredient:

Chlorfenapyr¹ 0.5%

Other Ingredients:

Total: 100.0%

¹4-bromo-2-[4-(chlorophenyl)-1-(ethoxymethyl)-5-(trifluoromethyl)-1H-pyrrrole-3-carbonitrile.

Contains chlorfenapyr, the active ingredient used in PT® Phantom® II. SPECTRE PS is not manufactured or distributed by BASF Corporation, seller of PT® Phantom® II.

EPA Reg. No.: 91234-23-53883

EPA Est. No: 53883-TX-002

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See side panel(s) for Precautionary Statements, Directions for Use, Storage and Disposal and Conditions of Sale and Warranty.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may also call 1-844-685-9173.

For Chemical Emergency Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers must wear baseline PPE consisting of (1) long sleeved shirt, (2) long pants and (3) shoes plus socks. **USER SAFETY REQUIREMENTS:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove and wash contaminated clothing before reuse.

USER SAFETY RECOMMENDATIONS

Users should (1) wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet, (2) remove contaminated clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing, and (3) remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to aquatic organisms, birds and wildlife. Do not apply directly to water or to areas where surface water is present or intertidal areas below the mean high water mark.

PHYSICAL OR CHEMICAL HAZARDS: Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray directly into any electronic equipment such as radios, televisions, computers, etc. due to the possibility of short circuits or shock hazard. Test on a small surface prior to use, as spray may mar plastic, painted or varnished surfaces.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Product Information: SPECTRE PS Insecticide is a pressurized product for localized use in and around apartments, food storage area, homes, hospitals, hotels, meat packing and food processing plants, motels, nursing homes, restaurants and other food handling establishments, schools, supermarkets, warehouses and other commercial and industrial buildings. SPECTRE PS Insecticide kills many nuisance pests, stored product pests and occasional pests that infest structures, including:

- a. **Ants:** Foraging fire ants, Harvester ants, and Pharaoh ants.
- b. **Beetles/Weevils:** Asian lady beetles, Dermestid beetles, Drugstore beetles, Elm leaf beetles, Flour beetles, Grain weevils and Khapra beetles.
- c. **Bees/Wasps:** Carpenter bees, Honey bees, Wasps, Yellow jackets.
- d. **Bugs:** Bed bugs, Box-elder bugs, Stink bugs.
- e. **Flies:** Cluster flies, Drain flies, House flies, Stable flies.
- f. **Other Insects Pests:** Booklice, Chocolate moths, Cockroaches, Crickets, Earwigs, Indian meal moths, Silverfish, Springtails.
- g. **Non-Insect Pests:** Centipedes, Pillbugs/Sowbugs and Spiders (excluding Brown Recluse).

USE RESTRICTIONS: Do not apply SPECTRE PS Insecticide in any manner other than specified on this label. •This product is not registered for use as a termiticide. •Do not compost an discarded plant materials that have been treated with SPECTRE PS Insecticide. •If it is raining, do not make outdoor applications. •Do not reapply to treated areas more than once every four (4) weeks. •Do not allow adults, children or pets to enter until vapors, mists and/or aerosols have dispersed and the treated area has been thoroughly ventilated. •Do not contact treated surfaces until dry. •Do not apply to school classrooms while in use. Allow treated areas to dry thoroughly before reoccupancy. •Remove patients in hospitals and nursing homes before treating rooms. Allow treated areas to dry thoroughly before returning patients to rooms.

•Remove or cover exposed food and drinking water in home treatment areas before application. Remove or cover dishes, utensils, food processing equipment and food preparations surfaces in the treatment area or wash them before use. Do not use in servicing areas while food is exposed. •Applications in food areas of food handling establishments are limited to spot and/or injection treatments as described below. Do not make applications when facility is in operation. Avoid contamination of food or feedstuffs. Do not apply directly to food, food packaging or food contact surfaces. During application, cover or remove food and cover food processing surfaces and equipment, or clean after treatment with a suitable detergent and rinse with potable water before use. Do not use in serving areas (dining areas where prepared foods are served) while food is exposed or facility is in operation. •Except when applying to sewers or drains as instructed on this label, do not apply directly into sewers or drains or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. •When used in zoos, remove animals and cover or remove animal food/water prior to application. •Remove pets and cover or remove pet food/water and fish bowls prior to application.

INJECTION TREATMENT: Use the supplied actuator and injection tubes to apply SPECTRE PS Insecticide. Inject into cracks and crevices or void spaces where pests may be hiding, living and breeding. After placing the injector tip into cracks, crevices, holes and other small openings, release approximately 1 sec of SPECTRE PS Insecticide. For light infestations, move injector tip along cracks and crevices or void spaces while treating at the rate of 3 Linear f/usc. For heavy infestations, move the injector tip along at 1 Linear f/usc. For closed voids, figure the cubic area of the void and apply SPECTRE PS Insecticide at 1 to 5 sec/3lit. Several holes may be required in long running voids. **SPOT TREATMENT:** Use the supplied actuator and injection tubes to apply SPECTRE PS Insecticide. Always hold the container upright while spraying and direct the nozzle about 12 inches from the surface being treated. Apply SPECTRE PS Insecticide directly on insects in these locations when possible. Limit individual spot treatments to an area no larger than 2 f2/room. SPECTRE PS could stain some surfaces or damage finishes. Prior to spot treating or plastic, painted or varnished surfaces, test a small, inconspicuous area and observe for staining or damage to finish. **FOOD HANDLING ESTABLISHMENTS:** Food handling establishments are places other than private residences in which food is held, processed, prepared or served. **FOOD AREAS:** Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) are also considered food areas. Apply as an Injection treatment or spot treatment directly into cracks, crevices, voids or onto surfaces where pests rest, nest, trail, run or hide. Deliver a pin-stream of SPECTRE Insecticide through an injection tube suitable for that purpose. Apply to different elements of construction: between equipment and floor; openings leading to hollow spaces in walls, equipment legs and bases; and where insects hide or nest. Limit individual spot treatments to an area no larger than 2 f2/room. **NON-FOOD AREAS:** Includes areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling). Apply SPECTRE Insecticide as an Injection treatment or spot treatment to selected surfaces such as baseboards and under elements of construction. **INDOOR TREATMENTS: ANTS** (including Foraging fire ant, Harvester ant and Pharaoh ant): Apply SPECTRE PS Insecticide as an Injection treatment into cracks, crevices and voids where ants rest, nest, trail, run or hide, including around doors, window frames, wall voids and other structural cracks and crevices. Apply SPECTER PS Insecticide as spot treatments to areas such as storage areas; closets; around water pipes,

NET CONTENTS: 15 OZ.