

Chickweed Clover & Oxalis Killer

TRUSTED SINCE 1926

BONIDE[®]

Concentrate



One pint covers 5,000 sq. ft.

Also kills speedwell, veronica, ground ivy
& many other broadleaf weeds

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid* 13.72%

Triethylamine Salt of 3, 5, 6-Trichloro-2-Pyridinyloxyacetic Acid** 1.56%

Dimethylamine Salt of Dicamba (3, 6-Dichloro-o-Anisic Acid)*** 1.35%

OTHER INGREDIENTS: 83.37%

TOTAL: 100.00%

See back panel for substatement of ingredient statement.

EPA No. 4-NY-1 EPA Reg. No. 228-334-4

Keep Out Of Reach Of Children

WARNING (See back panel for First Aid and
Additional Precautionary Statements)



Chickweed

Clover

Oxalis

Chickweed Clover & Oxalis Killer Concentrate

- ✓ Kills chickweed, clover, dandelions, ground ivy, oxalis, wild violet, and other tough weeds
- ✓ One pint covers 5000 sq. Ft.
- ✓ Contains mcpa, triclopyr and dicamba

By Isomer Specific Method, Equivalent to:

| | |
|--|-------------------------|
| *2-Methyl-4-Chlorophenoxyacetic Acid..... | 11.20%, 0.970 lbs./gal. |
| **3,5,6-Trichloro-2-Pyridinyloxyacetic Acid..... | 1.12%, 0.097 lbs./gal. |
| ***3,6-Dichloro-o-Anisic Acid..... | 1.12%, 0.097 lbs./gal. |

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

Store and transport in an upright position
EPA Reg. No. 228-334-4 EPA Est. No. 4-NY-1
Buyers Guarantee Limited to Label Claims.
©Bonide Products, Inc. All Rights Reserved

BN

Distributed by:
Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is for use on residential turf. **IMPORTANT:** For use on Bluegrass, Fescues, Rye, Bent, Bermuda, Bahia and Zoysia Lawns. Avoid application to St. Augustine or centipede lawns unless injury can be tolerated. Do not spray Carpet grass, Dichondra or desirable clovers. May cause temporary yellowing of some Bermudagrass turf. Do not exceed specified dosages for any area. Be particularly careful applying within the drip line of trees and other ornamental species. Avoid contact with exposed feeder roots of ornamentals and trees.

Do not apply more than 1.5 lb ae/acre per year. Do not apply more than 2 broadcast applications per year with a minimum retreatment interval of 21 days.

GENERAL APPLICATION RESTRICTIONS

Entry Restrictions: Do not apply this product in a way that will contact any person, or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

LAWN WEED CONTROL

When used as directed kills broadleaf weeds in lawns - roots and all. Kills weeds - not lawn grasses.

ORNAMENTAL LAWNS AND TURF

Apply this product at the rate of 2 pints in 4 to 50 gallons of water per 10,000 square feet (3.2 fluid ounces in 2.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to Bluegrass, Fescue, Rye, Bent (excluding golf course greens and tees), Bahia, Bermuda and Zoysia.

FOR BEST RESULTS

Before You Apply:

Do not mow for 1 to 2 days before or after application.

If soil is dry, water before application.

To treat entire lawn (broadcast application): Use a tank sprayer or hose-end sprayer. Measure lawn. Calculate square feet by multiplying length times width. Spray evenly over measured area.

HOW TO APPLY

To spot treat individual lawn weeds: Use a tank sprayer and spray to wet weeds. To reduce spray drift, adjust sprayer nozzle to deliver a coarse spray.

WHEN TO APPLY

- Spray when weeds are actively growing.
- Spray when temperature is below 90°F.
- Spray when air is calm to avoid drift to vegetables, flowers, ornamental plants, trees, shrubs and other desirable plants.
- Do not apply to newly seeded grasses until well-established (they have been mowed 3 times).
- Rain or watering 1 hour after application will not wash away effectiveness.
- Reseed no sooner than 3 weeks after the application.

WEEDS

Alder, Annual yellow sweet clover, Artichoke, Aster, Austrian fieldcress, Bedstraw, Beggartick, Bide, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Black-seed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbuttons, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinquefoil, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullein, Common sowthistle, Corn Chamomile, Creeping jenny, Crimson clover, Croton, Cudweed, Curly dock, Curly indigo, Dandelion, Dead nettle, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Fall dandelion, False dandelion, False flax, False sunflower, Fiddleneck, Field bindweed, Field pansy, Flea bane (daisy), Flixweed, Florida betony, Florida pusley, Frenchweed, Galinsoga, Garlic mustard, Goathead, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hairy bittercress, Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Honeysuckle, Hop clover, Horsenettle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, Lupine, Mallow, Marshelder, Matchweed, Mexicanweed, Milk vetch, Milkweed bloodflower, Mugwort, Morningglory, Mouseear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf vetch, Nettle, Orange hawkweed,

WEEDS Cont.

Oxalis, Oxeye daisy, Parsley-piert, Parsnip, Pearlwort, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Pineywoods bedstraw, Plains coreopsis, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem filaree, Rough cinquefoil, Rough fleabane, Roundleafed marigold, Rush, Russian pigweed, Russian thistle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spiny Amaranth, Spiny cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Tarweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Tumble mustard, Tumble pigweed, Tumbleweed, Velvet leaf, Venice mallow, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Western salsify, White clover, White mustard, Wild mustard, Wild aster, Wild buckwheat, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Wild vetch, Willow, Witchweed, Woolly morningglory, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed and other broadleaf weeds.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: To be stored in original container and placed in an area inaccessible to children.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. **If empty** - Place in trash or offer for recycling if available. **If partly filled** - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

FIRST AID

| | |
|-------------------------------|---|
| IF IN EYES | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person. |
| IF ON SKIN OR CLOTHING | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice. |

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for medical emergency information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING-AVISO: Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. This product may cause skin sensitization reactions in some people.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith.