

# EVOLVE™

## MOUSE

### A soft bait to reduce fertility in mice.

**Active Ingredients:**

Cottonseed Oil .....0.1%

**Other Ingredients:** ..... 99.9%

Oat Flour, Palm Oil, Hydrogenated

Soybean Oil, Cottonseed Meal,

Sucrose, Citric Acid

**Total:** 100.0%

**FIRST AID**

**If on skin:** Take off contaminated clothing. Wash with plenty of soap and water.

**If in eyes:** Rinse cautiously with water for several minutes.

**If ingested:** Do not induce vomiting. Call a poison center or doctor if you feel unwell.

**PRECAUTIONARY STATEMENTS****Hazards to Humans and Domestic Animals**

**CAUTION.** Contact with eyes may cause irritation. Wash hands thoroughly after handling.

**Personal Protective Equipment:**

Wear protective gloves and clothes. Safety glasses recommended.

**Treatment for Pet Ingestion:**

If animal consumes bait and adverse effects are observed, call a veterinarian.

### Keep Out of Reach of Children

## CAUTION

**Net Contents:** 1-40 lb. (0.5-18 kg)

**Contains:** approx. 26-1,037 pieces

**DIRECTIONS FOR USE**

READ THIS LABEL: Follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

USE RESTRICTIONS: **1.** Bait must not be deployed in areas where contamination of food, water, or surfaces that come in direct contact with food is possible. **2.** Do not dilute or add anything to the product. **3.** Store and deploy bait out of reach of children, pets, domestic animals, and non-target wildlife or in tamper-resistant bait stations. **4.** Dispose of product container, unused, and unconsumed bait as specified on this label.

SELECTION OF TREATMENT AREAS: This product is for use in commercial, residential, and agricultural sites. Determine areas where mice are most likely to find and consume bait. These areas can include concealed areas along walls, near burrows, near structural openings in the floor, wall, ceiling, or in hard to reach areas (e.g., in rafters, beams or attics) or where other signs of mouse activity have been observed. Reassess activity for changes in placement (e.g., due to seasonal changes or construction). Multiple bait stations may be placed in foraging areas or high activity locations as needed.

APPLICATION DIRECTIONS: **1.** Place 2 to 12 pieces per placement. **2.** Adjust the amount of bait applied at each placement based upon mouse activity levels. **3.** Maintain an uninterrupted supply of bait. Check and replace bait as necessary.

**STORAGE AND DISPOSAL**

**Storage:** Store away from water. Store only in original, tightly secured containers in a room temperature environment, inaccessible to unauthorized users, children, pets, livestock, and other non-target animals. **Disposal:** Place product packaging, unused bait, or unconsumed bait in trash.

**DISCLAIMER**

To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

This product has not been registered by the U.S. Environmental Protection Agency. SenesTech, Inc. represents that this product qualifies for exemption from registration as a minimum risk pesticide under FIFRA Section 25(b).

© 2024 SenesTech, Inc. US Patent Pending. Evolve™ is a trademark of SenesTech, Inc.

**Manufactured by:** SenesTech, Inc.  
777 W Pinnacle Peak Rd, Suite B104  
Phoenix, AZ 85027  
1-866-886-7287  
senestech.com



SCAN TO VIEW  
PRODUCT  
INFORMATION

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor immediately for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor immediately for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>TREATMENT FOR PET POISONING:</b> If animal eats bait, call veterinarian at once.	
<b>NOTE TO PHYSICIAN OR VETERINARIAN:</b> Contains the anticoagulant chlorophacinone. If swallowed, this material may reduce the clotting ability of blood and cause bleeding. For humans or animals ingesting bait and/or showing obvious poisoning symptoms (bleeding or prolonged prothrombin times), give Vitamin K <sub>1</sub> intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.	

## PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Harmful if absorbed through the skin. Avoid contact with eyes, skin or clothing. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear long pants, shoes, socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet, and change into clean clothing.

**ENVIRONMENTAL HAZARDS:** This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**Pesticide Storage:** Store in original container in a cool, dry place inaccessible to children and pets.  
**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of in trash or at an approved waste disposal facility.  
**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty container, then dispose of empty container in trash or at an approved waste disposal facility.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

# FLATLINE™

## SOFT BAIT

FLATLINE IS ATTRACTIVE TO RODENTS  
 FOR INDOOR AND OUTDOOR USE  
 KILLS NORWAY RATS, ROOF RATS, AND HOUSE MICE  
 KILLS RATS, MICE, DEER MICE, AND MEADOW VOLES\*  
 KILLS OTHER RODENT SPECIES AS LISTED ON THIS LABEL  
 CONVENIENT AND EASY-TO-USE  
 HELPS STOP DAMAGE CAUSED BY RODENTS

Active Ingredient: Chlorophacinone ..... 0.005%  
 Inert Ingredients ..... 99.995%  
 Total ..... 100.000%

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION** See side panel for additional precautionary statements.

**LIPHA**TECH™

Liphatech, Inc.  
 3600 W. Elm Street  
 Milwaukee, WI 53209  
 800-351-1476

This bait is packaged in 1.0 ounce  
 (28 grams) individual pouches

\*Use not permitted in California

Do not sell this product in individual containers  
 holding less than 4 pounds of bait.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
**READ THIS LABEL** and follow all use directions and precautions. Use only for the sites, pests, and application methods described on this label.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals, and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

**USE RESTRICTIONS:** This product can only be used to control Norway rats, roof rats, house mice, Deer mouse (*Peromyscus maniculatus*), White-footed mouse (*Peromyscus leucopus*), Cotton mouse (*Peromyscus gossypinus*), Cotton rat\* (*Sigmodon hispidus*), Eastern harvest mouse\* (*Reithrodontomys humuli*), Golden mouse\* (*Ochrotomys nuttalli*), Polynesian rat\* (*Rattus exulans*), Meadow vole\* (*Microtus pennsylvanicus*), White-throated woodrat\* (*Neotoma albigula*), Southern plains woodrat\* (*Neotoma micropus*) and Mexican woodrat\* (*Neotoma mexicana*) in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings, and related structures around and associated with these sites. Fence and perimeter baiting beyond 100 feet from a structure, as defined above, is prohibited. Do not sell this product in individual containers holding less than 4 pounds of bait. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feed, food or feed handling equipment, or milk or meat handling equipment. Do not apply directly to food or feed crops. Do not broadcast bait. Burrow baiting with this product is prohibited.

**Selection of Treatment Areas:** Determine areas where rats, mice, and/or meadow voles\* will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and concealed places, between floors or walls, beside burrows, or in locations where signs of rats, mice, and/or meadow voles\* have been observed. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:** Each pouch contains 1.0 oz. (28 g). Bait may be removed from the pouch and placed in a feeding device that secures the unwrapped bait in a bait station. Discard empty wrapper in the trash.

**Mice and Meadow Voles\*:** Apply 1 pouch per placement, usually spaced 8 to 12 feet apart. Up to 2 pouches may be needed at points of very high activity. Maintain a constant supply of fresh bait for 15 days or until signs of activity cease.

**Rats:** Apply 3 to 16 pouches per placement, usually spaced 15 to 30 feet apart. Maintain a constant supply of fresh bait for 10 days or until signs of rat activity cease.

**Follow-up:** Replace contaminated or spoiled bait. Wearing waterproof gloves, collect and dispose of all visible dead animals and leftover bait, as well as any bait that has become accessible to nontarget animals. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation does occur, repeat treatment. For a continuous infestation, set up permanent bait stations and replenish bait as needed. (062619)

EPA Reg. No. 7173 - 308  
 EPA Est. No. 7173-WI-1