

Devito™



For Both Indoor and Outdoor Use.

For use in, on and around buildings and structures for the control of listed pests, including on non-residential lawns, ornamental trees and shrubs around residential, institutional, public, commercial, agricultural and industrial buildings.

ACTIVE INGREDIENT:

(% by weight)

Lambda-cyhalothrin*: [1 α (S*), 3 α (Z)]-(\pm)-cyano-(3-phenoxyphenyl) methyl-3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethylcyclopropanecarboxylate. 9.7%

OTHER INGREDIENTS: 90.3%

TOTAL 100.0%

*Synthetic pyrethroid

EPA Reg. No.: 91234-250

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	<ul style="list-style-type: none">▪ Call a poison control center or doctor immediately for treatment advice.▪ Have person sip a glass of water if able to swallow.▪ Do not induce vomiting unless told to do so by the poison control center or doctor.▪ Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. To protect the environment, **do not** allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run-off to water bodies or drainage systems. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protect pollinating insects by following label directions intended to minimize drift and to reduce risk to these organisms.

PHYSICAL AND CHEMICAL HAZARDS

Do not use this product in or on electrical equipment due to the possibility of shock hazard. **Do not** mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

PRODUCT INFORMATION

Devito with EnduraCap Technology is a unique formulation utilizing EnduraCap™ Technology which is a proprietary blend of ingredients for use with the active ingredient Lambda-cyhalothrin.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For both indoor and outdoor use.

IMPORTANT: Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on lawns or grounds.

This product is RESTRICTED for use in New York State.

Application Restrictions for Residential Outdoor Surface and Space Sprays:

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

- **Do not** apply or irrigate to the point of runoff.
- **Do not** make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).
- Rainfall within 24 hours after application may cause unintended runoff of pesticide application.
- **Do not** apply the product into fish pools, ponds, streams, or lakes. **Do not** apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

- **Do not** allow to enter indoor or outdoor drains unless labeled for drain treatments.

No permita la entrada a desagües internos o externos a menos que el etiquetado indique que está permitido el uso del producto para tratamiento de desagües.



- Follow proper disposal procedures on this label.

Siga las indicaciones del etiquetado para el desecho apropiado del producto.

- **Do not** allow the product to enter any drain during or after application except where labelled for this use.
- **Do not** allow children or pets to contact treated surfaces until spray has dried.
- Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).
- **Do not** apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Treat surfaces to ensure thorough coverage but avoid runoff. To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

PERIMETER PEST CONTROL

Devito with EnduraCap Technology is for use as a general surface (nonfood/nonfeed areas), crack and crevice, or spot treatment in, on, and around buildings and structures and their immediate surroundings, and on modes of transport. Areas of use: industrial buildings, houses, patios and porches (spot and crack and crevice treatment only), closets, furniture, apartment buildings, mobile homes, laboratories, buses, greenhouses (non-commercial), stores, factories, warehouses, wineries, vessels, railcars, trucks, trailers, aircraft (cargo and other non-cabin areas only), schools, nursing homes, hospitals, mausoleums, restaurants, hotels, correctional facilities, livestock/poultry housing, pet kennels, food granaries, food grain mills and food manufacturing, processing, and servicing establishments.

Nonfood/nonfeed areas of food/feed handling establishments: garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling).

For indoor applications, retreat at 21-day intervals to maintain control. More frequent treatments are limited to spot treatments. Do not apply to carpets as a pin stream or coarse spray treatment.

Mixing Instructions

Dilute **Devito with EnduraCap Technology** with water for application using hand-held or power-operated application equipment as a coarse spray for crack and crevice or spot and general surface (nonfood/nonfeed areas only) treatments. Application equipment that delivers low volume treatments (such as the Micro-Injector® or Actisol® applicator) may also be used to make crack and crevice or spot and general surface treatments. Fill applicator tank with the desired volume of water and add **Devito with EnduraCap Technology**. Close and shake before use in order to ensure proper mixing. Shake or re-agitate applicator tank before use if application is interrupted. Mix only amount of treatment volume as required. **Devito with EnduraCap Technology** may be applied by using a paintbrush or other porous applicator attached to a handle as a general surface treatment.

Tank Mixing

Devito with EnduraCap Technology may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, perform a small volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Devito with EnduraCap Technology** last. If mixed with EC formulations, use within 24 hours. Fill tank to desired volume and continue to agitate while making applications.

Devito with EnduraCap Technology may be tank mixed with an Insect Growth Regulator (IGR). It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Foam Applications

Devito with EnduraCap Technology may be converted to a foam and the foam used to treat structural voids to control or prevent pests including ants, bees, termites (above ground only), wasps, or other listed arthropods harboring in walls, under slabs, or in other void areas.

RATES FOR PESTS (HAND APPLICATION EQUIPMENT)

PESTS	DILUTION RATE	SPECIFIC USE DIRECTIONS
Ants Bedbugs (adult) Bees Boxelder Bugs Carpenter Bees Carpet Beetles Centipedes Cigarette Beetles Clover Mites Cockroaches ¹ Confused Flour Beetles Crickets Earwigs Firebrats Fleas ² Flies Lesser Grain Borers Millipedes Mosquitoes Red Flour Beetles Rice Weevils Saw-toothed Grain Beetles Silverfish Sow Bugs Spiders Stink Bugs Ticks Wasps	0.2 fl. oz. (6 mL) to 0.4 fl. oz. (12 mL) per gallon of water (0.015 - 0.03% AI)	<p>Ants: Apply to any trails around doors and windows and other places where ants may be found. Locate and treat nests. Where ants are trailing inside, apply as a residual surface treatment to active areas: baseboards, corners, around pipes, in and behind cabinets, behind and under refrigerators, sinks, furnaces and stoves, cracks and crevices. When combining baits and residual surface insecticides, apply surface insecticides in cracks and crevices, along baseboards, and infested surfaces and outside barrier treatments. Use baits in other areas that are untreated by residual insecticides; also see Outdoor Surfaces Use.</p> <p>Bedbugs: Clean floors and surfaces by vacuuming. Apply as a coarse, low-pressure spray to harborage areas: crevices, baseboards, loose plaster, behind bed frames and headboards, beneath beds and furniture, and to bedsprings and bed frames.</p> <p>Do not apply to furniture surfaces or mattresses where people will be laying or sitting. Infested bedding should not be treated; remove bedding, place in sealed plastic bags, and take for laundering and drying at high temperatures.</p> <p>Cockroaches, Crickets, Earwigs, Firebrats, Silverfish, Spiders and Stink Bugs: Apply as a coarse, low pressure spray to areas where these pests hide: baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, cabinets, behind and under refrigerators, sinks, furnaces and stoves, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices; also see Outdoor Surfaces Use.</p> <p>Bees, Flies, Mosquitoes, and Wasps: Apply directly to walls, ceilings, window screens, and other resting areas as a residual surface treatment. May be used inside residential buildings as well as in and around carports, garages, and storage sheds; also see Outdoor Surfaces Use. Use caution when treating nests of stinging insects as Devito with EnduraCap Technology does not provide instant knockdown. Protective equipment for the applicator against stinging pests may be required.</p> <p>Treat bee, wasp and hornet nests late in the day when most insects will be present. Allow 2 - 3 days for the colony to die and repeat after 21 days, if necessary.</p> <p>Yards or other frequented areas enclosed by landscaping can benefit from the creation of a mosquito barrier to reduce invading mosquitoes by the treatment of perimeter vegetation. Apply Devito with EnduraCap Technology at specified rates in 2 - 5 gals. of water per 1,000 sq. ft.</p> <p>Carpenter Bees: Apply coarse spray to thoroughly wet wood surfaces where bees have been previously active or to provide protection against further damage. Apply early in the spring to prevent bees from invading wood. When bees have infested wood, surface applications can help control embedded larvae and bees that emerge from the wood.</p> <p>Pantry Pests (i.e., Carpet beetle, Cigarette beetle, Confused flour beetle, Lesser grain borer, Red flour beetle, Rice weevil, and Saw-toothed grain beetle): Apply to cupboards, shelving, and storage areas. Remove all utensils, uncovered foodstuffs (or any having original package opened), and shelf paper before making application. Allow treated surfaces to dry and cover shelves with clean paper before replacing any utensils, foodstuff, or other items. Destroy any food-stuff accidentally contaminated with treatment solution.</p> <p>Boxelder Bugs, Centipedes, Millipedes, Pillbugs, and Sow Bugs: Apply around doors and windows and other places where these pests may be found or where they may enter premises.</p> <p>Treat baseboards, storage areas, and other locations listed on this label. Apply treatments to prevent infestation as described below; also see Outdoor Surfaces Use.</p> <p>Fleas and Ticks: To control nuisance fleas and ticks (e.g. dog ticks) apply to kennels, yards, runs, and other areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent.</p> <p>Treat entire area rather than making spot treatments, and retreat as necessary to maintain control, as permissible on this label.</p> <p>Do not apply to pasture or cropland, and do not allow animals and people access to treated areas until the deposit has dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks; also see Outdoor Surfaces Use. Do not apply directly to animals.</p> <p>Cluster Flies: Apply in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors, paying particular attention to south-facing surfaces, but do not allow runoff to occur. Heavy precipitation prior to frost may require retreatments to maintain protection. In winter and spring when flies become active and are emerging, interior crack and crevice and void treatments can help reduce the infestation, along with ULV or general surface application in infested attics or unoccupied lofts.</p> <p>Litter Beetles (Darkling, Hide, and Carrion Beetles) and Flies In Animal Housing (Such As Poultry Houses): To control adult litter beetles, apply Devito with EnduraCap Technology to walls and floors at cleanout, before reintroduction of animals. This will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to areas where beetles frequently occur, walls, supports, cages, stalls, and around feeders. To help control flies, apply a directed application to horizontal surfaces and overhead areas and allow to dry before reintroduction of animals; also see Livestock/ Poultry Housing Structures and Pet Kennels.</p>
Cockroaches ¹ Crickets ⁶ Flies ⁵ Litter Beetles ³ (adults/immature stages such as darkling, hide, and carrion) Mosquitoes ⁴ Pillbugs Scorpions Spiders ⁵ Spider Mites (Two-spotted, Spruce) Stink Bugs ⁶ Ticks ⁶	0.8 fl. oz. (24 mL) per gallon of water (0.06% AI)	<p>Boxelder Bugs, Centipedes, Millipedes, Pillbugs, and Sow Bugs: Apply around doors and windows and other places where these pests may be found or where they may enter premises.</p> <p>Treat baseboards, storage areas, and other locations listed on this label. Apply treatments to prevent infestation as described below; also see Outdoor Surfaces Use.</p> <p>Fleas and Ticks: To control nuisance fleas and ticks (e.g. dog ticks) apply to kennels, yards, runs, and other areas where pets may frequent. To control ticks, apply using a coarse fan spray to vegetation brush, branches, rock walls, and other areas near habitation where ticks may harbor or frequent.</p> <p>Treat entire area rather than making spot treatments, and retreat as necessary to maintain control, as permissible on this label.</p> <p>Do not apply to pasture or cropland, and do not allow animals and people access to treated areas until the deposit has dried. Applications can begin in the spring and can continue until frost to control both larvae and adult ticks; also see Outdoor Surfaces Use. Do not apply directly to animals.</p> <p>Cluster Flies: Apply in late summer or early fall before flies are observed alighting on surfaces. Apply thoroughly on siding, under eaves, and around windows and doors, paying particular attention to south-facing surfaces, but do not allow runoff to occur. Heavy precipitation prior to frost may require retreatments to maintain protection. In winter and spring when flies become active and are emerging, interior crack and crevice and void treatments can help reduce the infestation, along with ULV or general surface application in infested attics or unoccupied lofts.</p> <p>Litter Beetles (Darkling, Hide, and Carrion Beetles) and Flies In Animal Housing (Such As Poultry Houses): To control adult litter beetles, apply Devito with EnduraCap Technology to walls and floors at cleanout, before reintroduction of animals. This will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to areas where beetles frequently occur, walls, supports, cages, stalls, and around feeders. To help control flies, apply a directed application to horizontal surfaces and overhead areas and allow to dry before reintroduction of animals; also see Livestock/ Poultry Housing Structures and Pet Kennels.</p>

PESTS	DILUTION RATE	SPECIFIC USE DIRECTIONS
Termites (above ground only)	0.2 fl. oz. (6 mL) to 0.4 fl. oz. (12 mL) per gallon of water (0.015 - 0.03% AI)	Devito with EnduraCap Technology may be used on termites to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. DO NOT use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.
¹ For cockroaches, the rate for maintenance treatments is 0.015% and for clean-out treatments is 0.03%. For control of severe infestations, use 0.06% rate. ² For outdoor use only and use 0.03% rate. ³ For control of LIGHT beetle infestations, use 0.03% rate. ⁴ For residual control, use 0.06% rate. ⁵ In all states except California, rates for flies may be increased to 0.06% when environmental conditions are severe and/or populations are high. ⁶ For clean-out/severe infestations, use 0.06% rate.		

Application within Food Handling Establishments

For places other than private residences in which exposed food is held, processed, prepared, or served: areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing foods, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups), and serving areas.

Use as a crack and crevice or spot treatment in and around both food and nonfood areas. Apply in small amounts directly into cracks and crevices, using equipment capable of delivering a pin stream of insecticide, in points between different elements of construction, between equipment and floor, openings leading to voids and hollow spaces in walls, equipment and bases. Clean food contact surfaces and equipment with an effective cleaning compound and rinse with potable water before using.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.). Take extreme care that the product is not introduced into the air. **Do not** contaminate food and food processing surfaces.

Application within Food Serving Areas

For facilities where foods are served, such as dining rooms. Apply as a crack and crevice or spot treatment to selective surfaces: baseboards, under elements of construction, and into cracks and crevices. **Do not** treat surfaces likely to be contacted by food. (**Do not** apply when facility is in operation or foods are exposed.) Food must be covered or removed in area being treated. **Do not** apply directly to food or allow applications to contaminate food.

NOTE: Application of this product in the Food Areas and/or Food Serving Areas of Food Handling Establishments other than as a spot and/or crack and crevice treatment is not permitted. Limit the use of application equipment (such as the Micro-Injector® or Actisol® applicator) in food areas to crack and crevice treatment only.

Livestock/Poultry Housing Structures and Pet Kennels

Apply as a general surface (including directed sprays) and/or crack and crevice treatment. Make interior and exterior perimeter applications in and around the livestock, poultry, and pet housing structures. Normal cleaning of the structure must also be done along with applications of **Devito with EnduraCap Technology** to effectively control the crawling and flying insect pests listed in the table.

With the exception of cattle and calves that may be present, only apply **Devito with EnduraCap Technology** to livestock barns or housing structures when unoccupied by animals. Apply to floors, vertical, and overhead surfaces where crawling or listed flying insect pests are or may be present. Cover feeders, waterers, and feed carts before application to prevent contamination. **Do not** apply to milk rooms or feed rooms. Pay attention to animal areas: stanchions, pipes, windows and doors, and areas where insect pests hide or congregate. Exterior applications to south facing walls and foundation perimeters can help prevent interior infestations of listed flying and crawling insect pests.

For poultry houses, apply to floor area (birds grown on litter) or to walls, posts, and cage framing (birds grown in cages). Application may also be made into cracks and crevices around insulation. Reapply after each growout or sanitization procedure. Indoor control can

be enhanced by making perimeter treatments around the outside of building foundations to prevent immigrating adult beetles. Apply in a uniform band 1 - 2 ft. up and 2 - 6 ft. out from foundation. Maintaining a year-round treatment program will prevent background populations from reaching problem levels.

RESTRICTIONS:

- **DO NOT** make interior applications of **Devito with EnduraCap Technology** in areas where animals, except cattle and calves, are present in the facility. Allow treated surfaces to completely dry before restocking the facility.
- **DO NOT** make applications to any animal feedstuffs, water, or watering equipment.
- **DO NOT** contaminate any animal food, feed, or water in and around livestock, poultry, or pet housing when making applications.
- **DO NOT** apply directly to animals.

Outdoor Surfaces Use

For control of ants, bees, centipedes, cockroaches, crickets, fleas, flies, millipedes, mosquitoes, scorpions, sowbugs, stink bugs, pillbugs, spiders, termites (above ground only), ticks, wasps, and other listed perimeter arthropod pests.

Apply with either hand or power application equipment as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches (crack and crevice only), screens, window frames, eaves, patios (crack and crevice only), garages, refuse dumps, and other similar areas where these insect pests are active.

For termites, **DO NOT** use this product as the sole source of control for active, structural infestations by subterranean termites; the purpose of this application is to kill workers or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control. It is not a substitute for mechanical alteration, soil and foundation treatment, but merely a supplement. For active, structural infestations by subterranean termites, this product can only be used to supplement a federally registered conventional product that is registered as a sole source for termite control. This product will not eliminate termite infestations or provide protection against future infestation.

Perimeter Barrier Treatments

Apply a continual band of insecticide around building foundations and around windows, doors, service line entrances, eaves, vents, and other areas to reduce the potential for entry by crawling pests. Before application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, deep mulch, or other potential pest harborage next to the foundation. Apply the band up to 7 ft. wide around the structure (or according to state regulations governing commercial pest control) and upwards along the foundation to 2 ft. and around windows, doors, and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active.

Amount of concentrate is dependent upon pest species (see pest table and comments), infestation levels, and service interval desired.

Rate Table for Perimeter Barrier Applications

Fl. oz. (mL) of Devito with EnduraCap Technology	Gals. of Water ¹	Area of Coverage (sq. ft.)
0.2 fl. oz. (6 mL)	1 - 5 gals.	800 - 1,600 sq. ft.
0.4 fl. oz. (12 mL)	1 - 5 gals.	800 - 1,600 sq. ft.
0.8 fl. oz. (24 mL)	1 - 5 gals.	800 - 1,600 sq. ft.

¹ Application volume may be greater than 5 gals./800 - 1,600 sq. ft. if required under heavy vegetative or landscaping materials in order to obtain desired coverage.

Examples of Dilutions for Perimeter Barrier Applications

Application Volume Gals. of Solution per 1,000 sq. ft.	Application Rate fl. oz. of Devito with EnduraCap Technology per 1,000 sq. ft.	Fl. oz. (mL) of Devito with EnduraCap Technology to Dilute in Water According to Spray Tank Volumes		
		5 gallons	10 gallons	50 gallons
1 gal. per 1,000 sq. ft.	0.2 fl. oz. (6 mL)	1 fl. oz. (30 mL)	2 fl. oz. (60 mL)	10 fl. oz. (300 mL)
	0.4 fl. oz. (12 mL)	2 fl. oz. (60 mL)	4 fl. oz. (120 mL)	20 fl. oz. (600 mL)
	0.8 fl. oz. (24 mL)	4 fl. oz. (120 mL)	8 fl. oz. (240 mL)	40 fl. oz. (1200 mL)
2 gal. per 1,000 sq. ft.	0.2 fl. oz. (6 mL)	0.5 fl. oz. (15 mL)	1 fl. oz. (30 mL)	5 fl. oz. (150 mL)
	0.4 fl. oz. (12 mL)	1 fl. oz. (30 mL)	2 fl. oz. (60 mL)	10 fl. oz. (300 mL)
	0.8 fl. oz. (24 mL)	2 fl. oz. (60 mL)	4 fl. oz. (120 mL)	20 fl. oz. (600 mL)
5 gal. per 1,000 sq. ft.	0.2 fl. oz. (6 mL)	0.2 fl. oz. (6 mL)	0.4 fl. oz. (12 mL)	2 fl. oz. (60 mL)
	0.4 fl. oz. (12 mL)	0.4 fl. oz. (12 mL)	0.8 fl. oz. (24 mL)	4 fl. oz. (120 mL)
	0.8 fl. oz. (24 mL)	0.8 fl. oz. (24 mL)	1.6 fl. oz. (48 mL)	8 fl. oz. (240 mL)

Example Calculation: To apply the mid-rate of **Devito with EnduraCap Technology** at a volume of 5 gal./1,000 sq. ft., mix 4 oz of concentrate in 50 gallons of water. The percent active ingredient in the finished **Devito with EnduraCap Technology** dilution can be calculated with the following formula:

$$\text{mL needed to add} = \frac{\text{times } 9.7\% \text{ active in concentrate}}{\text{finished dilution times } 3,785 \text{ mL/gal.} = \% \text{ active in dilution. (Example: 4 oz. in 50 gal. is 120 mL times 9.7 equals 1,164, and 50 gal. times 3,785 is 189,250. Dividing 1,164 by 189,250 equals 0.006\% active in the tank dilution.)}}$$

NOTE: Do not use water based sprays of **Devito with EnduraCap Technology** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard. Thoroughly wash out sprayer and screen with water and detergent before using **Devito with EnduraCap Technology**. **Devito with EnduraCap Technology** has not stained or caused damage to painted or varnished surfaces, plastics, fabrics, or other surfaces where water applied alone causes no damage. However, treat a small area and allow to dry, to determine whether staining will occur.

LET TREATED SURFACES DRY BEFORE ALLOWING HUMANS AND ANIMALS TO CONTACT SURFACES.

RESTRICTIONS:

- **DO NOT** use this product with oil.
- **DO NOT** apply this product in any room being used as living, eating, sleeping, or recovery area by patients, the elderly, or infirm when they are in the room.

- **DO NOT** apply to institutions (including libraries, sports facilities, etc.,) in the immediate area when occupants are present.
- **DO NOT** apply to classrooms when in use.
- **DO NOT** apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- **DO NOT** allow applications to contact water inhabited by fish, such as in aquariums and ornamental fish ponds that are located in/around structures being treated.
- **DO NOT** apply to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields or any other area frequented by the general public).

LAWNS/TURFGRASS

Devito with EnduraCap Technology may be used for applications to maintain non-residential indoor or outdoor areas where turf is grown. Treat non-residential landscapes around institutional, public, commercial and industrial buildings. Application rates for turf are lower than structural pest control rates, reflecting that treatment intervals are more frequent.

Applicators must ensure that they are certified in the necessary pesticide certification categories to allow application of **Devito with EnduraCap Technology** away from structures, such as to turf. Structural pest control certification categories may limit the distance away from structures for pesticide application. Consult your state extension office or pesticide regulatory officials for further information.

Mixing Instructions

Devito with EnduraCap Technology is to be mixed with water and may be used in all types of standard application equipment. Fill applicator tank with the desired volume of water and add **Devito with EnduraCap Technology**. It is suggested that the water be 5 - 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add **Devito with EnduraCap Technology** to applicator tank water with maximum agitation. Close and shake or re-agitate applicator tank before use if application is interrupted. Make up only amount of treatment volume as required.

Tank Mixing

Devito with EnduraCap Technology may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, perform a small volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Devito with EnduraCap Technology** last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Follow all restrictions and precautions which appear on the labels of these products.

RESTRICTIONS:

- **KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL SPRAY HAS DRIED FOLLOWING APPLICATION.**
- **DO NOT** apply when turfgrass is waterlogged or when soils are saturated with water (i.e. will not accept irrigation).
- **DO NOT** apply more than 0.36 lbs. of a.i. (52.4 fl. oz. of concentrate) per acre per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. For soil or foliar applications, **DO NOT** apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- **DO NOT** make outdoor broadcast applications to turf when wind speed is 15 mph or greater.
- In the state of New York, **DO NOT** apply within 100 ft. of coastal marshes or streams that drain into coastal marshes.
- **DO NOT** apply to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields, or any other area frequented by the general public).

RATES FOR LAWN AND TURFGRASS PESTS

PEST	AMOUNT OF DEVITO WITH ENDURACAP TECHNOLOGY*	APPLICATION INSTRUCTIONS
Ants (including Imported Fire Ants) Armyworms Centipedes Chinch Bugs ¹ Crickets Cutworms Earwigs Fleas (adult) Grasshoppers Japanese Beetles (adult) Millipedes Mites Mosquitoes (adult) Pillbugs Sod Webworms Sow Bugs Stink Bugs Ticks (including species which transmit Lyme disease)	3.4 - 7 mL per 1,000 sq. ft. or 5 - 10 fl. oz. per acre	<p>Application to turf should be started prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher labeled rates as pest pressure increases.</p> <p>Apply at 7-day intervals if retreatment is necessary. More frequent treatments should be limited to spot treatments.</p> <p>For spot treatments, use 0.5 fl. oz. of Devito with EnduraCap Technology per 1 - 2.5 gals. of water.</p> <p>Surface Insect Control (armyworm, cutworms, fleas, etc): Apply Devito with EnduraCap Technology at labeled rates in 2 - 5 gals. of water per 1,000 sq. ft. The use of a spreader-sticker may be useful if high rainfall amounts are forecast, otherwise the addition of adjuvants is not necessary under normal conditions for surface insect control in turf. Delay watering or mowing for 12 - 24 hours to control surface-feeding insect pests.</p> <p>Thatch Inhabiting Insect Control (chinch bugs, billbugs, etc): Apply Devito with EnduraCap Technology at labeled rates in 2 - 10 gals. of water per 1,000 sq. ft. The use of a nonionic wetting agent, penetrant or similar adjuvant is permitted at label rates. Lightly irrigate after application with up to 1/2 inch of water to move the Devito with EnduraCap Technology into the thatch layer. If irrigation is not available, then use high water application rates.</p> <p>Subsurface Insect Control (mole crickets, grubs, etc): Apply Devito with EnduraCap Technology at labeled rates in 4 - 10 gals. of water per 1,000 sq. ft. the use of a nonionic wetting agent, penetrant or similar adjuvant is permitted following label rates. Use the highest water application rates possible with your sprayer. Apply Devito with EnduraCap Technology to turf wet with dew, rain or irrigation. Water-in immediately after application with 1/4 - 1/2 inch of water.</p> <p>Fire Ant Control: Treat individual mounds with a drench application using a watering can. Use 0.5 fl. oz. of Devito with EnduraCap Technology per 2.5 gals. of water. Thoroughly soak each mound and a 3 ft. diameter circle around each mound. Gently apply the mixture to avoid disturbing the mound. Disturbing the mound may cause the ants to migrate and reduce the effectiveness of the treatment. Apply in early morning or late evening hours. You may make applications on a monthly basis to maintain protection of treated areas.</p> <p>Mosquito Control: Apply as a general spray around landscape plantings, turf, and building foundations to control mosquitoes. Apply Devito with EnduraCap Technology at labeled rates in 2 - 5 gals. of water per 1,000 sq. ft.</p>
Bluegrass Billbugs (adult) Black Turfgrass Ataenius (adult) Chiggers Fleas (adult) Grubs (suppression) Hyperodes Weevils (adult) Mole Crickets (nymphs and young adults)	7 mL per 1,000 sq. ft. or 10 fl. oz. per acre	
¹ Not for use on chinch bugs in New York State.		
* Example: To treat the listed turf pests at the 7 mL per 1,000 sq. ft. rate, determine the gallons of dilution per 1,000 sq. ft. needed to adequately cover the turf. At a 5 gallon per 1,000 sq. ft. dilution, take the 7 mL and divide it by the 5 (gallon) to get 1.4 mL to be added per gallon. For a 50 gallon tank, therefore, this would be 1.4 mL x 50 = 70 mL (or 2.5 oz.) in 50 gallons of water.		

Devito with EnduraCap Technology Mixing Chart for Turf Insect Pest Control

Rate of Devito with EnduraCap Technology	Application Rate Per 1,000 Square Feet of Turf				
	2 Gallons	4 Gallons	6 Gallons	8 Gallons	10 Gallons
5 oz. per acre	5.7 oz.	2.9 oz.	1.9 oz.	1.4 oz.	1.2 oz.
10 oz. per acre	11.5 oz.	5.7 oz.	3.8 oz.	2.9 oz.	2.3 oz.
Conversion Rate: 1 fluid ounce (fl. oz.) equals 29 milliliters (mL).					

ORNAMENTALS

Devito with EnduraCap Technology may be used for applications to maintain indoor or outdoor areas where ornamentals are grown. Treat non-turf areas of residential and non-residential landscaped areas around institutional, public, commercial and industrial buildings, parks, recreational areas, and athletic fields (trees, shrubs, flowers, evergreens, foliage plants and groundcovers). Application rates for ornamentals are lower than structural pest control rates, reflecting that treatment intervals are more frequent.

Applicators must ensure that they are certified in the necessary pesticide certification categories to allow application of **Devito with EnduraCap Technology** away from structures, such as to ornamental plantings. Structural pest control certification categories may limit the distance away from structures for pesticide application. Consult your state extension office or pesticide regulatory officials for further information.

IMPORTANT: Time application to flowering plants during periods when pollinating insects are not present, such as early morning or late evening.

NOTE: While phytotoxicity testing has been carried out on a wide range of ornamental plants under various environmental conditions, and no phytotoxicity has been observed, certain cultivars may be sensitive to the final spray solution.

It is advised to pre-spray a selection of ornamental plants and observe them for 7 - 10 days prior to treating large areas if local use experience is unavailable.

Mixing Instructions

Devito with EnduraCap Technology is to be mixed with water and may be used in all types of standard application equipment. Fill applicator tank with the desired volume of water and add **Devito with EnduraCap Technology**. It is suggested that the water be 5 - 7 pH. Adjust water pH with a buffering agent if necessary. Slowly add **Devito with EnduraCap Technology** to applicator tank water with maximum agitation. Close and shake or re-agitate applicator tank before use if application is interrupted. Make up only amount of treatment volume as required.

Tank Mixing

Devito with EnduraCap Technology may be tank mixed with other currently registered pesticides unless expressly prohibited by the product label. To ensure compatibility, perform a small volume mixing test with the other products. If other chemicals are added to the applicator tank, add **Devito with EnduraCap Technology** last. Fill tank to desired volume and continue to agitate while making applications. If mixed with EC formulations, use within 24 hours.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Follow all restrictions and precautions which appear on the labels of these products.

RESTRICTIONS:

- **DO NOT** apply more than 0.36 lbs. of the a.i. (52.4 fl. ozs. of concentrate) per acre per year.
- **DO NOT** apply this product through any type of irrigation system.
- **DO NOT** apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. For soil or foliar applications, **DO NOT** apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- **DO NOT** make outdoor broadcast applications to ornamentals when wind speed is 15 mph or greater.
- **DO NOT** apply to residential lawns and turf in residential settings (e.g., homes, parks, schools, athletic fields, or any other area frequented by the general public).

TANK DILUTION RATES FOR ORNAMENTAL PESTS

PEST	AMOUNT OF DEVITO WITH ENDURACAP TECHNOLOGY	APPLICATION INSTRUCTIONS
Ants (including Imported Fire Ants) Armyworms Azalea Caterpillars Aphids Bagworms Black Vine Weevils (adult) Boxelder Bugs Budworms California Oakworms Cankerworms Cockroaches Crickets Cutworms Eastern Tent Caterpillars Elm Leaf Beetles European Sawflies Fall Webworms Flea Beetles Forest Tent Caterpillars Gypsy Moth Larvae (continued)	1.5 - 5 fl. oz. per 100 gals. or 44 - 148 mL per 100 gals.	Application to ornamentals should be started prior to the establishment of high insect pest populations. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at labeled rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. For spot treatments use 0.5 fl. oz. Devito with EnduraCap Technology per 1 - 2.5 gals. of water. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. Scale: Thoroughly cover the plant with Devito with EnduraCap Technology spray, including trunks, stems, twigs, and foliage for control of scale insects (crawler stage). Bagworm: Apply Devito with EnduraCap Technology when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.

PEST	AMOUNT OF DEVITO WITH ENDURACAP TECHNOLOGY	APPLICATION INSTRUCTIONS
Japanese Beetles (adults) June Beetles (adults) Lace Bugs Leaf-feeding Caterpillars Leafhoppers Leafminers (adults) Leaf Rollers Leaf Skeletonizers Midges Mosquitoes Oleander Moth Larvae Pillbugs Pine Sawflies Pine Shoot Beetles Pinetip Moths Plant Bugs Root Weevils Sawflies Scale Insects (crawlers) Spiders Spittlebugs Stink Bugs Striped Beetles Striped Oakworms Thrips Tip Moths Tussock Moth Larvae Wasps	1.5 - 5 fl. oz. per 100 gals. or 44 - 148 mL per 100 gals.	<p>Application to ornamentals should be started prior to the establishment of high insect pest populations. Apply at 7-day intervals if retreatment is necessary. Limit more frequent treatment to spot treatments. Good spray coverage is necessary to provide the most effective level of control. Addition of a spreader-sticker at labeled rates may enhance the control of insects on certain species of ornamentals having waxy, hard to wet foliage. For spot treatments use 0.5 fl. oz. Devito with EnduraCap Technology per 1 - 2.5 gals. of water. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area.</p> <p>Scale: Thoroughly cover the plant with Devito with EnduraCap Technology spray, including trunks, stems, twigs, and foliage for control of scale insects (crawler stage).</p> <p>Bagworm: Apply Devito with EnduraCap Technology when bagworm larvae begin to hatch. Spray directly on the larvae. Application is the most effective when the larvae are young.</p>
Broadmites Brown Softscales California Redscale (crawlers) Clover Mites Mealybugs Pineneedlescales (crawlers) Spider Mites Whiteflies	3 - 5 fl. oz. per 100 gals. or 88 - 148 mL per 100 gals.	

Devito with EnduraCap Technology Mixing Chart for Ornamental Insect Pest Control
(Devito with EnduraCap Technology to add per spray tank)

Rate of Devito with EnduraCap Technology	25 Gallons	50 Gallons	100 Gallons	200 Gallons	300 Gallons
1.5 oz. ¹	0.4 oz.	0.8 oz.	1.5 oz.	3.0 oz.	4.5 oz.
3 oz. ²	0.8 oz.	1.5 oz.	3.0 oz.	6.0 oz.	9.0 oz.
5 oz. ³	1.3 oz.	2.5 oz.	5.0 oz.	10.0 oz.	15.0 oz.

¹ Equivalent to 3.5 mL per 1,000 sq. ft. (5 oz. per acre) when applied at 8 gal. per 1,000 sq. ft.

² Equivalent to 7 mL per 1,000 sq. ft. (10 oz. per acre) when applied at 8 gal. per 1,000 sq. ft.

³ Equivalent to 9.5 mL per 1,000 sq. ft. (14 oz. per acre) when applied at 8 gal. per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used, must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Devito™ with EnduraCap Technology is a trademark of Atticus, LLC.

20230816a

InTice¹⁰™

perimeter bait

**Kills Roaches, Crickets, Mole Crickets, Sowbugs, Pillbugs
Earwigs, Silverfish, Millipedes and Ants***

*Except Pharaoh ants, Carpenter Ants, Fire ants, Harvester ants

**For Indoor, Outdoor and Turf Uses
Mold and Moisture Resistant
Ready-to-Use Bait**

Active Ingredient: Orthoboric Acid 10%
Other Ingredients: 90%
Total:..... 100%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: **CAUTION:** Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product or label with you when calling a Poison Control Center or doctor or going for treatment. In case of emergency, for additional information call **800 858 7378**.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.



EPA Reg. No.: 73079-6
EPA Est. No.: 73079-MO-I
Net Contents: 10 lbs

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions

In food areas of restaurants, residential, health care, educational, daycare, agricultural buildings, animal rearing and handling establishments, food handling and food processing establishments, this product may only be applied as a crack and crevice treatment. Any product remaining outside of cracks and crevices must be cleaned up and removed.

Application Directions

Use Sites: In and around residential, multi-family, commercial, industrial, municipal, institutional, research, recreational, health care, educational, daycare, hospitality and agricultural buildings and other man-made structures, garages, home greenhouses, transport vehicles, sewers, animal rearing and handling establishments and food handling and food processing establishments. Turf sites include lawns, landscape beds, ornamental turf, parks, playing fields, right-of-ways, golf courses and home greenhouses.

Outdoor Application: Apply bait evenly over the affected area using a hand shaker or mechanical spreader at a rate of 1 lb per 1000 square feet or 45 lbs per acre. The product should not be watered in. For mole cricket treatments, results will be best if the area is irrigated prior to bait treatment. Mole crickets typically come to the surface to feed in the evening. Individual ant mounds may be treated by sprinkling 1/2 oz of product around each mound. Repeat treatment as needed. Apply bait early in the season in areas likely to be infested, such as flower, fruit and vegetable gardens, on ornamental turf and ground cover vegetation, and around shrubs. In home greenhouses, apply bait at a rate of 3 oz per 100 sq ft on and under benches. Do not apply bait directly in flats or pots. Reapply as needed.

Sewer Application: Apply bait along ledges inside sewers and around manhole covers.

Indoor Application: Apply bait only in areas inaccessible to children and pets, or in tamper-resistant bait stations. Apply bait at a rate of 3 oz (2/3 cup) per 100 square feet. Using a hand shaker or duster, apply bait in attics, crawl spaces, trash closets, utility closets, dropped ceilings, on dirt or gravel cellar floors, under and behind appliances and vending machines, in wall voids and voids under and behind cabinets, sinks and tubs, in drawer wells, garbage chutes, pipe chases, elevator pits, around drains, around electrical conduits and in cracks and crevices where insects may harbor. Focus application in areas where insects or their signs are seen. Reapply bait after one week if infestation is still present. A power duster may be used to apply bait in appropriate areas, such as attics and crawlspaces. The duster manufacturer's recommendations for settings should be followed.

Application tips for certain sites follow:

1. In attics, be sure to get bait near the eaves and vent pipes where insects often first enter, as well as around any pipes or potential access points between the attic and the main structure. Attics should have proper ventilation to insure that no food odor from the bait seeps into the living area, particularly in hot weather. If in doubt, apply the product in a manner such that it can be easily removed.
2. When treating cabinetry areas, if there is no gap between the lower cabinets and the kickplate, then small holes can be drilled to access the voids. Escutcheon plates around pipes can also be pulled back to allow access to voids.
3. The bottom drawers in kitchen and bathroom cabinets can be removed and bait dusted into drawer wells. No bait should be left exposed when drawers are replaced.

4. Electrical switch plate covers can be removed to allow access to voids, but bait shouldn't be applied directly in electrical boxes.
5. If there are cracks between baseboards, cabinets, door frames, hinges, counters or trim, and walls, these cracks can be lightly dusted with bait. If practical, these areas can be sealed once insect problems have been eliminated, to help prevent future problems.
6. Insects need moisture, so kitchen and bath areas and areas immediately surrounding them, and any other damp areas should be carefully inspected and treated if necessary. As a separate matter from the use of this product, it is advisable to fix any leaks to minimize structural damage and odors, and to prevent potential chronic insect problems.

To help prevent insect problems from developing in homes and similar structures, a preventative program is recommended. Once per year, apply bait in attics and crawlspaces at a rate of 8 oz per 1000 square feet.

Food Processing and Handling Establishment Application, including Federally-Inspected Meat and Poultry Plants: Follow instructions for other sites as appropriate. **In food areas, only apply bait in cracks, crevices, and other inaccessible areas.** Avoid contamination of food and feedstuffs. Never apply bait directly to a surface where food is stored, prepared or served. Any bait that is left visible after treatment should be swept into cracks and crevices or removed.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place. Do not store where children or animals may gain access.

Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or **1-800-CLEANUP** which is managed as a public-private partnership.

Conditions of Warranty and Limitations of Liability

IMPORTANT: READ BEFORE USE Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Manufacturer. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Manufacturer makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Manufacturer is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Manufacturer disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Manufacturer's election, the replacement of product.

Manufactured by:

 **Rockwell Labs Ltd**
creating the future of pest control

North Kansas City, MO 64116

Toll Free: 866 788 4101 • www.rockwelllabs.com

11-2010

D-FENSE NXT

PRESSURIZED SOLUTIONS

- Contact kill and residual control
- For use on mattresses, furniture and carpets
- Keeps working on listed cockroaches for up to 180 days
- Kills bed bugs where they hide
- Breaks the life cycle of listed cockroaches
- For indoor/outdoor use
- Broad spectrum insecticide with 2 IGRs

Kills Ants, Bees, Bed Bugs, Carpenter Ants, Carpenter Bees, Cockroaches, Fleas, Flies, Fruit Flies, Gnats, Grain Mites, Hornets, Indian Meal Moths, Mosquitoes, Spiders, Termites† (Subterranean, Formosan & Drywood), Ticks, Wasps

ACTIVE INGREDIENTS:

Novaluron	0.20%
Deltamethrin	0.06%
Pyriproxyfen	0.02%

OTHER INGREDIENTS*: 99.72%

TOTAL: 100.00%

*Contains petroleum distillates



Manufactured for:
Control Solutions, Inc.
5903 Genoa Red Bluff
Pasadena, TX 77507

EPA Reg. No: 53883-415
EPA Est. No: 53883-TX-002
Net Contents: 15 OUNCES

FOR USE IN AND AROUND: Apartments, Campgrounds, Non-Food Storage Areas, Homes, Hospitals†, Hotels, Motels, Nursing Homes†, Resorts, Restaurants and other Food Handling Establishments†, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes†), Utilities, Warehouses, and other Commercial and Industrial Buildings. † See special instructions for these sites under Directions for Use.

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
IF ON SKIN:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International for emergency medical treatment at (866) 897-8050.

NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. Do not apply directly to any body of water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flame. Do not puncture or incinerate container. Do not use in the vicinity of open flame or pilot lights. Do not apply to plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc. Do not apply where electrical shock hazards exist.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- Remove pets, birds, and cover fish aquariums before spraying.
- Test first for staining on an inconspicuous area prior to use.
- Hold container upright and DO NOT spray up into air.
- Remove protective cap and direct spray away from face.
- Hold can 12-15 inches from surface being sprayed.
- Press button down and spray as directed.

† **IN HOSPITALS AND NURSING HOMES:** Do not apply this product in hospital rooms while occupied or in any rooms occupied by the elderly or infirm. Apply according to specific directions for pest(s) to be treated. Patients should be removed from room prior to treatment. Room should be ventilated for two hours after spraying. Do not return patients to room until after ventilation.

† **FOOD/FEED HANDLING ESTABLISHMENTS (Places other than private residences in which food is held, processed, prepared or served. Including areas for receiving, storing, packing (canning bottling, wrapping, boxing) preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas where food/feed is exposed and the facility is in operation.):** Cover or remove exposed food/feed from the treatment area prior to application. All food processing surfaces should be covered during treatment or thoroughly washed before use. Do not apply directly to food or food handling surfaces. This product is for use in Federally Inspected Meat and Poultry Plants as a crack and crevice treatment. Except in federally inspected meat and poultry plants, spot or crack and crevice applications may be made

while the facility is in operation. 1 second spray per spot. General surface applications must be used only when the facility is not in operation provided exposed food/feed and all food/feed processing and/or handling equipment has been covered or removed prior to application. After application in food/feed processing plants, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food/feed will contact.

Non-Food/Non-Feed Areas: Examples of non-food/non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). All areas that insects inhabit or through which insects may enter should be treated.

† **PLANES:** Do not apply in aircraft cabins. Do not spray on plastic, painted, or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers, etc. Do not use this product in electrical equipment (where there is exposed wiring or contact points) due to possibility of short circuits and shock hazard. Product should only be used when can temperature is above 60°F. If can temperature is below 60°F, store at room temperature until a temperature above 60°F is reached.

CRACK AND CREVICE APPLICATION: Use D-Fense NXT with the supplied actuator and injection tubes. Inject D-Fense NXT into cracks and crevices or void spaces where insects may be harboring, living or breeding. Place injector tip into cracks, crevices, holes and other small openings. Apply product for one second. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For wall and equipment voids calculate the void's cubic area and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids. Reapply once every 6 months if necessary.

SPOT TREATMENT: Apply to storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, shelves, drawers and similar areas. Contact as many insects as possible. Hold can at arm's length and apply at the rate of 1 second per 2 sq. ft. Reapply once every 6 months if necessary.

GENERAL SURFACE APPLICATION: Hold can at arm's length and apply a light spray at the rate of 10 seconds per 100 sq. ft. Reapply once every 6 months if necessary.

INDOOR TREATMENTS

D-Fense NXT is acceptable as a pesticide for use in all areas of official establishments operating under the Federal meat, poultry, shell egg grading, and egg products inspection programs.

CRAWLING INSECTS (SUCH AS: ANTS, BOOKLICE, CARPET BEETLES, CENTIPEDES, CLOVERMITES, CRICKETS, MILLIPEDES, ROACHES, SILVERFISH, SOWBUGS, AND SPIDERS): Apply as a crack and crevice treatment into cracks and crevices in all hiding places, behind baseboards, sinks, cabinets, meter boxes, door frames, windows and closets. Apply as a spot treatment or general surface application to breeding areas, nesting sites, hiding places and other places where crawling insects may be found.

BED BUGS: Remove linens and wash before reuse. Take bed apart and treat cracks, joints and interior of framework. Treat mattresses and box springs; especially tufts, folds, and edges. Also treat picture frames, baseboards, moldings and all cracks and crevices in the room. Allow to dry

before remaking bed. Do not use in patient rooms in hospitals and nursing homes for treatment of bed bugs. May be used in conjunction with a bed bug proof mattress encasement. To treat furniture, such as couches and upholstered chairs, turn the piece upside down, lift the dust cover from the bottom and spray the interior frame. Replace the dust cover after frame treatment and turn piece right side up.

FOR EXPOSED ADULT AND LARVAL STAGES OF CARPET BEETLES, CIGARETTE BEETLES, CONFUSED FLOUR BEETLES, DRUGSTORE BEETLES, GRAIN MITES, GRANARY WEEVILS, RED FLOUR BEETLES, RICE WEEVILS, AND SAWTOOTHED GRAIN BEETLES: Apply as a Crack and Crevice treatment into cracks and crevices of pantries, cabinets, food processing and handling equipment and other places where insects harbor. Apply around cartons, and containers, and other areas as where these insects tend to congregate. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

FLEAS AND TICKS ON SURFACES: Remove pet bedding and destroy or clean thoroughly. Spray pet resting quarters. Adult fleas and larvae contacted by spray will be killed. Put fresh bedding down once spray has dried. DO NOT spray animals directly. Hold product 36" above floor and direct toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. making sure that all floor (rugs and carpets), sofa and chair surfaces are contacted. Treat adjacent cracks and crevices such as behind baseboards and edges of carpet and floor covers. Treat higher cracks such as upper door jams and window framing, behind pictures and other areas where females may crawl to lay eggs. Treatment may be repeated daily if necessary. Keep area closed for 15 minutes. Open and ventilate before reoccupying. The insect growth regulators in CSI 16-105 Delta + Tekko Pro work for up to 7 months by inhibiting the development of the immature stages of the flea and preventing them from reaching the biting adult stage. For best results, pets should be treated with an appropriate product before allowing them to return to the treated area.

OUTDOOR TREATMENTS

CRAWLING INSECTS ON OUTSIDE SURFACES OF BUILDINGS: Spray foundation of building where insects are active and may find entrance up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include areas such as screens, window frames, eaves, porches, patios, garages and refuse dumps. Spray infested surfaces of patio or picnic areas, hitting as many insects as possible. Also, spray legs of tables and chairs to repel ants.

ANTS: Apply to ant trails, around doors and windows and other places where ants may be found.

FLYING AND STINGING INSECTS: Use only on outdoor surfaces. Spray outside surfaces of windows and door frames and other areas where these pests may enter the home. Also spray localized resting areas, such as under eaves, porches, light fixtures and cords, railings, etc. where these insects may rest. Insects coming to rest on treated surfaces will be killed.

WOOD INFESTING INSECTS

CARPENTER ANTS, CARPENTERS BEES, AND TERMITES[†] (SUBTERRANEAN, FORMOSAN AND DRYWOOD): The following applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplement for controlling active infestations of wood destroying insects including Subterranean, Formosan, and Drywood termites. This

product is not recommended as sole protection against termites. For active indoor termite infestations, get a professional inspection. For control inside and outside of structures, apply as general surface, spot, crack and crevice or wall void application. To kill wood destroying insect infestations in localized above-ground areas of the structure, inject into galleries or voids in damaged wood, or into spaces between damaged wooden structural members. Locate galleries by visual signs of wood destroying insect infestation through the use of mechanical sounding techniques. Where necessary, penetrate the galleries or voids by drilling holes to receive the injector tip. Distribute drill holes to adequately treat the infestation. Applications must be made to inaccessible structure voids by drilling and then injecting into the infested void.

For termites: The purposes of such applications is to kill workers or winged forms which may be present in the treatment areas at the time of application.

For carpenter bees: Following treatment, the entrance holes must be left open 24 hours to be certain that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant.

Wood Destroying Insects Living Inside Trees, Utility Poles and Fences: CSI 16-105 Delta + Tekko Pro Insecticide must be used as a spot treatment for control of above-ground termites inside trees, utility poles, fencing and decking materials and similar structural members. Wherever possible, locate an inferior infested cavity and inject in the appropriate quantities. Repeat applications as needed to maintain control of re-infestations.

Aerial Termite Carton Nests in Trees: Termite carton nests in trees must be injected with CSI 16-105 Delta + Tekko Pro Insecticide using an injection applicator. Multiple injection points to varying depths may be necessary.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store upright at room temperature. Do not store near heat or open flame. Store in an area away from children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Do not puncture or incinerate! Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling, if available. **If partly filled:** Call your local solid waste agency for disposal instructions.

LIMITATION OF WARRANTY AND LIABILITY

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Control Solutions, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Control Solutions, Inc. and Seller harmless for any claims relating to such factors.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use

conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Control Solutions, Inc., and Buyer and User assume the risk of any such use. **CONTROL SOLUTIONS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

IT IS CONTROL SOLUTIONS INC., AND SELLERS INTENTION THAT in no event shall Control Solutions or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CONTROL SOLUTIONS, INC., AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CONTROL SOLUTIONS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Control Solutions, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Control Solutions, Inc.