be thoroughly washed with an effective cleaning compound and rinsed with potable water after use of this product. Apply this product indoors to fire ants on non-porous surfaces only. **DO NOT** apply more than once every 7 days.

Indoors: Apply directly to listed insects as a crack-and-crevice treatment application in all hiding places, including behind baseboards, sinks, cabinets, meter boxes, door frames, wall voids. plumbing chases, under sink cabinets, attics, windows and closets. Apply directly to listed insects as a spot treatment application to breeding and nesting sites for listed crawling insects (excluding fire or carpenter ants), hiding places and other places where crawling insects may be found. Apply this product indoors to fire ants on non-porous surfaces only. **DO NOT** apply more than once every 7 days.

Outdoors: Apply to surfaces on buildings, porches, patios, and other structures, openings ground doors and windows, eaves and attic vents, utility entry points, weep holes, soffit areas and other exterior openings (including foundation cracks or drilled holes) where listed pests may enter the structure or where they have been seen or found, or can find shelter. Apply to ant trails, around doors and windows, in crawl spaces and other places where ants may be found. **DO NOT** apply more than once every 7 days. **DO NOT** apply this product outdoors to fire or carpenter ants.

BED BUGS: Remove linens and wash before reuse. Spray bed bugs and eggs directly. Take bed apart and spray bed bugs and eggs directly that are found in or on cracks, joints and interior of framework las well as bedsprings, box springs and furniture. On infested mattresses, apply directly to bed byas and eags. Allow treatment to dru thoroughly before remaking bed. **DO NOT** use in patient rooms lin hospitals and nursing homes for treatment of bed bugs while occupied by patients. As part of a bed bug treatment program, this product may be used in conjunction with a bed bug proof mattress encasement. DO NOT apply more than once every 7 days.

PANTRY INSECTS/STORED PRODUCT INSECT PESTS: Apply around cartons, containers, shelving, storage areas, closets, cabinets, behind or under walls and baseboards and other areas where these insects tend to congregate. **DO NOT** apply to areas where food/feed, food utensils or food processing surfaces may come into contact and become contaminated. Infested products should be disposed of. DO NOT apply more than once every 7 days.

Indoors: Apply as a spot treatment to surfaces where house flies or stable flies congregate, collect or rest. **DO NOT** applu more than once every 7 days.

Outdoors: Apply directly to listed insects on outside surfaces of windows and door frames and other areas where these pests may enter the home. Also spray localized resting areas, such as under leaves, porches, light fixtures and cords, railings, etc. where these insects may rest, DO NOT apply more than once every 7 days.

CARPENTER BEES: Apply the spray directly into gallery entrance holes. Exterior wood surfaces may be treated to prevent nest cavity excavation by adult carpenter bees. Following treatment, the lentrance holes must be left open 24 hours to be certain that returning adult bees are killed. When there is no activity, the hole may be sealed with a suitable sealant. DO NOT apply more than once

WOOD DESTROYING INSECTS OUTDOORS: This product may be used as a spot treatment for control of above-ground termites and wood infesting beetles inside trees, utility poles, fencing and decking materials and similar structural members. Wherever possible, locate an interior infested cavity and spray directly into the entrance hole to listed wood destroying insects which are present at the time of treatment. Repeat applications as needed to maintain control of re-infestations. This product is not intended to provide structural termite pest control. For termites, the purpose of this application is to kill workers and/or winged reproductive forms of termites which are present at the time of treatment. It is not intended to provide structural pest control, and is not a substitute for mechanical EPA Reg. No. 53883-455 alteration, soil and foundation treatment, but merely as a supplement. DO NOT apply more than once every 7 days.

Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store upright at room temperature. Do not store near heat or open flame. Keep this product in its tightly closed original container. Store in a cool, dry (preferably locked) area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER HANDLING: Do not puncture or incinerate! Nonrefillable container. Do not reuse of refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedu of the user or buuer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc. election, the replacement of product.







$\mathsf{DOXEM}^{ exttt{@}}$ NX PRESSURIZED SOLUTIONS

- Kills pests on contact and leaves behind a residual
- Contains 4 unique modes of action
- For use in and around food handling establishments

KILLS: Ants, Bed Bugs, Booklice, Boxelder Bugs, Carpenter Bees, Carpet Beetles (Black and Varied), Centinedes, Cigarette Beetles, Clover Mites, Cockroaches, Crickets, Darkling Beetles, Drain Flies, Earwigs, Firebrats, Fruit Flies, Ground Beetles, House Flies, Lady Beetles, Leaffooted Bugs, Mealworms (Yellow or Dark), Millipedes, Pantry Insects, Pillbugs, Pharaoh Ants, Silverfish, Sowbugs, Spiders, Spider Beetles, Springtails, Stable Flies, Stink Bugs, Stored Product Pests, Termites (subterranean and drywood), Webbing Clothes Moths, Weevils and Wood Destroying Insects.

FOR USE IN AND AROUND: Apartments, Non-Food/Non-Feed Storage Areas, Food/Feed Handling Establishments, Homes, Hospitals, Hotels, Kennels, Animal Housing Facilities (poultry houses, cattle, sheep, swine, goat and dairy facilities, horse stables, and pet sleeping quarters), Medical Facilities, Motels, Nursing Homes, Resorts, Non-Food/Non-Feed and Food/Feed Areas of Restaurants and other Food/Feed Handling Establishments, Schools, Stores, Supermarkets, Modes of Transport (Buses, Boats, Ships, Trains, Trucks, Planes (cargo areas only)), Utility Rooms, Warehouses, and other Commercial and Industrial Buildings. For indoor and outdoor (excluding fire or carpenter ants) use

ACTIVE INGREDIENTS:

| Indoxacarb | 0.10% |
|---------------------------------|-----------|
| Novaluron | 0.20% |
| Imidacloprid | 0.05% |
| Pyriproxyfen | |
| OTHER INGREDIENTS* | 99.63% |
| TOTAL: | 100.00% |
| *Contains petroleum distillates | |

EPA Est. No: 53883-TX-002

KEEP OUT OF REACH OF CHILDREN

FIRST AID

day seven (7) days per week.

- Immediately call a poison control center or doctor Do not induce vomiting unless told to do so by a poison control center or doctor
- Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
- Have the product container or label with you when calling a poison control center or doctor, or going for treat-ment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), uou mau contact SafetuCall® International at 1866) 897-8050, twentu-four (24) hours per

NOTE TO PHYSICIAN: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.

PRODUCT INFORMATION: This product contains four active ingredients: two adulticides and two insect growth regulators. The adulticides give quick kill of existing indoor adult insects while the insect growth regulators work to prevent new adults from developing. Apply this product as an indoor crack-and-crevice, spot or void treatment. Spray may leave a slight residue with heavy applications that is easily removed with soap and water. For outdoor applications (excluding fire ants and carpenter ants) this product will kill and control listed pests as a crack-and-crevice, spot or void application.

NET CONTENTS: 15 FL OUNCES

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Wash hands thoroughly with soap and water after handling and

before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. DO NOT apply directly to any body of walter To protect the environment, **DO NOT** allow pesticide to enter or run-off into storm drains, drainage ditches, autters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the

PHYSICAL AND CHEMICAL HAZARDS Extremely Flammable. Contents under pressure. Keep away from fire, sparks, and heated surfaces DO NOT puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For indoor or outdoor use as either a crack-and-crevice injection or spot treatment. Use this product

with the supplied actuator. USE RESTRICTIONS: DO NOT use this product to treat drywood termites in New York State. DO NOT use this product as

the sole source of control for active, structural infestations by subterranean termites. DO NOT apply this product directly to companion animals (pets). DO NOT apply to rooms or structures while the are occupied by or easily accessible to companion animals (pets). **DO NOT** make applications to any animal feedstuffs, water or watering equipment. DO NOT contaminate animal food, feed or water in and around companion animal housing while making applications. DO NOT apply this product in a way that will contact adults, children or pets, either directly or through drift. DO NOT allow adults children or pets to enter the treated area until sprays have dried. **DO NOT** apply in classroom while they are in use. Room should be thoroughly ventilated after spraying before re-entry. DO NOT apply to institutions (e.g., libraries and recreational facilities) while occupants are present in the immediate area. **DO NOT** apply this product in patient rooms while occupied, or in any rooms while occupied, including rooms occupied by the elderly or the infirmed. DO NOT use in serving areas while food is exposed. In the home, cover or thoroughly wash all food processing surfaces and utensils following treatment if contaminated. Cover exposed food and drinking water or remove from area being treated. All outdoor applications must be limited to spot or cracks-and-crevices treatments only. **DO NOT** apply in aircraft cabins. Applications to aircraft must be limited to card areas and galley kitchen only. **DO NOT** use this product in electrical equipment (where there exposed wiring or contact points) due to possibility of short circuits and shock hazard. DO NOT make outdoor applications during rain. DO NOT apply directly to outdoor impervious horizontal surface such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. DO NOT apply this product to areas or surfaces where food/feed, food utensils, or food-processing surface may come into contact and become contaminated. **DO NOT** apply to food prep areas that a routinely washed, such as cracks and crevices in tops of tables, food/feed preparation and holding surfaces for prepared foods. DO NOT apply directly into sewers or drains, or to any area like a gutt where drainage to storm sewers, water bodies, or aquatic habitat can occur, unless labeled for drain treatment. Remove pets, birds and cover fish aquariums and ornamental ponds before spraying and turn aquarium systems off. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

second of spray per injection. For a heavy infestations, make injections approximately 12" apart. F light infestations or preventative treatment, make applications approximately three feet apart. long running cracks, move injector tip along the crack while spraying at the rate of three linear less per second. For heavy infestations move injector tip along at the rate of one linear foot per secon For inaccessible closed voids calculate the void's volume and treat at rate of five to ten seconds be three cubic feet. Several holes may be required in long running voids. Apply this product indoors to fire ants on non-porous surfaces only. **DO NOT** apply more than once every 7 days. SPOT TREATMENT: Apply to storage facilities, closets, around water pipes, doors and windows

CRACK-AND-CREVICE APPLICATION: Inject this product into cracks-and-crevices or void space

where listed insects may be harboring, living or breeding. Place injector tip into cracks, crevide

holes and other small openings for direct spray to target listed insects. Release approximately on

behind and under refrigerators, cabinets, sinks, stoves and other equipment, shelves, drawers and similar areas. Spray as many listed insects as possible directly. Hold can 12 to 15 inches from surface to be treated. Apply at the rate of 2 seconds per 2 sq. ft. Spot treatments must NOT exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.). Apply this product indoors to fire gnts on non-porous surfaces only. **DO NOT** apply more than once every 7 days.

FOOD/FEED HANDLING ESTABLISHMENTS - Non-Food/Non-Feed Areas: Examples of non-figo non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packagin canning or bottling). All areas that insects inhabit or through which listed insects may enter should be treated directly. This product may be used while the facility is in operation. Use as a crack-and crevice or spot treatment as instructed above. Apply this product indoors to fire ants on non-porbus surfaces only. DO NOT apply more than once every 7 days.

FOOD/FEED HANDLING ESTABLISHMENTS – Food/Feed Areas: Examples of food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible wdste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. This product may be used while the facility is in operation. Use as a crack-and-crevice treatment only, **DO NOT** apply to areas where food/feed, food utensils or food processing surfaces may come into contact and become contaminated. If food contact surfaces become contaminated, contact surfaces must



Maxf@rce FC MAGNUM

ROACH KILLER BAIT GEL

Note: Do not remove this label. Use only in areas not easily accessible to children and pets.

†CAS NO. 120068-37-3

EPA Reg. No. 432-1460 EPA Est. No. 071106-GA-003

KEEP OUT OF REACH OF CHILDREN

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash throughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Net Wt. 33g (1.16 Oz) 79432135 80211538C 140410AV1 May stain textiles, fabrics and/or porous surfaces. Refer to leaflet for complete First Aid, Directions for Use, Precautionary Statements and Storage and Disposal Instructions.

Produced for:

> Bayer Environmental Science A Division of Bayer CropScience LP 2 T. W. Alexander Drive Research Triangle Park, NC 27709

CONTAINER HANDLING: If empty: Do not reuse

container. Place in trash or offer for recycling if

available. If partly filled: Call your local solid waste

agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or

outdoor drain. If you have questions regarding the use

of this product, call toll free 1-800-331-2867.

Baver Product of China



ROACH KILLER BAIT GEL

79432135

80215495C 140410AV1

FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

Long lasting control For indoor and outdoor use See visible results overnight No vapor, fumes or odor Kills roaches quickly

ACTIVE INGREDIENT: Fipronil*: [5-Amino-1-(2, 6-dichloro-4-trifluoromethyl) phenyl)-4-(1, R, S) -(trifluoromethyl) sulfinyl)-1H-pyrazole-3-carbonitrile] OTHER INGREDIENTS: ... 99.95% 100.00% *CAS NO. 120068-37-3

EPA Reg. No. 432-1460

EPA Est. No. 071106-GA-003

STOP - READ THE LABEL BEFORE USE. KEEP OUT OF REACH OF CHII DRFN CAUTION

Product may stain textiles and clothing.

FIRST AID

| If on skin |
|--------------|
| or clothing: |
| |

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for
- 15-20 minutes Call a poison control center or doctor for treatment
- advice.

If in eves:

- . Hold eve open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using

PHYSICAL AND CHEMICAL HAZARDS: Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

Long lasting control
No vapor, fumes or odor

For indoor and outdoor use

Kills roaches quickly See visible results overnight

† American, Oriental, Brown, Smoky Brown

Provides FAST CONTROL in 2 days (German roaches).

Dual Action kills roaches two ways.

Contains Fipronil, which kills roaches by ingestion and contact. After foraging roaches either eat or contact the bait, they return to the harborage and kill other roaches.

Kills roaches in their harborage.

Inject bait into cracks and crevices or apply as a spot where roaches harbor or enter a structure.

This product effectively controls large roaches†.
This product contains Fipronil, a fast-acting active ingredient that kills American roaches. This product controls large roacht populations in 8

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Use Maxforce® FC Magnum Roach Killer Bait Gel as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches in residential areas and the nonfood/non-feed areas of institutional, warehousing and commercial establishments, including warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, apartment buildings, bakeries, beverage plants, breweries, bottling facilities, candy plants, canneries, cereal processing and manufacturing plants, dairies and dairy product processing plants, frozen food plants, schools, laboratories, computer facilities, sewers, "aircraft," buses, boats/ships, trains, homes, houses, industrial buildings, office buildings, kennels, kitchens, manufacturing facilities, spice plants, stores, wineries and similar structures, mausoleums, meat processing and packaging plants, meat and vegetable canneries, pet shops, and zoos. Nonfood/non-feed areas include bedrooms, living rooms, bathrooms, closets, media rooms, garbage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets, and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as outlined

May stain textiles, fabrics, and or porous surfaces.

USE RESTRICTIONS

Do not treat food preparation surfaces. Keep gel away from foods and food contact surfaces. Do not apply bait gel to areas which have been recently sprayed with insecticide, and do not spray insecticide over gel as it may cause the bait to become repellent. USE ONLY IN AREAS NOT EASILY ACCESSIBLE TO CHILDREN AND PETS. In pet shops and zoos place bait out of reach of pets and zoo animals.

APPLICATION DIRECTIONS

Application rate for all uses: Application rate varies according to the level of infestation and species to be controlled. Apply gel as round spots or thin beads into cracks and crevices or as a spot treatment. Place bait close to cockroach harborages, foraging, and feeding areas such as corners and cracks and crevices. Numerous smaller bait placements provide faster control than fewer larger placements. In cases of severe infestation and areas with excessive clutter or alternative food sources that cannot be eliminated, use the higher rate of application.

German Cockroaches, Brown Banded Cockroaches, and stored product pests:

For use in maintenance programs where regular [monthly, quarterly] applications are made for continuous control of cockroach populations,

use 0.75 grams of bait per square yard. This rate will not provide fast control of existing infestations

For light to moderate infestations (or for retreatments), use up to 1.5

grams of bait per square yard.
To clean out severe or heavy infestations, use up to 3.0 grams of bait per square yard.

Large cockroaches such as American, Smokey-Brown or Oriental: 1 to 2 grams of bait per square yard.

Poit Quantity Reference Table

1.5 gm = Quarter size spot

| But Quality Reference Tubic | | |
|-----------------------------|--------------------------------------|---------------------------------------|
| | Approximate weight per spot diameter | Approximate weight per bead length |
| | 0.1 gm = 1/4 inch diameter spot | 1/3 gram = 1.5 inch bead |
| | 1.0 gm = Dime size spot | 2/3 gram = 3.0 inch bead |

1 1/3 gram = 6 inch bead

Place bait under baseboards, sinks, countertops, seats, around water pipes, water heaters, dishwasher equipment, in crack and crevices as well as voids where insects harbor.

PANTRY PESTS: To control exposed, adult stages of beetles: cadelle, cigarette beetle, confused flour beetle, dermestid, drugstore beetle, flat grain beetle, granary weevil, lesser grain borer, lesser mealworm, mealworm, merchant grain beetle, red flour beetle, rice weevil, rusty grain beetle, saw-toothed grain beetle, spider beetle, or warehouse beetle. After removing infested foodstuffs and cleaning up spilled foodstuffs and dust, applications of this product will kill adult stages of beetles remaining in area. Apply in voids, cracks and crevices, behind and under cubboards, and other difficult to reach areas where insects may hide. Application rate is dependent on level of infestation.

Applications in Food/Feed Handling Establishments: Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. If gel contacts an exposed surface where food is handled, remove gel and wash exposed surface with an effective cleaning compound followed by a potable water rinse prior to use. Apply with Maxforce Bait Injector directly into cracks and crevices. Place applicator tip 1/2 inch into cracks. crevices, holes and other small openings where roaches may be a problem. Apply small amounts of gel.
Use Restrictions in Food/Feed Handling Establishments:

Do not apply bait to areas where food/feed, utensils, or processing surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation, and prepared food holding surfaces as bait may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 130°F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, fume hoods, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers, and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves, and other equipment where cockroaches hide. During follow-up visits inspect bait placements and reapply when necessary

Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not

permitted.
Gel placements should be at or near harborages or aggregation areas, such as corners or cracks and crevices. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control.

For outdoor use (adjacent to home -or- structure): Place bait in areas adjacent to structures where [large] roaches may nest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion

Apply bait in protected areas whenever possible. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness. Priority treatment areas may be identified by visual inspection of such

areas listed above or aided by trapping techniques or spot flushing. Refillable stations: Maxforce FC Magnum Roach Killer Bait Gel may be placed into refillable bait stations designed for interior or exterior use; when placing bait in stations do not exceed 1 gm (1/5 teaspoon) and do not exceed the capacity of the stations. Close the station securely after application. During follow up visits inspect placements and reapply as needed.

Directions for loading reservoirs and operation:

With the teeth the drive rod facing up, pull the drive rod all the way back. If the drive rod is initially hard to move simply un-screw (turn counter-clockwise) the black portion of the applicator that holds the gel reservoir until the drive rod moves easily.

2. Attach a pre-filled reservoir of gel with a twist-locking motion.

Remove the outlet cap and attach a dispensing tip. Break-tip Reservoir: Prior to use, break the snip-tip closed end of the gel reservoir at the scored line. After use, the reservoir may be resealed with the enclosed black outlet cap. The black outlet cap is reusable. 3. Turn the knob of the drive rod so that the teeth are facing down and

move it slightly forward so that the trigger mechanism will engage. Squeeze the trigger until the drive plunger seats against the piston in the gel reservoir. Gently squeeze the trigger until the gel flows out.

4. If gel continues to flow after the trigger is released, simply un-screw the portion of the bait applicator that holds the reservoir to a point where gel does not continue to flow after the trigger is released.

Directions for changing reservoirs: With the arrow on the drive rod pointing up, pull the drive rod out as far as it will go. Remove the empty reservoir and follow instructions in the Storage and Disposal box. Reservoirs that still contain gel can be sealed with the end cap and outlet cap for future use. To attach a new reservoir follow the above directions. Pest population reduction will be apparent within 1 day for German roaches and 3-4 days for American roaches.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store in a cool, dry area out of reach of children. CONTAINER HANDLING: If Empty: Do not reuse container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. If you have questions regarding the use of this product, please call

1-800-331-2867.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once

By using this product, user or buyer accepts the following Conditions. Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CronScience LP. All such risks shall be assumed by the user or buyer. DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, Bayer CropScience LP disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES. INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP 'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd) and MAXFORCE® are registered trademarks of Bayer.

Produced for: **Bayer Environmental Science** A Division of Bayer CropScience LP 2 T. W. Alexander Drive Research Triangle Park, NC 27709

Open this end

ROACH KILLER BAIT GEL







For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577 For PRODUCT USE Information Call 1-800-331-2867

Product may stain textiles and clothing.

| FIRST AID | | | |
|----------------------------|--|--|--|
| If on skin or clothing: | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. | | |
| If in eyes: | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. | | |
| Have the produ | uct container or label with you when calling | | |

a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION Harmful if absorbed through skin. Causes moderate eye irritation.

Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PHYSICAL AND CHEMICAL HAZARDS: Do not use this product in

or on electrical equipment where a possibility of shock hazard exists.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area out of reach of children.

CONTAINER HANDLING: If Empty. Do not reuse container.

Place in trash or offer for recycling if available. If Partly Filled:

Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indeport or united or the product of the product of

indoor or outdoor drain.

If you have questions regarding the use of this product, please call 1-800-331-2867.

See inside leaflet for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Bayer

Produced for: Bayer Environmental Science A Division of Bayer CropScience LP 2 T. W. Alexander Drive Research Triangle Park, NC 27709 Product of China



Net Weight 33g (1.16 Oz) Reservoir

....0.05%99.95%100.00%

STOP - READ THE LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. CAUTION

80183798C 140410AV1






ROACH KILLER BAIT GEL









ROACH KILLER BAIT GEL

Open this end

