

TALAKTM 7.9% F



When used as a Termiticide, Individuals/Firms must be licensed by the State to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the pest control regulatory agency of your State prior to use of this product.

ACTIVE INGREDIENT:	(% by weight)
Bifenthrin*	7.9%
OTHER INGREDIENTS:	92.1%
TOTAL:	100.0%
Contains 2/3 pound active ingredient per gallon.	
EPA Reg. No.: 91234-145	

* Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN CAUTION

See below for additional Precautionary Statements.

FIRST AID

If swallowed: ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by the poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person. **If inhaled:** ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. ▪ Call a poison control center or doctor for further treatment advice. **If on skin or clothing:** ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for treatment advice. **If in eyes:** ▪ Hold eye open and rinse slowly and gently with water 15 - 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continuing rinsing eye. ▪ Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.

Note to Physician - This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution - Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

All pesticide handlers (mixers, loaders, and applicators) must wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves. After the product is diluted in accordance with label directions for use, and/or when mixing and loading using a closed spray tank transfer system, or an in-line injector system, shirt, pants, socks, shoes, and waterproof gloves are sufficient. In addition, all pesticide handlers must wear a respiratory protection device¹ when working in a non-ventilated space. All pesticide handlers must wear protective eyewear when working in non-ventilated space or when applying termiticide by rodding or sub-slab injection.

¹ Use one of the following Mine Safety and Health Administration (MSHA)/National Institute for Occupational Safety and Health (NIOSH) air purifying respirator types with approval number prefixes: TC-23C, TC-21C, TC-19C, TC-13F and TC-14G, or a NIOSH approved respirator with any R, P or HE filter or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE prefilter.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow to drift to blooming crops if bees are visiting the treatment area.

Physical and Chemical Hazards

Do not apply water-based dilutions of **Talak 7.9% F** to electrical conduits, motor housings, junction boxes, switch boxes or other electrical equipment because of possible shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RESTRICTIONS:

- Do not apply by air.
- Do not apply in greenhouses, nurseries.
- Do not apply this product through any kind of irrigation system.
- Not for use on sod farm turf, golf course turf, or grass grown for seed.
- Do not water the treated area to the point of runoff.
- Do not make applications during rain.
- Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack and crevice treatment. During application, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters.

- All outdoor applications, if permitted elsewhere on this label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label:

- 1) Application to soil or vegetation, as listed on this label, around structures;
- 2) Applications to lawns, turf, and other vegetation, as listed on this label;
- 3) Applications to the side of a building, up to a maximum height of 3 feet above grade;
- 4) Applications to the underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
- 5) Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
- 6) Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch, or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests.

Application equipment that delivers low volume treatments, such as Patriot Injector® or Actisol® applicators, may also be used to make crack and crevice, deep harborage, spot and general surface treatments of **Talak 7.9% F**.

When treating adjacent to an existing structure, the applicator must check the area to be treated, and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean-up is completed.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended only for aesthetic purposes or climatic modifications and being grown in interior landscapes, ornamental gardens or parks, or lawns and grounds.

TERMITICIDE USES

Using this product in and around structures and building construction will prevent and control termite infestations.

To institute a barrier between the wood and the termites in the soil, the chemical dilution must be effectively dispersed in the soil. It is important to remove unnecessary materials that contain cellulose and wood from around foundation walls, crawl spaces (inside of structure), and porches, and fix damaged plumbing and construction grade in order to deny termite access to moisture.

To use this product effectively, it is important that the service technician be familiar with current termite control practices including trenching, rodding, sub-slab injection, low-pressure spray application, coarse fan spraying of soil surfaces, crack and crevice (void) injection, excavated soil treatment, and brush or spray applications to infested or susceptible wood. Using these techniques correctly is essential to prevent or control infestations by subterranean termite species of genera *Coptotermes*, *Heterotermes*, *Reticulitermes* and *Zootermopsis*. When determining what procedures to follow, the service technician should consider certain variables such as biology and behavior of the termite species, structure design, heating ventilation, and air conditioning (HVAC) systems, water table, soil type and compaction, grade conditions, and the location and type of domestic water supplies and utilities.

For information concerning the most up to date control practices in a given region or locale, consult the local resources for structural pest control, state cooperative extensions, and regulatory agencies.

SUBTERRANEAN TERMITE CONTROL

Important: The following precautions must be observed to avoid contamination of public and private water supplies:

- Use anti-backflow equipment or procedures to prevent siphonage of insecticide into water supplies.
- Do not contaminate cisterns, wells, or other water tanks by treating the soil beneath these structures.
- Do not treat soil that is water saturated or frozen.
- Do not treat soil where runoff may occur.
- Consult state and local specifications for recommended treatment practices in your area.
- If local or state specifications do not exist, consult the Federal Housing Administration Specifications (H.U.D.) guidance documents.

Note: For the purpose of this label, crawl spaces are defined as being inside of the structure.

Critical Areas: Points at which the foundation is penetrated or abuts another structure are Critical Areas. These include utility entry points, cracks and expansion joints, bath traps and adjacent structures such as stairs, patios and slab additions.

Structures with Wells/Cisterns Inside Foundations

Structures that contain wells or cisterns within the foundation of a structure can only be treated using the following techniques:

1. Do not treat soil while it is beneath or within the foundation or along the exterior perimeter of a structure that contains a well or cistern. The treated backfill method must be used if soil is removed and treated outside/away from the foundation. The treated backfill technique is described as follows:
 - a) Trench and remove soil to be treated onto heavy plastic sheeting or similar material or into a wheelbarrow.
 - b) Treat the soil at the rate of 4 gallons of dilute emulsion per 10 linear feet per foot of depth of the trench, or 1 gallon per 1.0 cubic feet of soil. See **Mixing Directions** section of this label. Mix thoroughly into the soil taking care to contain the liquid and prevent runoff or spillage.
 - c) After the treated soil has absorbed the diluted emulsion, replace the soil into the trench.
2. Treat infested and/or damaged wood in place using an injection technique such as described in the **Control of Wood-infesting Insects** section of this label.

Structures with Adjacent Wells/Cisterns and/or Other Water Bodies

Applicators must inspect all structures with nearby water sources such as wells, cisterns, surface ponds, streams, and other bodies of water and evaluate, at a minimum, the treatment recommendations listed below prior to making an application.

1. Prior to treatment, if feasible, expose the water pipe(s) coming from the well to the structure, if the pipe(s) enter the structure within 3 feet of grade.
2. Prior to treatment, applicators are advised to take precautions to limit the risk of applying the termiticide into subsurface drains that could empty into any bodies of water. These precautions include evaluating whether application of the termiticide to the top of the footer may result in contamination of the subsurface drain. Factors such as depth to the drain system and soil type and degree of compaction should be taken into account in determining the depth of the treatment.

3. When appropriate (i.e., on the water side of the structure), the treated backfill technique (described above) can also be used to minimize off-site movement of termiticide.

Before these techniques are used close to cisterns, wells, or other bodies of water, seek advice from local, state or federal agencies for information on treatment practices that are acceptable in your area.

Application Rate: Use a 0.06% dilution for subterranean termites. For other pests on the label use specific listed rates.

Mixing Directions: Mix the termiticide use dilution in the following manner:

1. Fill tank 1/4 to 1/3 full.
2. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose.
3. Add appropriate amount of **Talak 7.9% F**.
4. Add remaining amount of water.
5. Let pump run and allow recirculation through the hose for 2 to 3 minutes.

Talak 7.9% F may also be combined into full tanks of water. If combined into full tanks of water, allow sufficient time for agitation and/or recirculation to ensure consistency or dilution.

To prepare a 0.06% water dilution, ready to use, dilute 3 quarts of **Talak 7.9% F** with 99.25 gallons of water.

Mixing: Using the chart below, determine the volume of **Talak 7.9% F** and water to produce the desired volume of finished dilution.

Amount of Talak 7.9% F (Gallons Except Where Noted)			
Emulsion Concentrate	Amount of Talak 7.9% F	Amount of Water	Desired Gallons of Finished Emulsion
0.06%	1 oz.	127 oz.	1
	5 oz.	4.9	5
	10 oz.	9.9	10
	25 oz.	24.8	25
	1.5 qt.	49.6	50
	2.25 qt.	74.4	75
	3 qt.	99.25	100
	4.5 qt.	148.8	150
0.12%*	6 qt.	198.5	200
	2 oz.	126 oz.	1
	10 oz.	4.9	5
	19.5 oz.	9.8	10
	1.5 qt.	24.6	25
	3 qt.	49.2	50
	4.5 qt.	73.8	75
	6 qt.	98.5	100
	9 qt.	147.7	150
	3	197	200
*When treating for termites, use this rate only in conjunction with volume adjustments, foam applications or underground services applications as listed below.			
Common units of measure: 1 pint = 2 cups = 16 fluid ounces (oz.) 1 quart = 2 pints = 4 cups = 32 fluid ounces (oz.)			

Application Volume: To provide maximum control and protection against termite infestation apply the specified volume of the finished water emulsion and active ingredient as set forth in the **DIRECTIONS FOR USE** section of this label. If soil will not accept the labeled application volume, the volume may be reduced provided there is a corresponding increase in concentration so that the amount of active ingredient applied to the soil remains the same.

Note: Large reductions of application volume reduce the ability to obtain a continuous barrier. Variance is allowed when volume and concentration are consistent with the label directed rates and a continuous barrier can still be achieved.

The volume of the 0.12% emulsion may be reduced by 1/2 the labeled volume where desirable for pre- and post-construction applications. When the volume is reduced, the hole spacing for subslab injection and soil rodding may also need to be adjusted to account for lower volume dispersal of the termiticide in the soil. Consult the following **Volume Adjustment Chart** for details.

Volume Adjustment Chart		
Rate (% Emulsion)	0.06%	0.12%
Volume allowed		
Horizontal (gallons emulsion/10 ft ²)	1.0 gallons	0.5 gallons
Vertical (gallons emulsion/10 linear ft.)	4.0 gallons	2.0 gallons

After Treatment: All holes in commonly occupied areas into which **Talak 7.9% F** has been applied must be plugged. Plugs must be of a non-cellulose material or covered by an impervious, non-cellulose material.

Foam Applications

Talak 7.9% F emulsion, from 0.06 to 0.12% may be converted to foam with 2X - 40X expansion characteristics and used to control or prevent termite infestations.

Depending on the circumstances, foam applications may be used alone or in combination with liquid emulsion applications. Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block voids or structural voids, under slabs, stoops, porches, or to the soil in crawlspaces, and other similar voids.

Foam and liquid application must be consistent with volume and active ingredient instructions in order to insure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the labeled liquid emulsion volume of product must be applied, with the remaining percent delivered to appropriate areas using foam application. Refer to label and use recommendations of the foam manufacturer and the foaming equipment manufacturer.

Foam applications are generally a good supplement to liquid treatments in difficult areas, but may be used alone in difficult spots.

Application Under Slabs or to Soil in Crawlspaces to Prevent or Control Termites

When making applications, **Talak 7.9% F** foam can be used alone or in combination with liquid emulsion. Whether applied as a emulsion, foam, or some of both, the equivalent of at least 4 gallons of 0.06% emulsion (4 ounces of **Talak 7.9% F** concentrate) per 10 linear feet must be applied for vertical barrier, or at least 1 gallon of 0.06% emulsion (1 ounce of **Talak 7.9% F** concentrate) per 10 square feet must be applied for a horizontal barrier. For a foam only application, apply **Talak 7.9% F** concentrate in sufficient concentration and volume to equal 4 ounces of concentrate per 10 linear feet or 1 ounce of concentrate per 10 square feet. For example, 2 gallons of 0.12% emulsion converted to foam and used to cover 10 linear feet is the equivalent of 4 gallons of 0.06% emulsion per 10 linear feet.

Sand Barrier Installation and Treatment

As long as termites have access to soil that has not been treated and can avoid soil that has been treated with **Talak 7.9% F**, they can build mud tubes over surfaces that have been treated. Cracks and spaces should be filled with play box or builder's sand and then treated in the same manner as soil. Follow the rates listed in the **Talak 7.9% F** label.

Retreatment for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide barrier in the soil. These vulnerable or reinfested areas may be retreated in accordance with application techniques described in this product's labeling. The timing and type of these retreatments will vary depending on factors such as termite pressure, soil types, soil conditions and other factors which may reduce the effectiveness of the barrier.

Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.

Application in Conjunction with Termite Baits

As part of an IPM (integrated pest management) program for termite control, **Talak 7.9% F** may be applied to areas of the structure with known or suspected infestations such as plumbing, utility entry sites, bath traps, expansion joints, and foundation cracks at a rate of 0.06% as a spot treatment or complete barrier treatment. Applications may be made as described in the **Post-construction Subterranean Termite Treatment** section of this label.

Pre-construction Subterranean Termite Treatment

Use Precautions:

- The treatment site must be covered prior to a rain event in order to prevent run-off of the pesticide into non-target areas.
- The applicator must either cover the soil him/herself or provide written notification of the above requirement to the contractor on site and to the person commissioning the application (if different than the contractor). If notice is provided to the contractor or person commissioning the application, then they are responsible under FIFRA to ensure that: 1) if the concrete slab cannot be poured over the treated soil within 24 hours of application the treated soil is covered with a waterproof covering (such as polyethylene sheeting), and 2) the treated soil is covered if precipitation is predicted to occur before the concrete slab is scheduled to be poured.

Use Restrictions:

- Do not treat soil that is water-saturated or frozen.
- Do not treat when raining.
- Do not allow treatment to runoff from the target area.
- Do not apply with 10 feet of storm drains. Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or ponds; estuaries; and commercial fish farm ponds).
- Do not make on-grade applications when sustained wind speeds are above 10 mph (at application site) at nozzle end height.
- Do not apply at a lower dosage and/or concentration than specified on this label for applications prior to the installation of the finished grade.

When treating foundations deeper than 4 feet, apply the termiticide as the backfill is being replaced, or if the construction contractor fails to notify the applicator to permit this, treat the foundation to a minimum depth of 4 feet after the backfill has been installed. The applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements, at the rate prescribed from grade to a minimum depth of 4 feet. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the

bottom of the footing. However, in no case should a structure be treated below the footing.

To produce effective pre-construction subterranean termite control, create vertical and/or horizontal chemically treated zones of protection using 0.06% emulsion of **Talak 7.9% F**. Follow the current edition of the Housing and Urban Development Minimum Property Standards to assure that F.H.A. termite-proofing requirements are met.

Horizontal Barriers

Establish a horizontal chemical barrier wherever treated soil will be covered by a slab, such as basement floors, carports, entrance platforms, footing trenches, and slab floors and the soil below stairs and crawl spaces.

Apply 1 gallon of 0.06% emulsion per 10 square feet, or use 1 fluid ounce of **Talak 7.9% F** per 10 square feet in sufficient water (no less than 1/2 gallon or more than 2 gallons) to provide a uniform treated barrier for the area being treated.

If the fill is coarse aggregate, such as washed gravel, a sufficient volume of emulsion must be applied to allow it to reach the soil beneath for coarse fill.

Make applications with a low-pressure spray (less than 50 p.s.i.) using a coarse spray nozzle. If foundation walls have not been installed around the treated soil and the slab will not be poured the same day as treatment, the treated soil must be covered with a water-proof barrier. Polyethylene sheeting may be used for this purpose.

Vertical Barriers

Vertical barriers must be established in Critical Areas such as along the inside of foundation walls, plumbing, bath traps, utility services and other features that will penetrate the slab.

Using a 0.06% emulsion, apply 4 gallons of emulsion per 10 linear feet per foot of depth or 4 fluid ounces of **Talak 7.9% F** per 10 linear feet per foot of depth from grade level to the top of the footing in sufficient water (not less than 2 gallons or more than 8 gallons) to provide a uniform treated barrier.

When trenching and rodding into the trench, or trenching, ensure that the emulsion reaches the top of the footing. Space rod holes so that a continuous treated barrier is created, but not exceeding 12 inches apart. Avoid soil wash-out around the footing. Trenches should be about 6 inches wide and 6 inches deep. Mix the chemical emulsion with the soil as it is being replaced in the trench. Inside vertical barriers may not be required for monolithic slabs.

When treating hollow block voids, use 2 gallons of emulsion per 10 linear feet to assure that the emulsion reaches the top of the footing.

Prior to each application, applicators must notify the general contractor, construction superintendent, or similar responsible party, of the intended termiticide application and intended sites of application and instruct the responsible person to notify construction workers and other individuals to leave the area to be treated during application and until the termiticide is absorbed into the soil.

Post-construction Subterranean Termite Treatment

For a post-construction treatment, use a 0.06% emulsion. Post-construction treatments shall be made by injection, trenching and rodding into the trench or trenching using a low-pressure spray not exceeding 25 p.s.i. at the nozzle. Proper precautions should be taken to avoid soil wash-out around the footing.

Locate, identify, and mark wells, electrical conduits, water and sewer lines, and radiant heat pipes prior to application of **Talak 7.9% F**. Do not puncture or inject **Talak 7.9% F** into these structures.

Foundations

For applications made after the final grade is installed, the applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements, at the rate prescribed from grade to the top of the footing. When

the footing is more than four (4) feet below grade, the applicator must trench and rod into the trench or trench along the foundation walls at the rate prescribed to a minimum depth of four feet. The actual depth of treatment will vary depending on soil type, degree of compaction, and location of termite activity. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

Slabs

Create vertical barriers by trenching and rodding into the trench or trenching outside at the rate of 4 gallons of emulsion per 10 linear feet per foot of depth and by sub-slab injection within the structure. Ensure an even distribution of chemical. Applications must not be made below the bottom of the footing.

Apply beside the outside of the foundation and under the slab on the inside of foundation walls, where needed. Treatment of slabs may also be necessary under and beside both sides of any interior footing-supported walls, in all cracks and expansion joints, and beside one side of interior partitions. By long-rodding or grid pattern injection vertically through the slab, horizontal barriers may be created where necessary.

- To permit the creation of an uninterrupted insecticidal barrier, drill holes in the foundation and/or slab.
- For foundations that are less than or equal to 1 foot, dig a narrow trench about 6 inches wide beside the outside of the foundation walls. Do not dig beneath the bottom of the footing. As the soil is placed back into the trench, apply 4 gallons of emulsion per 10 linear feet per foot of depth to the trench and soil.
- For foundations that are deeper than 1 foot, follow the rates stated above for basements.
- A 0.06% emulsion may be used to treat exposed and wood in bath traps.

Basements

Treatment must be made by trenching and rodding into the trench, or trenching at the rate of 4 gallons of emulsion per 10m linear feet per foot of depth wherever the footing, from grade to the bottom of the foundation, is greater than 1 foot of depth. When the footer is greater than four feet below grade, the applicator may trench and rod into the trench, or trench beside foundation walls at the rate designated for four feet of depth. Space rod holes to create a continuous insecticidal barrier, but in no case more than 12 inches apart. Depending on the type of soil, degree of compaction and location of termite activity, the actual depth of treatment will differ. However, a structure should never be treated below the footer. Sub-slab injection may be needed beside the inside of foundation walls, around conduits, piers, and pipes, beside both sides of interior footing-supported walls, and beside cracks and partition walls.

Accessible Crawl Spaces: For crawl spaces, apply vertical termiticide barriers at the rate of 4 gallons of emulsion per 10 linear feet per foot of depth from grade to the top of the footing, or if the footing is more than 4 feet below grade, to a minimum depth of 4 feet. Apply by trenching and rodding into the trench, or trenching. Treat both sides of foundation and around all piers and pipes. Where physical obstructions such as concrete walkways adjacent to foundation elements prevent trenching, treatment may be made by rodding alone. When soil type and/or conditions make trenching prohibitive, rodding may be used. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. Read and follow the mixing and use direction section of the label if situations are encountered where the soil will not accept the full application volume.

- Rod holes and trenches must not extend below the bottom of the footing.
- Rod holes must be spaced so as to achieve a continuous termiticide barrier but in no case more than 12 inches apart.

3. Trenches must be a minimum of 6 inches deep or to the bottom of the footing, whichever is less, and need not to be wider than 6 inches. When trenching in sloping (tiered) soil, the trench must be stepped to ensure adequate distribution and prevent termiticide from running off. The emulsion must be mixed with the soil as it is replaced in the trench.
4. When treating plenums or crawl spaces, turn off the air circulation system of the structure until application has been completed and all termiticide has been absorbed by the soil.

Inaccessible Crawl Spaces: For inaccessible interior areas, such as areas where there is insufficient clearance between floor joists and ground surfaces to allow operator access, excavate if possible, and treat according to the instruction for accessible crawl spaces. Otherwise, apply one or a combination of the following two methods.

1. To establish a horizontal barrier, apply to the soil surface, 1 gallon of emulsion per 10 square feet overall using a nozzle pressure of less than 25 p.s.i. and a coarse application nozzle (e.g., Delavan® Type RD Raindrop®, RD-7 or larger, or Spraying Systems Co. 8010LP TeeJet® or comparable nozzle). For an area that cannot be reached with the application wand, use one or more extension rods to make the application to the soil. Do not broadcast or powerspray with higher pressures.
2. To establish a horizontal barrier, drill through the foundation wall or through the floor above and treat the soil perimeter at a rate of 1 gallon of emulsion per 10 square feet. Drill spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals, so check state regulations which may apply.

When treating plenums and crawl spaces, turn off the air circulation systems of the structure until application has been completed and all termiticide has been absorbed by the soil.

Masonry Voids: Drill and treat voids in multiple masonry elements of the structure extending from the structure to the soil in order to create a continuous treatment barrier in the area to be treated. Apply at a rate of 2 gallons of emulsion per 10 linear feet of footing, using a nozzle pressure of less than 25 p.s.i. When using this treatment, access holes must be drilled below the sill plate and should be as close as possible to the footing as is practical. Treatment of voids in block or rubble foundation walls must be closely examined. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy the contaminated areas of the structure until the clean-up is completed.

Note: When treating behind veneer structures (walls, etc.) take proper care to not drill beyond the veneer. If concrete blocks exist behind the veneer, both can be drilled and treated simultaneously.

Talak 7.9% F may not be used in voids insulated with rigid foam insulation.

Excavation Technique: If treatment must be made in difficult situations, along field-stone or rubble walls, along faulty foundation walls, and around pipes and utility lines which lead downward from the structure to a well or pond, apply using the following technique:

- a. Prepare a trench, placing the removed soil into heavy-weight plastic sheeting or similar, water-impermeable material.
- b. Treat the soil with 4 gallons of emulsion per 10 linear feet per foot of depth of the trench. Completely mix the emulsion into the soil exercising care to avoid liquid running off the sheeting.
- c. Place the treated soil back into the trench after it has absorbed the emulsion.

Attention: Wear MSHA/NIOSH approved unvented goggles and a respirator when applying **Talak 7.9% F** in a confined area.

SPECIFIC PEST CONTROL APPLICATIONS

Underground Services: Wires, cables, utility lines, pipes, conduits, etc. Services may be within structures, in right-of-ways or to protect long range (miles) of installations of services.

To prevent attack by termites and ants, apply 0.06 to 0.12% **Talak 7.9% F** emulsion to the soil.

Apply to the bottom of the trench at the rate of 2 gallons of emulsion per 10 linear feet. Let the emulsion be absorbed into the soil. Lay services on the treated soil and cover with about 2 inches of fill soil. Apply another 2 gallons per 10 linear feet over the soil surface to complete the chemical barrier. In wide trenches, only the soil near the services should be treated. A continuous barrier of treated soil surrounding the services must be established.

In cases where the soil will not accept the above labeled volume, 1 gallon of 0.12% **Talak 7.9% F** may be used per 10 linear feet of trench. Apply both to the bottom of the trench and over the soil on top of the services.

Fill the trench with treated fill soil. Treat the soil where each service sticks out from the ground by trenching/rodding of not more than 1 to 2 gallons of emulsion into the soil.

Precautions: Do not treat electrically active underground services.

Posts, Poles, and Other Constructions: Around wooden constructions such as signs, fences and landscape ornamentation an insecticidal barrier can be established by treating with a 0.06% emulsion. Sub-surface injection and gravity-flow through holes in the bottom of the trench, are two treatment methods that can be used on poles and posts that have already been installed. A complete chemical barrier around the pole can be established by treating on all sides. For poles and posts less than six inches in diameter use 1 gallon of emulsion per foot of depth and for larger poles, use 1.5 gallons of emulsion per foot of depth. Apply to a depth of 6 inches below the bottom of the wood. For larger constructions, use 4 gallons per 10 linear feet per foot of depth.

Treatment of Wood-in-Place for Control of Wood-infesting Insects (Localized Areas in Structure): For the control of insects such as Termites, Ants, Carpenter Ants, and wood-infesting beetles such as Old House Borer and Powder Post in localized areas of infested wood in and around structures, apply a 0.06% emulsion to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is at risk. Paint on or fan spray applications may also be used. Place plastic sheeting under overhead areas that are spot treated except for soil surfaces in crawl spaces. Areas in which access is difficult may be treated by drilling, and then injecting the emulsion with a crack and crevice injector into the damaged wood or void spaces. This type of application is not intended to be a substitute for soil treatment, mechanical alteration or fumigation to control extensive infestation of wood-infesting insects.

Termite Carton Nests in Trees or Building Voids: To control termite carton nests in trees or building voids inject with a 0.06% emulsion. Multiple injection points to varying depths may be necessary. Carton nest material should be removed from building voids when nests are discovered.

Control of Bees, Wasps Hornets, and Yellow Jackets Indoors: To control Bees, Wasp, Hornets, and Yellow-Jackets, apply a 0.06% emulsion. Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity.

Important: Do not apply emulsion until location of heat pipes, ducts, water and sewer lines and electrical conduits are known and identified. Caution must be taken to avoid puncturing and injection into these structural elements. Do not apply into electrical fixtures, switches, or sockets.

In the home, all food processing surfaces and utensils in the treatment area should be covered during treatment or thoroughly washed before re-use. Remove pets, birds, and cover aquariums before spraying. Do not permit humans or pets to contact treated surfaces until the spray has dried.

During any overhead applications to overhead interior areas of structures, cover surfaces below with plastic sheeting or similar materials.

Wear protective clothing, unvented goggles, gloves and respirator, when applying to overhead areas or in poorly ventilated areas. Avoid touching sprayed surfaces until spray has completely dried.

Outside of Structures

Broadcast Treatment of Wood for the Control of Wood-infesting Insects and Nuisance Pests: In order to control wood-infesting insects active inside trees, utility poles and/or fence posts, drill to the interior infested cavity and inject a 0.06% emulsion. If treating nuisance pests on the exterior of the structure, apply a 0.06% emulsion with a fan spray using a maximum pressure of 25 p.s.i. and apply just to the point of run-off. To control Bees, Wasps, Hornets, and Yellow-Jackets, direct the spray at nest openings in the ground, bushes, and in cracks and crevices. Saturate nest openings and contact as many insects as possible. Apply in late evening when insects are at rest.

Pests Under Slabs: To control infestations of Arthropods, including ants, cockroaches and scorpions living beneath the slab area, drill and inject, or horizontal rodding and then inject 1 gallon of a 0.06% to 0.12% emulsion per 10 square feet or 2 gallons of emulsion per 10 linear feet.

Formula for Determining the Active Ingredient Content of the Finished Spray Mixture: The following formula may be used to determine the percent active ingredient that is in the spray tank after mixing **Talak 7.9% F**:

$$\frac{(7.9\%) \times (\text{fl. oz. of Talak 7.9\% F Added to Tank})}{(\text{Gallons of Finished Spray Mix}) \times (128)} = \text{Percent Active Ingredient of Spray Mix}$$

LAWNS AND ORNAMENTALS

Application Instructions

Talak 7.9% F may be mixed with water and other aqueous carriers for the control of insects and mites on trees, shrubs, foliage plants, non-bearing (perennial crops that will not produce a harvestable raw agricultural commodity during the season of application) fruit and nut trees, and flowers in interiorscapes including hotels, shopping malls, office buildings, etc., and outdoor plantscapes, such as around residential dwellings, parks, institutional buildings, recreational areas, athletic fields and home lawns.

Talak 7.9% F may be tank-mixed with insect growth regulators as well as other pesticides. Observe all precautions and the directions for use for each tank mix product. Physical compatibility of **Talak 7.9% F** may vary with different combinations of products, and local cultural practices. Prepare a small scale (pint or quart jar) test sample for any combination not tested previously. Use the proper proportions of pesticides and water to make sure of the compatibility of the mixture.

Tank Mix Compatibility Testing

A jar test is recommended prior to tank mixing to ensure compatibility of **Talak 7.9% F** and other products. Use a clear glass quart jar with lid and mix the tank mix ingredients in their relative proportions. Always use water from the intended source. Invert the jar

containing the mixture several times and observe the mixture for approximately 1/2 hour. Evaluate the solution for uniformity and stability. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combination should not be used.

Tank Mix Preparation:

Unless otherwise noted, follow the procedure below to prepare a tank mix:

- 1) Add wettable powders to tank water
- 2) Agitate
- 3) Add liquids and flowables
- 4) Agitate
- 5) Add emulsifiable concentrates
- 6) Agitate

Formula for Determining the Active Ingredient Content of the Finished Spray Mixture: The following formula may be used to determine the percent active ingredient that is in the spray tank after mixing **Talak 7.9% F**:

$$\frac{(7.9\%) \times (\text{fl. oz. of Talak 7.9\% F Added to Tank})}{(\text{Gallons of Finished Spray Mix}) \times (128)} = \text{Percent Active Ingredient of Spray Mix}$$

Resistance: Some insects may develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state pest management authorities for details.

If resistance to this product develops in your area, this product, or other products with similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and suspect that resistance is a reasonable cause, immediately consult your local company representative or pest management advisor for the best alternative method of control in your area.

In the State of New York, for application uses outdoors on ornamentals and lawns in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields:

The Following Precautionary Measures Must be Obeyed.

A 100 foot buffer must be maintained between the application site and the waters of the State. A 100 foot buffer is required for all waters except those entirely privately owned with no outlet to State waters. The buffer must consist of well maintained, established vegetation (i.e. grass, etc.) growth and must be maintained to prevent the development of channels.

In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but no sooner than two weeks after the first application.

Lawn Application

Apply **Talak 7.9% F** as a broadcast treatment. For uniform control when applying to dense grass foliage, use volumes of up to 10 gallons per 1,000 square feet.

To ensure control of sub-surface pests, including Mole Crickets, using low volume applications (less than 2 gallons per 1,000 square feet) immediately follow treatment with irrigation of treated area with at least 0.25 inches of water.

Application Rates

Under typical conditions, the application rates shown in the table below will provide control of the listed pests. Follow the application rates listed in the table below for typical pest pressure. **Talak 7.9% F** may be applied at up to 1 fl. oz. per 1,000 square feet at the discretion of the applicator. Use the higher application rate for maximum residual control.

Pest	Application Rate	Application Instructions
Armyworms Cutworms Sod Webworm	0.18 - 0.25 fl. oz. per 1,000 sq. ft.	For best results, postpone watering (irrigation) or mowing for 24 hours after application. Higher treatment rates (up to 1 fluid oz. per 1,000 square feet) may be necessary if high pest pressure exists and grass is maintained taller than 1 inch.
Annual Bluegrass Weevil (Hyperodes) (Adult) Banks Grass Mite Billbugs (Adult) Black Turfgrass Ataenius (Adult) Centipedes Chinch Bugs Crickets Earwigs Fleas (Adult) Grasshoppers Leafhoppers Mealybugs Millipedes Mites Pillbugs Sowbugs	0.25 - 0.50 fl. oz. per 1,000 sq. ft.	Annual Bluegrass Weevil (<i>Hyperodes</i>) Adults: Treatment should be timed as they travel into grass away from their overwintering sites. Travel usually begins when <i>Forsythia</i> is in full bloom and concludes when flowering dogwood (<i>Cornus florida</i>) is in full bloom. For additional detailed information, check with your State Cooperative Extension Service. Billbug Adults: Treatment should be made when adult billbugs are first noticed in April and May. To optimize treatment, degree day models have been developed. For additional detailed information, check with your State Cooperative Extension Service. Spring treatments for billbug adults will also offer control of over-wintered chinch bugs in temperate climates. Black Turfgrass Ataenius Adults: To control the first and second generation of black turfgrass ataenius adults, respectively, treatments should take place in May and July. Time the May treatment to coincide with the full bloom stage of Vanhoutte spiraea (<i>Spiraea vanhoutte</i>) and horse chestnut (<i>Aesculus hippocastanum</i>). Time the July treatment to coincide with the blooming Rose of Sharon (<i>Hibiscus syriacus</i>). Chinch Bugs: Mostly found in the thatch layer, Chinch bugs infest the base of grass plants. In order to optimize the penetration of the insecticide to location of the chinch bugs, irrigation of the grass prior to treatment may be necessary. If the grass is kept at a long mowing height or if the thatch layer excessive, use higher volume treatments. It may be necessary to use higher application rates (up to 1 fluid oz. per 1,000 square feet) to control Chinch Bug populations made up of both nymphs and adults in mid summer. Mites: Apply in combination with a labeled application rate of a surfactant to achieve optimal control of eriophyid mites. A second application may be needed, five to seven days after the first, to ensure optimal control.
Crane Flies	0.5 fl. oz. per 1,000 sq. ft.	Applications should be made August - February to control early to mid-season larvae as they feed on plant crowns. Applications made March - April to late-season larvae may aid in suppression.
Ants Fleas (Larvae) Imported Fire Ants Japanese Beetle (Adult) Mole Cricket (Adult) Mole Cricket (Nymph) Ticks	0.5 - 1.0 fl. oz. per 1,000 sq. ft.	Flea Larvae: Flea larvae mature in shaded areas accessible to pets or other animals. When treating these areas use a higher volume treatment so that the insecticide penetrates into the soil. Note: If the lawn area is being treated with Talak 7.9% F at 0.25 fluid ounces per 1,000 square feet for adult flea control, then the larval application rate can be accomplished by a two- to four-fold increase in spray volume. Imported Fire Ants: For best control use broadcast treatments in combination with mound drenches. This will control present colonies along with foraging workers and newly mated fly-in queens. It is critical either to use high volume treatments or to irrigate prior to application if the soil is dry. Apply 1 fluid oz. per 1,000 square feet when using broadcast treatments. For mound drenches, dilute 1 teaspoon of Talak 7.9% F per gallon of water and use 1 to 2 gallons of finished spray per mound using sufficient force to penetrate the top and allow the emulsion to flood ant channels. Treat a four foot diameter around each ant mound. Application should be made in late evening or early morning when it is cooler (65 - 80°F). Note: A spray rig calibrated to apply 1 fluid oz. per 1,000 square feet of Talak 7.9% F in 5 gallons per 1,000 square feet contains the equivalent emulsion (1 teaspoon per gallon) required for fire ant mound drenches in the spray tank. Mole Cricket Adults: Since the preferred grass areas are subject to constant invasion in early spring by the active adult stage, it can be difficult to maintain control of adult mole crickets. Make treatments as late in the day as possible and follow with up to 0.5 inches of water after treatment. To ensure maximum contact with the insecticide when the soil is dry, irrigate prior to treatment to bring the adult mole crickets closer to the soil surface. To obtain optimal control of potential nymphal populations, the grass areas preferred by adult mole crickets in the spring should be treated immediately prior to peak hatch stage. (See below).

(continued)

Pest	Application Rate	Application Instructions
Ants Fleas (Larvae) Imported Fire Ants Japanese Beetle (Adult) Mole Cricket (Adult) Mole Cricket (Nymph) Ticks	0.5 - 1.0 fl. oz. per 1,000 sq. ft.	Mole Cricket Nymphs: To obtain optimal control of potential nymphal populations, the grass areas preferred by adult mole crickets in the spring should be treated immediately prior to peak hatch stage. Young nymphs are more vulnerable to insecticidal treatment at this stage because they are close to the soil surface where the insecticide is most effective. Use higher application rates and frequent applications to control larger, more damaging, nymphs later in the year. Make treatments as late in the day as possible and water immediately with up to 0.5 inches of water. To ensure maximum contact with the insecticide when the soil is dry, irrigate prior to treatment to bring the adult mole crickets closer to the soil surface. Ticks (including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever): Make application to the entire area where contact with ticks may occur. Do not make spot treatments. When applying to areas with heavy leaf litter or dense ground cover use higher spray volumes. To attain and/or sustain control in times of high pest pressure, retreatments may be necessary; retreat only if signs of continued or renewed tick activity are present. Repeat treatments must not be made more often than once per seven days. Deer Ticks (<i>Ixodes</i> sp.) have a four-stage life cycle spanning 2 years. Treat in the late fall and/or early spring to control adult ticks located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that live in the soil and leaf litter. American Dog Ticks invade suburban settings in areas where residences and dwellings are constructed on former fields or wooded areas. These pests normally gather by paths or roadways where they are likely to find a host. To control tick larvae, nymphs and adults, treatments should take place, as needed, from mid spring to early fall.

Calculating Dilution Rates: To determine the proper dilution of **Talak 7.9% F** that is required to control specific pests, follow the steps below:

- 1) Determine the target pest requiring the highest application rate for effective control in the **Application Rates** table.
- 2) Choose the treatment rate in terms of fluid oz. of **Talak 7.9% F**.
- 3) Determine the dilution volume necessary for the treatment in the **Dilution Chart**.
- 4) Use the proper amount of **Talak 7.9% F** that must be mixed in your desired volume of water as shown in the **Dilution Chart**.

For example, to control ticks the **Application Rates** table shows that 0.5 to 1.0 fluid ounces of **Talak 7.9% F** must be applied per 1,000 square feet. You select an application rate of 1.0 fluid oz. per 1,000 square feet because maximum residual control is desired. Your application volume is approximately 10 gallons per 1,000 sq. ft. Consulting the **Lawn Application Dilution Chart** reveals that you should dilute 1.0 fluid oz. of **Talak 7.9% F** in 10 gallons of water.

Lawn Application Dilution Chart

Application Volume	Application Rate	Fluid Ounces* of Talak 7.9% F Diluted to these Volumes of Finished Spray			
Gallons per 1,000 sq. ft.	Fluid Ounces per 1,000 sq. ft.	1 gallon	5 gallons	10 gallons	100 gallons
1.0	0.18	0.18	0.90	1.8	18.0
1.0	0.25	0.25	1.25	2.5	25.0
1.0	0.5	0.5	2.5	5.0	50.0
1.0	1.0	1.0	5.0	10.0	100.0
2.0	0.18	-	0.45	0.90	9.0
2.0	0.25	0.13	0.63	1.25	12.5
2.0	0.5	0.25	1.25	2.5	25.0
2.0	1.0	0.5	2.5	5.0	50.0
3.0	0.18	-	0.30	0.60	6.0
3.0	0.25	-	0.42	0.83	8.3
3.0	0.5	0.17	0.83	1.67	16.7
3.0	1.0	0.33	1.67	3.33	33.3
4.0	0.18	-	0.23	0.45	4.5
4.0	0.25	-	0.31	0.63	6.3
4.0	0.5	0.13	0.63	1.25	12.5
4.0	1.0	0.25	1.25	2.5	25.0

(continued)

Lawn Application Dilution Chart *(continued)*

Application Volume	Application Rate	Fluid Ounces* of Talak 7.9% F Diluted to these Volumes of Finished Spray			
Gallons per 1,000 sq. ft.	Fluid Ounces per 1,000 sq. ft.	1 gallon	5 gallons	10 gallons	100 gallons
5.0	0.18	-	0.18	0.36	3.6
5.0	0.25	-	0.25	0.5	5.0
5.0	0.5	0.1	0.5	1.0	10.0
5.0	1.0	0.2	1.0	2.0	20.0
10.0	0.18	-	-	0.18	1.8
10.0	0.25	-	0.13	0.25	2.5
10.0	0.5	-	0.25	0.5	5.0
10.0	1.0	0.1	0.5	1.0	10.0

* To convert fluid ounces to milliliters, multiply by 29.57
1 fluid ounce = 29.57 ml = 2 tablespoons = 6 teaspoons
Do not use household utensils to measure **Talak 7.9% F**.

Ornamentals and Trees Application

Talak 7.9% F is for use on trees, shrubs, ground covers, bedding plants, and foliage plants. Treat with 0.125 to 1.0 fl. oz. of **Talak 7.9% F** per 1,000 square feet or 5.4 to 43.5 fl. oz. per 100 gallons. **Talak 7.9% F** may be diluted and used in different volumes of water as long as the maximum label rate (1.0 fluid oz. per 1,000 square feet or 43.5 fl. oz. per 100 gallons) is not exceeded. If diluted with water or other carriers, **Talak 7.9% F** may be applied through low volume application equipment as long as the maximum label rate (1.0 fluid oz. per 1,000 square feet or 43.5 fl. oz. per 100 gallons) is not exceeded.

Treat as a full coverage foliar spray using the stated application rate. If pest pressure and density of foliage increases, repeat treatments using higher rates may be needed to reach the desired control. Repeat treatments must not be made more often than once per 7 days.

Before application to entire planting, test treat a small number of plants and watch for signs of sensitivity. Some plant species may be sensitive to the final spray solution.

To avoid or delay pest resistance, it is recommended to use an alternate class of pesticide in any application program.

In the State of New York, for application uses outdoors on ornamentals and lawns in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields:

The Following Precautionary Measures Must be Obeyed.

A 100 foot buffer must be maintained between the application site and the waters of the State. A 100 foot buffer is required for all waters except those entirely privately owned with no outlet to State waters. The buffer must consist of well maintained, established vegetation (i.e. grass, etc.) growth and must be maintained to prevent the development of channels.

In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but no sooner than two weeks after the first application.

Application Rates

Under typical conditions, the application rates in the table below will offer optimal control of the listed pests. **Talak 7.9% F** may be applied at up to 1 fl. oz. per 1,000 square feet (43.5 oz. per 100 gallons) at the discretion of the applicator. When maximum residual control is preferred, use the higher treatment rates.

Pest	Application Rate	Application Instructions
Bagworms Cutworms Elm Leaf Beetles Fall Webworms Gypsy Moth Caterpillars Lace Bugs Leaf Feeding Caterpillars Tent Caterpillars	0.125 - 0.250 fl. oz. per 1,000 sq. ft. (5.4 - 10.8 fl. oz. per 100 gal.)	Bagworms: For optimum control treat when larvae have started to hatch and are young, directing spray to contact as many larvae as possible.

(continued)

Pest		Application Rate	Application Instructions
Adelgids	Leafrollers	0.25 - 0.50 fl. oz. per 1,000 sq. ft. (10.8 - 21.7 fl. oz. per 100 gal.)	Beetles, Scale Crawlers, Twig Borers, and Weevils: Apply to plant foliage; also treat trunks, stems and twigs.
Ants	Mealybugs		
Aphids	Millipedes		
Bees	Mites		
Beet Armyworm	Mosquitoes		
Beetles	Orchid Weevil		
Black Vine Weevil (Adults)	Pillbugs		
Broad Mites	Pine Needle Scales (Crawlers)		
Brown Soft Scales	Plant Bugs		
Budworms	(Including <i>Lygus</i> spp.)		
California Red Scale (Crawlers)	Psyllids		
Centipedes	San Jose Scales (Crawlers)		
Cicadas	Scorpions		
Citrus Thrips	Sowbugs		
Clover Mites	Spider Mites		
Crickets	Spiders		
Diaprepes (Adults)	Spittlebugs		
Earwigs	Thrips		
European Red Mite	Tip Moths		
Flea Beetles	Treehoppers		
Fungus Gnats (Adults)	Twig Borers		
Grasshoppers	Wasps		
Japanese Beetle (Adult)	Weevils		
Leafhoppers	Whiteflies		
Imported Fire Ants**		0.5 - 1.0 fl. oz. per 1,000 sq. ft. (21.7 - 43.5 fl. oz. per 100 gal.)	Spider Mites: Apply during spring and summer for most effective control of twospotted spider mites. During mid- to late-summer it may be necessary to make more frequent treatments, possibly at higher rates to achieve suitable control. Control may be enhanced by adding a surfactant or horticultural oil or by combining Talak 7.9% F with other products registered to control mites. Applications of Talak 7.9% F may be alternated with chemicals offering other modes of action in programs that are designed to manage resistance by twospotted spider mites. For recommendations on resistance management in your region check with your local Cooperative Extension Service.
Leafminers			
Pecan Leaf Scorch Mite			
Pine Shoot Beetle (Adults)			
Spider Mites			
** For foraging ants.			

Calculating Dilution Rates: To determine the proper dilution of **Talak 7.9% F** that is required to control specific pests, follow the steps below:

- 1) Determine the target pest requiring the highest application rate for effective control in the **Application Rates** table.
- 2) Choose the treatment rate in terms of fluid oz. of **Talak 7.9% F**.
- 3) Determine the dilution volume necessary for the treatment in the **Dilution Chart**.
- 4) Use the proper amount of **Talak 7.9% F** that must be mixed in your desired volume of water as shown in the **Dilution Chart**.

For example, to control black vine weevil adults on rhododendron, the **Application Rates** table shows that 0.25 to 0.5 fluid ounces of **Talak 7.9% F** must be applied per 1,000 square feet. You select an application rate of 0.5 fluid oz. per 1,000 square feet because maximum residual control is desired. Your application volume is approximately 300 gallons per acre, which is equivalent to 6.9 gallons per 1,000 square feet. Consulting the **Ornamental Application Dilution Chart** reveals that you should dilute 0.72 fluid oz. of **Talak 7.9% F** in 10 gallons of water.

Ornamental Application Dilution Chart

Application Volume Gallons per		Application Rate fl. oz. per	Fluid Ounces* of Talak 7.9% F Diluted to these Volumes of Finished Spray			
1,000 sq. ft.	Acre	1,000 sq. ft.	1 gallon	5 gallons	10 gallons	100 gallons
2.3	100	0.125	–	0.27	0.54	5.4
2.3	100	0.25	0.11	0.54	1.08	10.8
2.3	100	0.5	0.22	1.09	2.17	21.7
2.3	100	1.0	0.44	2.17	4.35	43.5
4.6	200	0.125	–	0.14	0.27	2.7
4.6	200	0.25	–	0.27	0.54	5.4
4.6	200	0.5	0.11	0.54	1.09	10.9
4.6	200	1.0	0.22	1.09	2.17	21.7
6.9	300	0.125	–	–	0.18	1.8
6.9	300	0.25	–	0.18	0.36	3.6
6.9	300	0.5	–	0.36	0.72	7.2
6.9	300	1.0	0.15	0.72	1.45	14.5

*To convert fluid ounces to milliliters, multiply by 29.57

300 gallons per acre is a typical application volume for landscape ornamental applications.

1 fluid ounce = 29.57 ml = 2 tablespoons = 6 teaspoons

Do not use household utensils to measure **Talak 7.9% F**.

Pest Control on Outside Surfaces and Around Buildings

Apply **Talak 7.9% F** to outside surfaces of buildings including exterior siding, foundations, porches, window frames, eaves, patios, garages, refuse dumps, lawns such as grass areas adjacent or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, fence lines, storage sheds, barns, and other residential and non-commercial structures, soil, trunks of woody ornamentals and other areas where pests congregate or have been seen.

Talak 7.9% F may be used to control the following pests:

Ants	Chinch Bugs	Gnats	Silverfish
Carpenter Ants	Clover Mites	Grasshoppers	Sod Webworms
Fire Ants	Crickets	Hornets	Sowbugs (Pillbugs)
Armyworms	Cutworms	Japanese Beetles [†]	Spider Mites
Bees	Dichondra Flea Beetles	Midges	Spiders
Beetles	Earwigs	Millipedes	(Including Black Widow Spiders)
Biting Flies	Elm Leaf Beetles	Mosquitoes	Springtails
Boxelder Bugs	Firebrats	Moths	Ticks (Including Brown Dog Ticks)
Centipedes	Fleas	Roaches (Including Cockroaches)	Wasps
Chiggers	Flies	Scorpions	

[†] Not for use in California.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack and crevice treatment. During application, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters.

Additional Application Restrictions for Residential Outdoor Surface and Space Sprays:

All outdoor applications, if permitted elsewhere on this label, must be limited to spot or crack-and-crevice treatments only, except or the following permitted uses, if allowed elsewhere on this label:

1. Application to soil or vegetation, as listed on this label, around structures;
2. Applications to lawns, turf, and other vegetation, as listed on this label;
3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
4. Applications to the underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch, or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters, in order to control occasional invaders or aggregating pests.

Application

Use a 0.02 to 0.06% dilution to spray outside surfaces of buildings. Use a spray volume of up to 10 gallons of dilution per 1,000 square feet. Use higher application volumes if vegetation or landscape materials are dense.

Mixing Directions: For 0.02% suspension, mix 0.33 fluid oz. of **Talak 7.9% F** per gallon of water. For 0.06% suspension, mix 1 fluid oz. **Talak 7.9% F** per gallon of water (1 fluid oz. = 2 tablespoons). Do not use household utensils to measure **Talak 7.9% F**. Use the higher rates for heavy pest infestation, quicker knockdown or longer residual control. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application only if there are signs of renewed insect activity. Repeat application limited to once per seven days.

Perimeter Treatment: Treat a band of soil and vegetation 6 to 10 feet wide around and next to the structure and the foundation of the structure to a height of 2 to 3 feet. Use 0.33 to 1.0 fluid oz. of **Talak 7.9% F** per 1,000 square feet in enough water to provide sufficient coverage (refer to **Perimeter Application Dilution Chart**).

For Ant and Fire Ant Mounds use Talak 7.9% F 0.06% Dilution as Drench Method: Use 1 to 2 gallons of dilution for each mound area. Sprinkle the mound until wet and apply to a 4 foot diameter circle around the mound. For mounds larger than 12", use a higher volume. Applications should be made in cool weather, such as in early morning or late evening hours, not in the heat of the day.

Mosquito Control: To control mosquitoes around buildings, landscapes, and lawns, dilute 0.33 to 1.0 fl. oz. of **Talak 7.9% F** per gallon of water and apply at the rate of one gallon of dilution per 1,000 square feet as a general spray. **Talak 7.9% F** may be diluted at lower concentrations and applied at higher volumes to ensure the proper amount of product per area (refer to the **Ornamental or Perimeter Application Dilution Charts**).

Calculating Dilution Rates: The following steps should be taken to determine the appropriate dilution of **Talak 7.9% F** that is required to control specific pests:

- 1) Select an application rate in of fluid oz. of **Talak 7.9% F**.
- 2) Determine your application volume and amount of spray mix you want to prepare in the **Dilution Chart**.
- 3) Use the **Dilution Chart** to determine the appropriate volume of **Talak 7.9% F** that must be mixed in your desired volume of water.

Perimeter Application Dilution Chart

Application Volume	Application Rate	Fluid Ounces* of Talak 7.9% F Diluted to these Volumes of Finished Spray			
Gallons per 1,000 sq. ft.	Fluid Ounces per 1,000 sq. ft.	1 gallon	5 gallons	10 gallons	100 gallons
1.0	0.33	0.33	1.67	3.3	33.3
1.0	0.5	0.5	2.5	5.0	50.0
1.0	0.67	0.67	3.33	6.7	66.7
1.0	0.75	0.75	3.75	7.5	75.0
1.0	1.0	1.0	5.0	10.0	100.0
2.0	0.33	0.17	0.83	1.65	16.5
2.0	0.5	0.25	1.25	2.5	25.0
2.0	0.67	0.33	1.67	3.35	33.5
2.0	0.75	0.38	1.88	3.75	37.5
2.0	1.0	0.5	2.5	5.0	50.0
3.0	0.33	0.11	0.55	1.10	11.0
3.0	0.5	0.17	0.83	1.67	16.7
3.0	0.67	0.22	1.11	2.23	22.3
3.0	0.75	0.25	1.25	2.5	25.0
3.0	1.0	0.33	1.67	3.33	33.3
4.0	0.33	-	0.42	0.83	8.3
4.0	0.5	0.13	0.63	1.25	12.5
4.0	0.67	0.17	0.84	1.67	16.7
4.0	0.75	0.19	0.94	1.88	18.8
4.0	1.0	0.25	1.25	2.5	25.0
5.0	0.33	-	0.33	0.67	6.7
5.0	0.5	0.1	0.5	1.0	10.0
5.0	0.67	0.13	0.67	1.33	13.3
5.0	0.75	0.15	0.75	1.5	15.0
5.0	1.0	0.2	1.0	2.0	20.0

(continued)

Perimeter Application Dilution Chart *(continued)*

Application Volume	Application Rate	Fluid Ounces* of Talak 7.9% F Diluted to these Volumes of Finished Spray			
Gallons per 1,000 sq. ft.	Fluid Ounces per 1,000 sq. ft.	1 gallon	5 gallons	10 gallons	100 gallons
10.0	0.33	-	0.17	0.33	3.3
10.0	0.5	-	0.25	0.5	5.0
10.0	0.67	-	0.33	0.67	6.7
10.0	0.75	-	0.38	0.75	7.5
10.0	1.0	0.1	0.5	1.0	10.0

* To convert fluid ounces to milliliters, multiply by 29.57
1 fluid ounce = 29.57 ml = 2 tablespoons = 6 teaspoons
Do not use household utensils to measure **Talak 7.9% F**.

INDOOR USE

Talak 7.9% F may be used for residual pest control in buildings and structures and on modes of transport. For control of ants, bees, beetles, boxelder bugs, carpet beetles, centipedes, clothes moths, cockroaches, crickets, earwigs, firebrats, flies, gnats, midges, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders, ticks and wasps.

In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before reuse. Exposed food should be covered or removed. Do not permit humans or pets to contact treated surfaces until the spray has dried. During any overhead applications to overhead interior areas of structures, cover surfaces below with plastic sheeting or similar materials.

Application

Apply to areas where pests hide. These areas include baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, cabinets, sinks, furnaces, stoves, the underside of shelves, drawers and similar areas. Treat with a low-pressure spray (25 p.s.i. or less), coarse, crack and crevice, pinstream, spot or with a paint brush. Pay close attention to cracks and crevices. Not for use as a space spray.

Mixing Directions: Prepare a dilution of **Talak 7.9% F** for spray or brush application. See **Mixing Directions** in **Pest Control on Outside Surfaces and Around Buildings** section.

- Fill sprayer with the required amount of water.
- Add **Talak 7.9% F**.
- Close sprayer and shake to ensure proper mixing.
- Prepare only the amount of solution necessary for treatment.

In order to achieve and/or maintain control in times of high pest pressure, retreatment may be needed. Repeat application should only take place if there are signs of renewed insect activity and should not exceed one application per 7 days.

Application Rates: For 0.02% suspension, mix 0.33 fluid oz. of **Talak 7.9% F** per gallon of water. For 0.06% suspension, mix 1 fluid oz. **Talak 7.9% F** per gallon of water (1 fluid oz. = 2 tablespoons). Do not use household utensils to measure **Talak 7.9% F**. Use the higher rates for heavy pest infestation, quicker knockdown or longer residual control.

Pest	Application Rate	Application Instructions
Ants Bedbugs Bees Beetles Boxelder Bugs Carpet Beetles Centipedes Clothes Moths Cockroaches Crickets Earwigs Firebrats	Flies Gnats Midges Millipedes Pillbugs Scorpions Silverfish Sowbugs Spiders Ticks Wasps	0.33 - 1.0 fl. oz. per gallon of water (0.02% - 0.06% suspension)
Ants: Apply to ant trails, around doors and windows and other places that ant frequent. Bedbugs: To help control of Bedbugs, apply thoroughly to crack and crevices where bedbugs frequent. This includes bed frames, box springs, inside empty dressers and clothes closets, carpet edges, wall moldings (high and low), and wallpaper edges. Do not use on bed linens, pillows, mattresses, or clothes. Before application, remove all clothes and other articles from dressers or clothes closets. Allow all treated areas to dry before use. Not recommended for use as sole protection against bedbugs. If evidence of bedbugs is found in/on mattresses, use a product approved for this use. Bees and Wasps: Apply to nest in late evening when pests are at rest. Spray nests and surrounding areas thoroughly. Spray nests, entrances to nests and surrounding areas thoroughly. Contact as many insects as possible. Retreat if signs of renewed activity exist. Boxelder Bugs, Centipedes, Earwigs, Beetles, Millipedes, Pillbugs, and Sowbugs: Treat around doors, windows, baseboards, storage areas and other locations where pests may be found. Cockroaches, Crickets, Firebrats, Scorpions, Silverfish, Spiders, and Ticks: Use a coarse, low pressure spray to areas where pests hide. These areas include baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, cabinets, sinks, furnaces, and stoves, the underside of shelves, drawers, and similar areas. Pay close attention to cracks and crevices.		

Food/Feed Handling Establishments

Talak 7.9% F, when used as a general spot, surface, or crack and crevice treatment, may be applied in both food/feed and nonfood areas of food/feed handling establishments.

Food/feed handling establishments are any place other than private residences where exposed food/feed is held, processed, prepared or served, including areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, enclosed processing systems (mills, dairies, edible oils, syrups) of food and edible waste storage. Serving areas where food is exposed and the facility is in operation are also considered food areas.

Non-food areas in which applications are allowed include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets and storage (after canning or bottling).

Some of the use sites that are allowed include: Aircraft (do not use in aircraft cabins), apartment buildings, bakeries, bottling facilities, breweries, buses, cafeterias, candy plants, canneries, dairy product processing plants, food manufacturing plants, food processing plants, food service establishments, granaries, grain mills, hospitals, hotels, industrial buildings, laboratories, meat/poultry/egg processing plants, mobile/motor homes, nursing homes, offices, railcars, restaurants, schools, ships, trailers, trucks, vessels, warehouses and wineries.

Surface Application: Do not use this application method in food/feed handling establishments when the facility is in operation or foods/feeds are exposed. During treatment, remove or cover all food processing and/or handling equipment and do not apply directly to food products. All equipment, benches, shelving and other surfaces in food processing plants, bakeries, cafeterias, and other facilities, which food will contact must be washed after treatment. Clean food handling equipment or processing equipment and rinse completely with fresh, clean water.

Spot, Crack and Crevice Application: These types of treatments can be done when the facility is operating, but food should be covered or removed from the treatment area. Do not apply directly to food.

Foam Applications: Converting **Talak 7.9% F** to foam will allow it to be used to treat structural voids. To produce a 0.02% to 0.06% foam concentration, dilute 0.33 to 1.0 fl. oz. of **Talak 7.9% F** per gallon of water and add the manufacturer's recommended amount of foaming agent. Before application, make sure that the foaming agent is compatible with **Talak 7.9% F**.

TERMITE CONTROL (ABOVE GROUND ONLY)

The treatment methods that are expressed below are intended to kill termite workers or winged reproductives present at the time of application. These methods should supplement, not substitute for, mechanical alteration, soil treatment or foundation treatment.

Controlling winged reproductive termites and exposed workers in localized areas can be accomplished by diluting 1.0 fluid ounce of **Talak 7.9% F** per gallon of water and applying the dilution at the rate of one gallon per 1,000 square feet to attics, crawl spaces, unfinished basements and other void areas as a coarse fan spray. Both swarming termites and the areas where they gather should be treated.

Controlling above-ground termites in localized areas of infested wood may be accomplished by diluting 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and applying as a liquid or foam to voids and galleries in wood that is damaged, in addition to spaces between wooden structural members and between the sill plate and foundation where the wood is at risk to attack. Drilling and then injecting the foam or dilution into damaged wood or wall voids with an appropriate directional injector will help reach those areas that are not easy to access. After treatment is completed, securely plug the holes that are in regularly occupied areas in the construction elements.

Controlling termite carton nests in building voids can be accomplished by diluting 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and apply as a liquid or foam using a pointed injection tool. To obtain control, various depths of injection and numerous

injection points may be needed. After treatment is complete and when feasible, remove the carton nest material from the building void.

ANT CONTROL

Nuisance Ants Indoors: Apply to ant nests for best results. Apply a dilution of 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per gallon of water at the rate of one gallon of dilution per 1,000 square feet to places where ants have been seen or are believed to forage as a general surface, crack and crevice or spot treatment. Some of these areas include baseboards, in and behind cabinets, under and behind dishwashers, furnaces, refrigerators, sinks and stoves, around pipes, cracks and crevices and in corners. Pay close attention when treating entry points into the home or around doors and windows. When combining liquid **Talak 7.9% F** treatments with bait treatments use **Talak 7.9% F** as instructed above and apply baits in those areas where **Talak 7.9% F** has not been applied. Do not apply as a broadcast or general surface treatment in residential areas.

Nuisance Ants Outdoors: Apply to ant nests for best results. Treat ant trails, around doors and windows, and other places where ants have been seen or are likely to forage. Treat using a low or high volume perimeter treatment depending on the density of vegetation and landscape materials as described in the **Pest Control on Outside Surfaces and Around Buildings** section of this label. When treating concrete surfaces, more frequent treatments, higher dilutions and/or application volumes may be needed for ant control. The following procedure will normally allow optimal control:

1. Non-porous surfaces should be treated with low volume applications using 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and applying this dilution at the rate of one gallon per 1,000 square feet.
2. Vegetation and porous surfaces should be treated with high volume applications using dilutions that are calculated to deliver 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per 1,000 square feet (refer to the **Ornamental and Perimeter Application Dilution Charts**).
3. Dilute 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and apply at a rate of up to 10 gallons of dilution per 1,000 square feet for maximum residual control.

Carpenter Ants Indoors: Treat areas where carpenter ants are seen or are predicted to forage, such as baseboards, in and behind cabinets, under and behind dishwashers, furnaces, refrigerators, sinks, and stoves, around pipes, cracks and crevices, and in corners by diluting 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and applying at the rate of one gallon of dilution per 1,000 square feet as a general surface, crack and crevice or spot and/or foam treatment. Pay close attention to treating entry points into the home or premises such as around doors and windows. Spray or foam into cracks into crevices or drill holes and spray, mist or foam into voids and galleries where carpenter ants or their nests are present. When combining liquid **Talak 7.9% F** treatments with bait treatments use **Talak 7.9% F** as instructed above and apply baits in those areas where **Talak 7.9% F** has not been applied. Do not apply as a broadcast or general surface treatment in residential areas.

Carpenter Ants Outdoors: Treat carpenter ant nests for best results. Treat areas where carpenter ants are seen or are believed to forage, such as ant trails, around doors and windows. As stated in **Pest Control on Outside Surfaces and Around Buildings** section, apply using a low or high volume perimeter treatment of this label. When treating concrete surfaces, more frequent treatments, higher dilution and/or application volumes may be needed for carpenter ant control. Following the procedure below will normally allow optimal control:

1. Treat non-porous surfaces with low volume applications using 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and applying this dilution at the rate of one gallon per 1,000 square feet.
2. **Talak 7.9% F** may be used as a treatment for the trunks of trees that have carpenter ant trails, or where carpenter ants are foraging. Use 0.5 to 1.0 fl. oz. of **Talak 7.9% F**

per gallon of water and apply this dilution to wet the bark from the base of the tree to as high as possible on the trunk.

3. Vegetation and porous surfaces should be treated with high volume applications using dilutions that are calculated to deliver 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per 1,000 square feet (refer to the **Ornamental and Perimeter Application Dilution Charts**).

4. Dilute 0.5 to 1.0 fluid oz. of **Talak 7.9% F** per gallon of water and apply at a rate of up to 10 gallons of dilution per 1,000 to obtain maximum residual control.

To control carpenter ants inside trees, utility poles, fencing or deck materials and similar structural members, drill to find the inside infested cavity and inject or foam a 0.06% dilution (1.0 fl. oz. of **Talak 7.9% F** per gallon of water) into the cavity with adequate volume and a proper treatment tool with a splash-back guard.

Where there are ants tunneling below the surface, dilute 0.5 to 1.0 fluid ounces of **Talak 7.9% F** per gallon of water and applying as a drench or inject the dilution or foam at intervals of 8 to 12 inches. A uniform vertical barrier should be established where there are ants tunneling below surfaces such as at the edges of walls, driveways or other hard surfaces.

Apply a 0.06% dilution to stored lumber or wood piles using a sprinkling can or a hose-end sprayer to deliver a coarse drenching spray. This wood may be used for lumber or burned after 30 days. Do not use this method of application in structures.

The soil under the area where firewood will be stacked may be treated with a dilution of 1.0 fluid oz. of **Talak 7.9% F** per gallon of water to protect the firewood from carpenter ants (and termites). Apply at the rate of one gallon of dilution per 8 square feet. DO NOT treat firewood with this product.

IMPREGNATION AND APPLICATION OF TALAK 7.9% F ON DRY BULK LAWN FERTILIZERS

Talak 7.9% F may be impregnated on dry bulk fertilizers. When applied as directed, **Talak 7.9% F**/dry bulk fertilizer mixtures provide insect control equal to that provided by the same rates of **Talak 7.9% F** applied in water.

Impregnation: Apply using a minimum 2.3 pounds of dry bulk fertilizer per 1,000 square ft. with the recommended amount of **Talak 7.9% F** per 1,000 square ft. Use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform, fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with **Talak 7.9% F** provides a satisfactory dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, flowable mixture. Microcel E (Johns-Manville Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. DO NOT impregnate **Talak 7.9% F** onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the insecticide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with **Talak 7.9% F**.

The amount of **Talak 7.9% F** actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with **Talak 7.9% F** should be applied immediately, not stored.

All individual Federal and State regulations relating to bulk dry fertilizer blending, registration, labeling, and application of the mixtures are the responsibility of the individual and/or company selling the fertilizer and **Talak 7.9% F** mixture.

Fertilizer for this use should be Turf fertilizers recommended for specific regions.

USE IN LIVESTOCK/POULTRY HOUSING STRUCTURES AND PET KENNELS

For control of pests including biting flies, filth-breeding flies, fleas, litter beetles, hide beetles, bed bugs, mites, and ticks.

Application may be made as a general surface spray (including directed spray) and/or as a crack and crevice treatment. For best results, make interior and exterior applications at or around the same time. In addition to applications of **Talak 7.9% F**, ensure that normal cleaning practices are followed.

Occupied Areas

Indoors, apply only to indoor cracks and crevices. For exteriors, apply to walls and foundation perimeters to help prevent interior infestations of pests. Use **Talak 7.9% F** at a rate equivalent to 0.33 to 1 fl. oz. per 1,000 sq. feet.

Unoccupied Areas

Apply to areas where crawling or flying pests may be present, such as floors, vertical surfaces, and overhead surfaces, paying special attention to areas such as stanchions, pipes, windows, and doors. Cover feeders, waterers, and feed carts before application, to avoid contamination. Do not apply to milk rooms. Make exterior applications to walls and foundation perimeters to help prevent interior infestations of pests. Use **Talak 7.9% F** at a rate equivalent to 0.33 to 1 fl. oz. per 1,000 sq. feet.

Bed Bugs, Mites and Ticks - Treat cracks and crevices, walls, posts, nest boxes, and mobile side curtains. Do not apply this product directly to animals.

Adult Flies - Make applications to areas where flies will rest, such as the ceiling, rafters, and trusses; also treat windows, walls (interior and exterior), supports, fences, and vegetation. **Talak 7.9% F** may be applied to manure in situations where fly larvae are abundant and the area cannot be cleaned.

Poultry Houses - Make applications to the floor (where birds are grown on litter), walls, posts, and cage framing (where birds are grown in cages); apply also into cracks and crevices around insulation. Reapply after each growout or sanitization procedure, but not more often than every 8 weeks. For improved indoor control, apply to the outside of building foundations to keep adult beetles from moving indoors. Apply in a uniform band 2 to 3 feet up the foundation, and 6 to 10 feet out from the structure. A routine, year-round treatment program will prevent pests from reaching problem levels.

Where Birds are Grown on Litter - Apply **Talak 7.9% F** to litter after birds are removed and during tilling at a rate equivalent to 0.33 to 1 fl. oz. per 1,000 sq. feet. If litter is removed and replaced with fresh litter, make an application to bare soil or concrete at a rate equivalent to 0.33 to 1 fl. oz. per 1,000 sq. feet, and treat the new litter once it is spread. Spray inside walls, posts, and exterior perimeter. Reapply between each flock.

Broiler-breeder Houses - To control beetles; apply as directed above for litter and soil/floor treatment.

Caged-layer Houses - For control of beetles, do not treat accumulated manure because it may disrupt natural enemies that control fly breeding. Treat the perimeter of the manure at a rate equivalent to 0.33 to 1 fl. oz. per 1,000 sq. feet. Also spray pit walls, posts, and the exterior of the structure. Reapply between each flock.

Before applying disinfectants, ensure that the **Talak 7.9% F** treatment is dry.

DO NOT apply **Talak 7.9% F** as a general surface spray when animals are present in the facility. Allow applications to dry before restocking the facility. Crack and crevice treatment may be made when animals are present.

DO NOT apply **Talak 7.9% F** to any animal feed, water, or watering equipment.

DO NOT contaminate any animal feed, food, or water in and around livestock, poultry, or pet housing when making applications.

Attention:

- Do not apply a broadcast application to interior surfaces of homes.
- Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm.
- Do not apply in classrooms when in use.
- Do not apply when occupants are present in the immediate area in institutions such as libraries, sports facilities, etc.
- Do not allow people or pets on treated surfaces until spray has dried.
- Let surfaces dry before allowing people and pets to contact surface.
- During any application to overhead areas of structure, cover surface below with plastic sheeting or similar material except for soil surfaces in crawlspaces.
- Do not allow spray to contact food, foodstuffs, food-contacting surfaces or food utensils or water supplies.
- Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product.
- Do not treat areas where food is exposed.
- During indoor surface applications do not allow dripping or runoff to occur.
- **Talak 7.9% F** will not stain, or damage any surface that water alone will not stain or damage.
- Do not apply to pets, crops, or sources of electricity.
- Firewood is not to be treated.
- Use only in well ventilated areas.
- Do not use on edible crops.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use replace lids and close tightly. Do not put concentrate or dilute material into food or drink container.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To Confine Spill: If liquid, dike surrounding area or absorb with sand, cat litter, or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal: Pesticide wastes are toxic. Do not contaminate water, food, or feed by storage or disposal. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. Dispose of excess or waste pesticide by use according to label directions, or contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling: Plastic Container: Non-refillable Containers. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

Containers 5 gallons or less: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once cleaned, offer for recycling if available.

STORAGE AND DISPOSAL (continued)

Containers larger than 5 gallons: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Once cleaned, offer for recycling or reconditioning if appropriate.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. All such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Talak is a trademark of Atticus, LLC.

Actisol® is a registered trademark of Environmental Delivery Systems, Inc.

Delavan® is a registered trademark of Delavan Spray, LLC.

Patriot Injector® is a registered trademark of Unifex USA, LLC.

Raindrop® is a registered trademark of GP Companies, Inc.

TeeJet® is a registered trademark of Spraying Systems Company.

F20130613a



NixloTM IGR Concentrate

Contains pyriproxyfen, the active ingredient used in NyGuard® IGR Concentrate.

Insect Growth Regulator breaks the cycle of insect reproduction to control Fleas, Stored Product Pests, Cockroaches, Mosquitoes, Flies, Lone Star Ticks and other listed insect pests.

Effective long-term control against fleas for 7 months and cockroaches for 6 months

- Labeled for use in food areas of food handling establishments
- Works to break the life cycle of listed insects
- Effective indoors and outdoors
- Reduces egg production in house flies and prevents development of larvae (maggots) into adults

ACTIVE INGREDIENT:	(% by weight)
2-[1-Methyl-2-(4-phenoxyphenoxy) ethoxy] pyridine [Pyriproxyfen]	10.0%
OTHER INGREDIENTS:	90.0%
TOTAL	100.0%

EPA Reg. No.: 91234-336

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
If in eyes:	▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.
If on skin or clothing:	▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
NOTE TO PHYSICIAN	
Vomiting may cause aspiration pneumonia.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

NixloTM IGR Concentrate is not manufactured, or distributed by MGK Company, seller of NyGuard® IGR Concentrate.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear protective eyewear such as goggles, face shield, or safety glasses. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and waterproof chemical-resistant (Barrier Laminate or Viton $\geq$ 14 mils) gloves. Do not induce vomiting because of aspiration pneumonia hazard. Obtain medical attention if irritation persists. Avoid contamination of food or feedstuffs.

Cover or remove all food and food processing surfaces. Thoroughly wash all food processing surfaces before reuse. After spraying in bakeries, meat packing plants, food processing plants, etc., all benches, shelving, equipment, etc. Where exposed food will be handled must be washed with an effective compound followed by a potable water rinse to remove all traces of contamination. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection program.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users Should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT use this product in or on electrical equipment due to the possibility of shock hazard. Do not mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- DO NOT use this product in drinking water systems.
- DO NOT use this product in or on electrical equipment due to shock hazard.

READ AND FOLLOW ALL LABEL DIRECTIONS WHEN APPLYING THIS PRODUCT BEFORE USE.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Nixlo IGR Concentrate is a non-repellent Juvenile Hormone (JH) Mimic. It disrupts the normal development of the immature life stages of listed insects, with either complete or incomplete metamorphosis, preventing successful development of the adult stage. Depending upon the insect species and life stage at the time of exposure, the observed results can include: decreased egg viability, sterile adult insects incapable of reproduction, suspension of insect in immature life stages, reduced fitness of insect, behavioral changes, reduced egg production, morphological defects – deformed insects and death. When used in an Integrated Pest Management (IPM) program, **Nixlo IGR Concentrate** breaks the reproductive cycle of listed insects, and aids in the prevention of insect resistance.

RESISTANCE MANAGEMENT

For resistance management, **Nixlo IGR Concentrate** contains a Group 7C insecticide. Any insect population may contain individuals naturally resistant to **Nixlo IGR Concentrate** and other Group 7C insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly at the same use sites. Appropriate resistance management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of **Nixlo IGR Concentrate** or other Group 7C insecticides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, you may contact your local Atticus, LLC representative or by calling 984-465-4800.

SPRAY DRIFT

Boom-less Ground Applications:

- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.



SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Spray Nozzle- Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

SHIELDED SPRAYERS

Shielding the individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boom-less Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications:

- Take precautions to minimize spray drift.

STAINING

Nixlo IGR Concentrate is generally non-staining. If staining is a concern, test final spray solution in an inconspicuous area prior to use, as some fibers or surfaces (such as vinyl siding) may be adversely affected by an application of this, or any liquid product.

Prior to using in commercial or residential food areas, please refer to the **Food Handling Establishment Applications** section of the label. The preparations prior to treatment and the cleanup after application described are also to be followed for residential use.

In hospitals, nursing homes or in any rooms occupied by the elderly or infirm, remove patients before treating rooms. If application is done as a surface spray, or crack and crevice application, patients may re-enter once the spray has dried. If application is a ULV fog or space spray, patients may return to the room after it has been thoroughly ventilated.

In institutions (including libraries, sports facilities, etc.) applications may be made while the facility is in use, provided treatment is not made in the immediate area where people are present.



USE SITES

Nixlo IGR Concentrate is effective when used in both indoor and outdoor areas such as:

Agricultural Buildings,	Chests,	Food Handling Areas,	Mass Merchandising Stores,	Septic Tanks,
Aircraft (cargo areas only),	Child Care Centers,	Food Packing Plants,	Mattresses,	Ship Cargo Holds,
Animal Housing Facilities,	Closed Porches,	Food Processing Plants,	Mausoleums,	Ships,
Animal Quarters,	Closets,	Food Storage Areas,	Meat & Produce Canneries,	Shutters,
Animal Research facilities,	Clothes Storage,	Food Warehouses,	Meat Processing Plants,	Side Yards,
Apartment Buildings,	Coffee Houses,	Foundations,	Mess Halls,	Soil,
Apartments,	Colleges,	Funeral Parlors,	Mobile Canteens,	Solariums,
Appliances,	Commercial Buildings,	Furniture,	Mobile Homes,	Spas,
Assisted Living Facilities,	Compost,	Garages,	Morgues,	Sports Stadiums,
Atriums,	Composting Sites,	Garbage Areas,	Mortuaries,	Stockyards,
Attics,	Condominiums,	Garbage Cans,	Motels,	Storage Areas,
Automobiles,	Convenience Stores,	Gas Stations,	Mulch,	Stored Food Areas,
Backyards,	Crawl Spaces,	Gazebos,	Nursing Homes,	Stored Food Warehouses,
Bakeries,	Crematoriums,	Grain Storage Facilities,	Office Buildings,	Stores,
Barns,	Cupboards,	Grass, Grocery Stores,	Offices,	Sun Porches,
Basements,	Dairies,	Headboards,	Other Public Buildings,	Sunrooms,
Bathrooms,	Day Care Centers,	Health Clubs,	Outdoor Eating Establishments,	Supermarkets,
Bed Frames,	Decks,	Homes,	Pantries,	Swine Production Facilities,
Bedrooms,	Delicatessens,	Hospices,	Parks,	Taverns,
Boats,	Dens,	Hospitals,	Patios,	Theaters,
Boiler Rooms,	Detention Centers,	Hotels,	Pet Areas,	Tobacco Plants and Warehouses,
Bookshelves,	Dining Halls,	Indoor Eating Establishments,	Pet Bedding,	Town Houses,
Bottling Facilities,	Dog Houses,	Industrial Buildings,	Pet Carriers,	Trailers,
Box Springs,	Dog Kennels,	Industrial Plants,	Pet Grooming Parlors,	Trains,
Boxcars,	Dog Runs,	Institutional Buildings,	Pet Quarters,	Transportation Vehicles,
Breweries,	Dormitories,	Institutional Dining Areas,	Pet Sleeping Areas,	Tree Houses,
Building Exteriors,	Drains,	Institutions,	Pet Stores,	Trucks,
Building Foundations,	Draperies,	Jails,	Playhouses,	Universities,
Building Perimeters,	Dressers,	Kennels,	Playrooms,	Upholstered Furniture,
Buses,	Drive-Ins,	Kitchens,	Porches,	Utility Room Indoor,
Cabinets,	Drugstores,	Laboratories,	Poultry Houses,	Utility Room Outdoor,
Cabins,	Egg Production Facilities,	Lanais,	Prisons,	Vending Machines,
Cafeterias,	Equipment,	Landscape Plants,	Public Buildings,	Verandas,
Calf Hutches,	Equipment Voids,	Landscaping,	Railroad Cars,	Veterinaries,
Campers,	Fabric Storage Areas,	Lawns,	Recreation Vehicles,	Warehouses,
Campsites,	Factories,	Libraries,	Recreational Areas Indoors,	Washrooms,
Candy Plants,	Farm Buildings,	Livestock Housing,	Recreational Areas Outdoors,	Yards,
Cargo Holds,	Farms,	Livestock Production Facilities,	Residential Homes,	Zoos
Carpets,	Farmyards,	Living Rooms,	Restaurants,	
Casinos,	Feedlots,	Locker Rooms,	Rooms,	
Cemeteries,	Floors,	Manufacturing Plants,	Rugs,	
Cereal Processing Facilities,	Flour Mills,	Marine Vessels,	Schools,	

Pests in the Acari subclass, Anoplura superfamily, Blattodea, Coleoptera, Dermaptera, Diptera, Heteroptera suborder, Hymenoptera, Lepidoptera, Orthoptera, Psocoptera, Siphonaptera, as well as certain other orders are susceptible in varying degrees to **Nixlo IGR Concentrate's** active ingredient. Specific pests that **Nixlo IGR Concentrate** has shown to significantly disrupt the life cycle are:

Almond Moths,	Cheese Skippers,	Filter Flies,	Lesser Grain Borers,	Sawtoothed Grain Beetles,
American Cockroaches,	Cigarette Beetles,	Fleas,	Litter Beetles,	Sewage Flies,
Angoumois Grain Moths,	Cinereous Cockroaches,	Flies,	Lone Star Ticks,	Silverfish,
Ants,	Clothes Moths,	Florida Stink roaches,	Madeira Cockroaches,	Small Dung Flies,
Asian Cockroaches,	Cockroaches,	Fruit Flies,	Mealworms,	Smoky Brown Cockroaches,
Asian Lady Beetles,	Confused Flour Beetles,	Fungus Gnats,	Merchant Grain beetles,	Stable Flies,
Australian Cockroaches,	Copra Beetles,	German Cockroaches,	Midges,	Surinam Cockroaches,
Beetles,	Crickets,	Granary Weevils,	Mosquitoes,	Tobacco Moths,
Black Carpet Beetles,	Darkling Beetles,	Ground Beetles,	Moths,	Turkestan Cockroaches,
Booklice,	Dermeid Beetles,	Hide Beetles,	Pennsylvania Wood Cockroaches,	Vinegar Flies,
Bottle Flies,	Drain Flies,	Horn Flies,	Phorid Flies,	Warehouse Beetles
Boxelder Bugs,	Drugstore Beetles,	Horntails,	Psocids,	
Brown Cockroaches,	Elm Leaf Beetles,	House Flies,	Red Flour Beetles,	
Brownbanded Cockroaches,	Face Flies,	Indian Meal Moths,	Rice Moths,	
Cat Fleas,	Field Cockroaches,	Khapra Beetles,	Rice Weevils,	

MIXING INSTRUCTIONS

Mix concentrate with water and apply with equipment capable of delivering coarse wetting surface spray, fogging (space spray/mist), spot, crack and crevice spray applications, professional foam applicator or with a paintbrush, roller or similar device. Partially fill the sprayer with water, add required amount of concentrate in sprayer, agitate and fill to final volume. Agitate before each spray.

Tank Mixing

Nixlo IGR Concentrate may be tank mixed with other registered pest control products, unless either product label prohibits tank mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Make applications in accordance with those label instructions that are consistent for each product. To prepare the tank mix, add the tank mix partner first and make sure it is well mixed and in suspension, then add the correct amount of **Nixlo IGR Concentrate** according to the label instructions below.

Tank Mix Restrictions:

- Do not tank mix with Vapona or Malathion. These products are antagonistic to **Nixlo IGR Concentrate's** active ingredient.

APPLICATION INSTRUCTIONS

Use **Nixlo IGR Concentrate** both indoors and outdoors as a general surface spray, ULV/fog or space spray application, spot treatment or crack and crevice application. Application may be made using power sprayers, hand pump sprayers, void injectors, in carpet shampoo, using a paintbrush, roller, professional foam applicator, or similar device.

SURFACE APPLICATIONS

Apply diluted spray as a spot treatment, broadcast surface application and crack and crevice spray. Thoroughly treat areas where insects travel, breed, rest, hide or harbor. Special attention should be paid to applying spray solution to harborage and hiding places where conditions support insect population growth and reproduction.

ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION

Nixlo IGR Concentrate may be applied as a fog or an ultra-low volume (ULV) application with equipment such as foggers, ULV equipment, mechanical misting sprayers or aerosol generators. To prepare the fogging solution, follow the equipment manufacturer's solution preparation instructions. For fogging applications, **Nixlo IGR Concentrate** may be diluted with water or an approved petroleum distillate. Apply at a rate of 1 ml of **Nixlo IGR Concentrate** per 1,000 cu. ft. of space to be treated. Remove all persons from the area to be treated. Close area to be treated. Shut off ventilation systems. Apply the product in the air and direct fog or mist toward all areas that may harbor target pests. After treatment, leave the room closed for 30 minutes or longer, as instructed by the labels of other products that may be tank mixed with **Nixlo IGR Concentrate**. Exit the treated area after fogging application is completed. Thoroughly ventilate area before re-entry.

To treat voids in equipment and structures with ULV and fog injection systems (void injectors): Apply to harborage sites in wall voids and other inaccessible spaces in equipment and structures using injection equipment designed for deep void ULV (Ultra Low Volume) applications. Follow the injection equipment manufacturer's recommendations for the proper set up and air pressures. Place the applicator tip at or into the void space or crack and crevice to be treated. Apply in short bursts while keeping the tip in position. Allow the air stream to push the material deep into the space. For applications along baseboards, trim, molding, etc. move applicator tip along the crevice to be treated, applying a constant burst of material. Move the tip at a steady rate of one foot per second. Avoid applying to the point of runoff or drip. Ventilate area thoroughly before re-entry.

FOOD HANDLING ESTABLISHMENT APPLICATIONS

Apply **Nixlo IGR Concentrate** as a general surface spray, space spray, spot and/or crack and crevice treatment in both food and non-food areas of food handling establishments where food and food products are held, processed, prepared and served. It also includes areas where food items are: received, prepared, packaged, and stored, as well as enclosed food processing systems (mills, dairies, etc.). Prior to application, remove or cover all food items in area to be treated.

Restriction: Do not apply directly to food or food-handling surfaces.

In the home, cover all food-processing surfaces and utensils during treatment or thoroughly wash before use. Thoroughly wash food contact surfaces with an effective cleaning compound and rinse with potable water after use of the product.

SPOT OR CRACK AND CREVICE APPLICATION

Since active ingredient is stable and not volatile, spot, and/or crack and crevice treatment may be used while the facility is in operation, provided exposed food is covered or removed from the area being treated prior to application.

SPACE SPRAY AND GENERAL SURFACE APPLICATION

Use space and general surface applications only when the facility is not in operation provided exposed food is covered or removed from the area being treated prior to application.

PANTRY/CABINETS, SMALL FOOD STORAGE AREAS

Apply as a surface spray, fog, crack and crevice spray, or spot treatment to pantry and cupboard shelves or other areas where food is stored. Prior to spraying, remove all food containers and foodstuffs. Replace food containers or foodstuffs after spray has dried.

STORED FOOD WAREHOUSES

Apply **Nixlo IGR Concentrate** as a general surface spray, fog, spot, or crack and crevice treatment in warehouses where packaged food items are stored in such a way that direct contact with the foodstuff is not anticipated.

Integrated Pest Management in Schools

The health of school residents and long-term suppression of pests must be the primary objectives that guide pest control in school settings. To accomplish these objectives, an IPM program must always look for alternatives first and use pesticides only as a last resort. There are many chemical products to choose from that pose less risk to humans and the larger environment and are, at the same time, effective against target pests.

Consult the EPA Document: IPM for Schools: A How-to Manual to learn more. This and other resource materials are available at <https://www.epa.gov/ipm>.

Nixlo IGR Concentrate may be used in schools and day care facilities. Unless it is an emergency situation, perform applications when only maintenance staff are present, and the building is otherwise unoccupied. Apply to classrooms which are not in use. Vacate rooms prior to treatment. If application is a surface spray, spot or crack and crevice application, re-entry is permitted once the spray has dried. If application is a ULV fog or space spray, re-entry is permitted after the treated area has been thoroughly ventilated.

Always consult federal, state and local governing bodies to understand any and all requirements prior to application.



Rate Chart

Amount of Nixlo IGR Concentrate per gallon of water to treat 1,500 sq. ft.	Amount of Nixlo IGR Concentrate per unit area	
	1,000 sq. ft.	1,500 sq. ft.
4 ml per gallon (0.01%)	3.0 ml	4.0 ml
8 ml per gallon (0.02%)	5.5 ml	8.0 ml
12 ml per gallon (0.03%)	8.0 ml	12.0 ml

Alternate mixing instructions for outdoor power spraying		
18 ml per 50 gallons of water		0.001%
38 ml per 50 gallons of water		0.002%
56 ml per 50 gallons of water		0.003%

FOR INDOOR AND OUTDOOR CONTROL OF LISTED FLYING INSECTS SUCH AS MOTHS, MOSQUITOES, FLIES AND SMALL NUISANCE FLIES

Nixlo IGR Concentrate decreases the number of eggs laid by treated adults and reduces hatch rates. Larvae show physiological and functional defects leading to the inability to develop into normal adults. When used alone, treatment will result in a reduction in adult fly populations after 3 to 4 weeks, when previously hatched adult flies have died naturally. Apply spray dilution to known and suspected breeding sites (such as moist areas with decaying organic matter), as well as areas where these insects are known to travel, rest, hide or harbor. Make application following cleaning and sanitation practices. Tank mix **Nixlo IGR Concentrate** with an adulticide to get immediate control of adult stages while inhibiting successful development of immature stages. Monitor insect population and repeat applications at 2 to 4 week intervals, if necessary, depending upon the severity of infestation. If application is being made in food areas, read and follow the “**FOOD HANDLING ESTABLISHMENT APPLICATIONS**” instructions in this label.

Surface Application (including Broadcast, Spot and Crack and Crevice):

Apply **Nixlo IGR Concentrate** in a surface application at the rate of 4 ml in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 sq. ft. Use up to 12 ml per 1,500 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Apply solution as a coarse, wetting spray to breeding and harborage sites. For drains, apply dilution into and around the drain and catch basin area. When possible, treat vertical surfaces surrounding breeding sites, where small flies tend to alight.

Space Spray, Void Injection, ULV and Fog Application:

Read and follow the “**ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION**” in the Application Instructions section of this label. To achieve uniform coverage, apply **Nixlo IGR Concentrate** as a ULV or fogging treatment, alone or tank mixed with an adulticide. Apply at the rate of 1 ml of concentrate per 1,000 cu. ft. of space to be treated. Direct application to areas where insects are known to travel, rest, hide or harbor.

Foam Application:

For the control of Drain/ Fruit Flies, combine **Nixlo IGR Concentrate** at the rate of 4 to 8 ml concentrate with one gallon of diluent. To convert this diluent to a foam, add foaming agent at the manufacturer's recommended rate. Using a professional foam applicator, apply finished foam to areas of fly breeding such as drains, traps of sinks, bathtubs and in floor drains. For best results, foam should be applied thoroughly, ensuring all void areas and fly breeding material is treated. Follow the requirements of the foam equipment manufacturer.

Outdoor Application:

Apply as a surface application (including Broadcast, Spot and Crack and Crevice) at the rate of 4 ml in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,000 sq. ft. Use up to 12 ml per 1,000 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Alternate mixing directions for power spraying: use 18 to 56 ml of concentrate per 50 gallons of water.

Apply to surfaces where insects breed, travel, rest, hide or harbor. Treat known and suspected breeding sites on, under and around exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, stagnant water, moist/shaded areas, fence lines, and other areas where target insects are active or may be hiding. Since **Nixlo IGR Concentrate** is photo-stable, outdoor applications on soil surfaces will provide residual IGR activity, depending upon soil type and weather conditions.

FOR INDOOR AND OUTDOOR CONTROL OF LISTED CRAWLING INSECTS SUCH AS FLEAS, LONE STAR TICKS, BEETLES AND OTHER LISTED INSECTS

Nixlo IGR Concentrate breaks the reproductive cycle by acting on the immature life stages of fleas, Lone Star ticks, beetles and other crawling insects, preventing the successful development of adult insects. Flea eggs and flea larvae that contact treated areas will not develop into adult fleas. **Nixlo IGR Concentrate** reduces egg production and hatch and inhibits the laying of eggs by engorged Lone Star tick nymphs. **Nixlo IGR Concentrate** is a valuable component of an integrated program to control fleas, beetles, lone star ticks and other crawling insects. If application is being made in food areas, read and follow the “**FOOD HANDLING ESTABLISHMENT APPLICATIONS**” instructions in this label.

PREVENTIVE FLEA TREATMENT

Nixlo IGR Concentrate can be used as a preventive treatment prior to the flea season. Application of this product in areas where pets and other animals are known to frequent and where previous infestations have been known to occur will reduce the emergence of adult fleas and also control immature stages of fleas. Indoor treatments of **Nixlo IGR Concentrate** provide up to 7 months of residual flea control, so this early usage could reduce the need for conventional insecticides when used alone as directed. Apply as a surface application at the rate of 4 to 8 ml in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 sq. ft.

TREATING ESTABLISHED INFESTATIONS

Surface Application (including Broadcast, Spot and Crack and Crevice):

Apply as a surface application at the rate of 4 ml in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 sq. ft. Use up to 12 ml per 1,500 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Treat all areas that may harbor target insects, such as carpets, furniture, pet sleeping areas and throw rugs. Treat under cushions of upholstered furniture. Use with an adulticide to get immediate control of adult insects. Keep children or pets away from contact with treated surfaces until spray has thoroughly dried.

In Conjunction with Carpet Shampoo:

Nixlo IGR Concentrate may be combined with a liquid, non-heat extracting, carpet shampoo application. Mix 4 ml of **Nixlo IGR Concentrate** per gallon (3.8 L) of diluted shampoo to be applied to carpet. Mix well and shampoo carpet as per instructions on the shampoo product label. Allow carpet to completely dry and vacuum thoroughly.

In Kennels and Doghouses:

Apply as a surface application (including Broadcast, Spot and Crack and Crevice) at the rate of 4 ml in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 sq. ft. Use up to 12 ml per 1,500 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Remove animals before spraying. Apply in and around buildings, resting areas, walls, floor, animal bedding and run areas. Return animals to treated areas when spray has thoroughly dried.

Space Spray, Void Injection, ULV and Fog Application:

Read and follow the “**ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION**” in the Application Instructions section of this label. To achieve uniform coverage, apply **Nixlo IGR Concentrate** as a ULV or fogging treatment, alone or tank mixed with an adulticide. Apply at the rate of 1 ml of concentrate per 1,000 cu. ft. of space to be treated. Direct application to areas where target insects are known to travel, rest, hide or harbor.

Outdoor Application:

Apply as a surface application (including Broadcast, Spot and Crack and Crevice) at the rate of 4 ml in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,000 sq. ft. Use up to 12 ml per 1,000 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Alternate mixing directions for power spraying: use 18 to 56 ml of concentrate per 50 gallons of water. Remove animals before spraying. Apply to listed outdoor areas where target insects are suspected or known to breed, travel, rest, hide or harbor.

To help prevent insect infestation of buildings:

Treat building exteriors, foundations and soil or other substrate adjacent to buildings. Application on soil surfaces will provide 30 to 60 days of IGR activity, depending upon insect type, soil type and weather conditions.

FOR CONTROL OF LISTED COCKROACHES, CRICKETS

Nixlo IGR Concentrate acts on the immature life stages and prevents nymphal cockroaches and crickets from developing into reproductive adults. Shortly after application of this concentrate, cockroaches with twisted wings and other deformities will begin to appear, indicating that **Nixlo IGR Concentrate** has taken effect – those cockroaches are unable to reproduce. May be used in a program with cockroach baits. One treatment works for 6 months against cockroach nymphs and crickets. If application is being made in food areas, read and follow the “**FOOD HANDLING ESTABLISHMENT APPLICATIONS**” instructions in this label.

Surface Application (including Broadcast, Spot and Crack and Crevice):

Apply as a surface application at the rate of 12 ml in one gallon of water to treat 1,500 sq. ft. Apply diluted spray as a spot treatment, surface application and crack and crevice spray. Treat all areas where target insects are known or suspected to travel, rest, hide or harbor. Tank mix with an adulticide to provide immediate control of adult stages. Contact as many insects as possible with this spray. Special attention should be paid to hiding places such as beneath sinks, behind and beneath stoves and refrigerators, cracks and crevices, around garbage cans, cabinets, along the outside of baseboards, door and window sills, door and window frames and floors, around and on drains, pipes, plumbing, behind bookcases, storage and other utility installation areas, and the inside of cabinets and closets hitting insects with spray whenever possible. Repeat application in 6 months, or more frequently if necessary.

Space Spray, Void Injection, ULV and Fog Application:

Read and follow the “**ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION**” in the Application Instructions section of this label. To achieve uniform coverage, apply **Nixlo IGR Concentrate** as a ULV or fogging treatment, alone or tank mixed with an adulticide. Apply at the rate of 1 ml of concentrate per 1,000 cu. ft. of space to be treated. Direct application to areas where insects are known to travel, rest, hide or harbor.

Outdoor Application:

Apply as a surface application (including Broadcast, Spot and Crack and Crevice) at the rate of 12 ml of concentrate in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 sq. ft. For building perimeter applications, apply in a band 2-6 feet adjacent to and around the structure and apply to a height of 2 to 3 feet on the foundation where pests may be active or may find entrance. Apply spray on, around and beneath window frames, doors, garage doors, patios, decks, turf, landscaping, compost areas, garbage areas and other areas where target insects travel, feed, rest, hide or harbor. Alternate mixing directions for outdoor power spraying: use 56 ml of concentrate per 50 gallons of water.

FOR CONTROL OF ANTS

Surface Application Indoors:

Nixlo IGR Concentrate may be used as part of an integrated approach to control ants. Apply broadcast, spot or crack and crevice treatments of **Nixlo IGR Concentrate** at the rate of 12 ml diluted in water per 1,500 sq. ft. to interior areas where ants are actively foraging and traveling. May be used in conjunction with baits and/or residual treatments. If application is being made in food areas, read and follow the “**FOOD HANDLING ESTABLISHMENT APPLICATIONS**” instructions in this label.

Outdoor Application:

Apply as an outdoor surface application (including Broadcast, Spot and Crack and Crevice) at the rate of 12 ml of concentrate in one gallon of water (or sufficient volume of water to get adequate coverage) to treat 1,500 sq. ft. Alternate mixing directions for outdoor power spraying: use 56 ml of concentrate per 50 gallons of water.

For building perimeter applications, apply in a band 2-6 feet adjacent to and around the structure and apply to a height of 2 to 3 feet on the foundation where ants may travel or find entrance. Apply directly to known or suspected ant forage, trails and forage sites. Treat ant mounds or nests directly, with a wetting spray. **Nixlo IGR Concentrate** may be used in a program with baits or in a tank mix combination with adulticide insecticide.



FOR CONTROL OF LISTED STORED PRODUCT PESTS

(Including Indian Meal Moths, Rice Moths, Tobacco Moths, Red Flour Beetles, Confused Flour Beetles, Lesser Grain Borers, Merchant Grain Beetles, Sawtoothed Grain Beetles, Warehouse Beetles, Cigarette Beetles, Dermestid Beetles). If application is being made in food areas, read and follow the “**FOOD HANDLING ESTABLISHMENT APPLICATIONS**” instructions in this label.

Surface Application (including Broadcast, Spot and Crack and Crevice):

Apply as a surface application at the rate of 8 ml in one gallon of water to treat 1,500 sq. ft. Use up to 12 ml per 1,500 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Apply diluted spray as a spot treatment, surface application and crack and crevice spray. Treat all areas where target insects are known or suspected to travel, rest, hide or harbor such as cracks and crevices in walls and floors, under and between pallets, areas at the base of machinery and equipment, under furniture, in mop and equipment closets, behind cabinets, behind and under refrigerators, sinks, stoves, drawers under shelves and racks, around pipes and openings in floors and walls, along baseboards, in wall voids, and other areas where insects seek harborage. Since **Nixlo IGR Concentrate** is stable and not volatile, spot, and/or crack and crevice treatment may be used while a food processing facility is in operation; cover or remove exposed food from the area being treated prior to application. Tank mix with an adulticide to provide immediate control of adult stages. Contact as many insects as possible. Repeat application in 6 months, or more frequently if necessary.

Space Spray, Void Injection, ULV and Fog Application:

Read and follow the “**ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION**” in the Application Instructions section of this label. To achieve uniform coverage, apply **Nixlo IGR Concentrate** as a ULV or fogging treatment, alone or tank mixed with an adulticide. Apply at the rate of 1 ml of concentrate per 1,000 cu. ft. of space to be treated. Direct application to areas where insects are known to travel, rest, hide or harbor.

Use in Livestock and Animal Production

FOR CONTROL OF INSECT PESTS IN AND AROUND DAIRIES, LIVESTOCK AND OTHER ANIMAL HOUSING AND FACILITIES

(Including Kennels, Animal Shelters, Stables and Zoos):

Nixlo IGR Concentrate may be used by pest control professionals and livestock producers in, on and around animal housing and facilities to control listed flying insect pests. **Nixlo IGR Concentrate** mimics naturally occurring insect growth hormones, disrupting the reproductive processes of listed insects, resulting in reduced egg production, reduced egg hatch rates, disruption or inhibition of normal molting processes, deformity, sterile adults and death. **Nixlo IGR Concentrate** effectively breaks the cycle of insect reproduction. It is an important component in an IPM program and is a means to prevent or manage known resistant insect populations. Used as directed, **Nixlo IGR Concentrate*** will not harm beneficial insects such as predatory wasps.

*Pyriproxyfen

Apply as a spot treatment; general surface broadcast treatment, ULV/ fog, spot or crack and crevice application to interior and exterior areas where insect pests collect, travel, rest, hide, harbor or breed. Apply when animals are not in the immediate area being treated. Allow animals to reenter treated area once spray solution has thoroughly dried. **Restriction:** Do not contaminate food, feed, potable water or watering equipment.

RATES TO CONTROL MOSQUITOES AND OTHER FLYING INSECTS

Nixlo IGR Concentrate affects both adult and juvenile insect stages. It decreases the number of eggs laid by treated adults and reduces hatch rates. Larvae (maggots) show physiological and functional defects leading to the inability to develop into normal adults. When used alone treatment will result in a reduction in adult fly populations after 3 to 4 weeks, when previously hatched adult flies have died naturally. For faster results, **Nixlo IGR Concentrate** may be tank mixed with an adulticide. Apply alone or tank mixed with an adulticide at the rate of 4 ml per 1,500 sq. ft. of surface area in sufficient volume of water to thoroughly wet the surface, although not to the point that runoff occurs. Use up to 12 ml of concentrate per 1,500 sq. ft. of surface for insect populations that are difficult to control or in conditions that make it difficult to achieve desired results.

RATES TO CONTROL LISTED CRAWLING INSECTS

Apply **Nixlo IGR Concentrate** alone or tank mixed with an adulticide at the rate of 4 ml per 1,500 sq. ft. of surface in sufficient volume of water to thoroughly wet the surface, although not to the point that runoff occurs. Use up to 12 ml of concentrate per 1,500 sq. ft. of surface for insect populations that are difficult to control or in conditions that make it difficult to achieve desired results.

Space Spray, Void Injection, ULV and Fog Application:

Read and follow the “**ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION**” in the Application Instructions section of this label. Apply **Nixlo IGR Concentrate** with a ULV generator or fogger at a rate of 1 ml per 1,000 cu. ft. If used alone, dilute concentrate with water at the rate of 1 ml of concentrate with 1 fl. oz. (30 ml) of water or oil, and fog solution at a rate of 1 fl. oz. (30 ml) per 1,000 cu. ft. When tank mixing with a pyrethrum fogging concentrate, mix **Nixlo IGR Concentrate** directly with the pyrethrum fogging concentrate solution. Exit the treated area after fogging. Keep facility closed for 1 hour and thoroughly ventilate prior to reentry.

INTERIOR AND PERIMETER APPLICATION DIRECTIONS

Inspect and identify harborage and breeding areas. Treat all interior areas where crawling insects may collect, travel, rest, hide, harbor or breed, including vertical walls, ceilings, posts, pipes, conduit, around doors and windows, cages, stalls, stanchions, floors, under watering equipment and feeders, cracks and crevices along floors, animal bedding, manure and other known or potential breeding sites. Control will be enhanced when facilities are cleaned, and interior applications are supplemented with exterior perimeter treatments. Apply solution to exterior perimeter of facility in a uniform band at least 1 foot up and 6 to 10 feet out from foundation. Apply around the exterior surfaces of doors and windows, overhead areas and other potential points of harborage or entry. If there are raised floors with soil or debris beneath them, be sure to treat the area beneath the floor.

EXTERIOR AND OUTDOOR APPLICATION DIRECTIONS:

Inspect and identify known or suspected harborage and breeding areas. Treat known or potential exterior breeding sites such as garbage bins or refuse piles, around grain or feed storage facilities, soil and vegetation in fence lines, compost piles, manure piles, beneath watering equipment or feeders, in, around and under calf hutches, in and beneath outdoor cages, in corrals, farmyards and feed lots.

Reapply every 30 days, or when target insect populations return to economic threshold levels.



Use in Poultry Production

FOR CONTROL OF INSECT PESTS IN AND AROUND POULTRY HOUSING AND FACILITIES – APPLICATIONS ARE PERMITTED WHEN POULTRY ARE PRESENT

Nixlo IGR Concentrate may be used by pest control professionals and poultry producers in, on and around poultry houses and processing facilities to control listed flying insect pests. Applications can be made while poultry are present. **Nixlo IGR Concentrate** mimics naturally occurring insect growth hormones, disrupting the reproductive processes of listed insects, resulting in reduced egg production, reduced egg hatch rates, disruption or inhibition of normal molting processes, deformity, sterile adults and death. **Nixlo IGR Concentrate** effectively breaks the cycle of insect reproduction. It is an important component in an IPM program and is a means to prevent or manage known resistant insect populations. Used as directed, **Nixlo IGR Concentrate*** will not harm beneficial insects such as predatory wasps.

*Pyriproxyfen

Apply as a spot treatment; general surface broadcast treatment, ULV/fog, spot or crack and crevice application to interior and exterior areas where insect pests collect, travel, rest, hide, harbor or breed.

Restriction: Do not contaminate food, feed, potable water or watering equipment. Initiate egg collection process prior to **Nixlo IGR Concentrate** application.

RATES TO CONTROL MOSQUITOES AND OTHER FLYING INSECTS

Nixlo IGR Concentrate affects both adult and juvenile insect stages. It decreases the number of eggs laid by treated adults and reduces hatch rates. Larvae (maggots) show physiological and functional defects leading to the inability to develop into normal adults. When used alone treatment will result in a reduction in adult fly populations after 3 to 4 weeks, when previously hatched adult flies have died naturally. For faster results, **Nixlo IGR Concentrate** may be tank mixed with an adulticide. Apply alone or tank mixed with an adulticide at the rate of 4 ml per 1,500 sq. ft. of surface area in sufficient volume of water to thoroughly wet the surface, although not to the point that runoff occurs. Use up to 12 ml of concentrate per 1,500 sq. ft. of surface for insect populations that are difficult to control or in conditions that make it difficult to achieve desired results.

RATES TO CONTROL LISTED CRAWLING INSECTS:

Apply **Nixlo IGR Concentrate** alone or tank mixed with an adulticide at the rate of 4 ml per 1,500 sq. ft. of surface in sufficient volume of water to thoroughly wet the surface, although not to the point that runoff occurs. Use up to 12 ml of concentrate per 1,500 sq. ft. of surface for insect populations that are difficult to control or in conditions that make it difficult to achieve desired results.

Space Spray, Void Injection, ULV and Fog Application:

Read and follow the “**ULV AND FOGGING APPLICATION INSTRUCTIONS, INCLUDING SPACE SPRAY AND VOID INJECTION**” in the Application Instructions section of this label. **Nixlo IGR Concentrate** may be applied with a ULV generator or fogger at a rate of 1 ml per 1,000 cu. ft. If used alone, dilute concentrate with water at the rate of 1 ml of concentrate with 1 fl. oz. (30 ml) of water, and fog solution at a rate of 1 fl. oz. (30 ml) per 1,000 cu. ft. When tank mixing with a pyrethrum fogging concentrate, mix **Nixlo IGR Concentrate** directly with the pyrethrum fogging concentrate solution. Exit the treated area after fogging. Keep facility closed for 1 hour and thoroughly ventilate prior to reentry.

INTERIOR AND PERIMETER APPLICATION DIRECTIONS:

Inspect and identify harborage and breeding areas. Treat all interior areas where crawling insects may collect, travel, rest, hide, harbor or breed, including vertical walls, ceilings, posts, pipes, conduit, around doors and windows, cages, stalls, stanchions, floors, under watering equipment and feeders, cracks and crevices along floors, animal bedding, manure and other known or potential breeding sites. Control will be enhanced when facilities are cleaned, and interior applications are supplemented with exterior perimeter treatments. Apply solution to exterior perimeter of facility in a uniform band at least 1 foot up and 6 to 10 feet out from foundation. Apply around the exterior surfaces of doors and windows, overhead areas and other potential points of harborage or entry. If there are raised floors with soil or debris beneath them, be sure to treat the area beneath the floor.

EXTERIOR AND OUTDOOR APPLICATION DIRECTIONS:

Inspect and identify known or suspected harborage and breeding areas. Treat known or potential exterior breeding sites such as garbage bins or refuse piles, around grain or feed storage facilities, soil and vegetation in fence lines, compost piles, manure piles, beneath watering equipment or feeders, in, around and under calf hutches, in and beneath outdoor cages, in corrals, farmyards and feed lots.

Reapply every 30 days, or when target insect populations return to economic threshold levels.

FOR CONTROL OF LITTER BEETLES

(Such as Darkling, Hide and Carrion Beetles)

Nixlo IGR Concentrate can be applied to litter, such as in poultry houses infested with litter beetles, to break the life cycle of these insect pests. Make uniform application of **Nixlo IGR Concentrate** to litter at the rate of 4 ml of **Nixlo IGR Concentrate** per 1,500 sq. ft. of surface in sufficient volume of water to thoroughly wet the surface, although not to the point that runoff occurs. Use up to 12 ml of concentrate per 1,500 sq. ft. of surface for insect populations that are difficult to control or in conditions that make it difficult to achieve desired results. **Nixlo IGR Concentrate** can be used prior to bird placement and/or used as a mid-flock treatment while birds are present.

For effective prevention of beetle larvae in litter, remove old litter from facility as often as possible. After new litter has been placed, treat top of litter with **Nixlo IGR Concentrate** and an adulticide. If litter is not changed between flocks, treat litter immediately prior to bird placement. Surface treatments immediately after animals are removed and again just prior to new animal placement are most effective. Adult egg-laying beetles can be killed with an adulticide. Application of adulticide to walls and floors at cleanout, before reintroduction of animals, will suppress beetles that escaped earlier treatment and will help delay onset of future infestations. Pay attention to areas where beetles frequently occur, such as walls, supports, cages, stalls and around feeders.

Mid-flock treatments for litter beetle control: Mix Nixlo IGR Concentrate with an adulticide approved for use while birds are present. Apply mixture to areas where beetles frequently occur, such as walls, supports, cages, under water lines, and around feeders.

Apply **Nixlo IGR Concentrate** uniformly to floor area and to walls, posts, and cage framing (birds grown in cages only). Application should also be made into cracks and crevices around insulation. Reapply after each grow-out or sanitization procedure. Indoor control can be enhanced by making perimeter treatments around the outside of building foundations to prevent immigrating adult beetles. Apply **Nixlo IGR Concentrate** or adulticide insecticides, or both, in a uniform band at least 1 foot up and 6 to 10 feet out from foundation. Maintaining a year-round treatment program will prevent background populations from reaching problem levels.

USE AS A MOSQUITO LARVICIDE

Active only on the immature life stages of mosquitoes. Tank mix or use with an adulticide for immediate kill of biting (adult) mosquitoes.

Restrictions:

- Not for use in natural holding waters or active waterways.
- Not for wide area mosquito use.
- Do not apply using truck mounted sprayers or aerial application equipment.

RESIDENTIAL AND COMMERCIAL SITES

Apply using a backpack sprayer, ULV cold fogger, compressed air sprayer, power sprayer or handheld sprayer at a rate of 4 to 8 ml of **Nixlo IGR Concentrate** per 1,500 sq. ft. in sufficient amount of water to get good coverage of target site. Use up to 12 ml per 1,500 sq. ft. for insect populations that are difficult to control or in conditions that make it difficult to get control of target pests. Alternate mixing instructions: use 16 to 48 ml of concentrate per 50 gallons of water.

Treat moist, shady areas where mosquitoes rest, harbor or breed, such as trees, shrubs, landscape plants, tall grass or plant foliage, non-drinkable standing water, gutters, drains, under decks, in-ground sprinkler heads, pet areas, fountains, birdbaths, landscape timbers, rocks, pavers, planters and fences. Tank mix **Nixlo IGR Concentrate** with an adulticide to get immediate control of the biting adult stages, while inhibiting successful development of immature stages. Monitor insect emergence and re-apply every 28 days, or when insect populations reach treatment threshold levels.



STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For Totes/55 gal containers: Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Nixlo™ is a trademark of Atticus, LLC.

NyGuard® is a registered trademark of McLaughlin Gormley King Company.

20230309ap1

