

INDOXACARB GROUP 22A INSECTICIDE



Advion[®] Cockroach

Gel Bait

syngenta.

Insecticide

FOR USE IN RESIDENTIAL, INSTITUTIONAL, COMMERCIAL, AND INDUSTRIAL AREAS

Use sites include single and multi-family residential buildings, schools, commercial and industrial facilities (including warehouses, apartments, supermarkets, restaurants, motels, hotels, hospitals, food handling/storage establishments), and transportation equipment such as aircraft, trains, ships, boats, buses.

**KEEP OUT OF REACH OF CHILDREN
CAUTION/
PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DO NOT TREAT AREAS THAT ARE EASILY ACCESSIBLE TO CHILDREN AND PETS

Active Ingredient:
Indoxacarb: (S)-methyl 7-chloro-2,5-dihydro-2-[[methoxycarbonyl] [4(trifluoromethoxy) phenyl]amino] carbonyl]indeno[1,2-e] [1,3,4]oxadiazine-4a-(3H) carboxylate 0.6%

Other Ingredients 99.4%
Total: 100.0%

EPA Reg. No. 100-1484
EPA Est. 54450-IL-002
Product of USA

**SCP 1484A-L1C 0620
4123793**

**4 x 1.06 oz (30g)
syringes**

**Net Weight
Non-refillable Container**



(01) 0 07 02941 53206 9

Advion Cockroach Gel Bait

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance
(Human or Animal) Or Chemical Emergency Assistance
(Spill, Leak, Fire or Accident) Call **1-800-888-8372**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 24 hours. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Do not apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters).

Physical and Chemical Hazards

Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

PRODUCT INFORMATION

Advion® Cockroach Gel Bait is designed to control pest species of cockroaches such as German, American, Australian, Smoky-Brown, Brown, Asian, Oriental, and Brown-Banded.

INSECT RESISTANCE MANAGEMENT

INDOXACARB	GROUP	22A	INSECTICIDE
------------	-------	-----	-------------

For resistance management, Advion Cockroach Gel Bait contains a Group 22A insecticide. Any insect population may contain individuals naturally resistant to Advion Cockroach Gel Bait and other Group 22A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same treated areas. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of Advion Cockroach Gel Bait or other Group 22A insecticides with different groups that control the same pests.
- Contact your local Syngenta representative, retailer, or extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Syngenta at 1-866-SYNGENT(A) (866-796-4368).

Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

Use Instructions and Precautions:

Do not treat areas that are easily accessible to children and pets.

1. Advion Cockroach Gel Bait can be used indoors or outdoors applied as a crack and crevice treatment within residential homes, industrial facilities, offices, warehouses, commercial kitchens, hospitals, schools, nursing homes, hotels, buses, trains, aircraft, retail and commercial establishments, and other areas infested with cockroaches.
2. Apply Advion Cockroach Gel Bait to areas frequented by cockroaches such as between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, and behind and under appliances, cabinets, sinks, closets, and any other equipment that cockroaches are typically known to hide.
3. Do not apply Advion Cockroach Gel Bait to surfaces recently treated with residual sprays.
4. Do not apply residual sprays to surfaces treated with Advion Cockroach Gel Bait.
5. Inspect bait placements during follow-up inspections and apply fresh bait as needed.
6. Do not apply to surfaces that food contacts.
7. Do not allow open foods to contact gel material.

Advion Cockroach Gel Bait

Treatment Rates for All Uses

- For heavy infestations of cockroaches, apply 3 to 5 spots of Advion Cockroach Gel Bait per 10 linear feet.
- For light to moderate infestations of cockroaches, apply 1-3 spots of Advion Cockroach Gel Bait per 10 linear feet.
- Each spot of Advion® Cockroach Gel Bait must equal about 0.5 grams (approx. 1/4" diameter)
- For all levels of cockroach infestations, when a bead of Advion Cockroach Gel Bait is used, apply a small thin bead that is less than 1/8" wide and approximately 2 inches long.

Indoor Use Directions

- Infested areas must be carefully inspected to determine overall infestation levels, locations of harborage areas, and most appropriate application points.
- Apply Advion Cockroach Gel Bait as a small spot of material, approximately 0.5 grams (approx. 1/4" in dia.), to areas not able to be contacted by children or pets.
- Several small applications distributed within an area are more effective than 1 or 2 applications of large quantities. In some cases a small thin bead (less than 1/8" wide and 2" long) may be appropriate such as between construction elements or other cracks and crevices.
- Indoor applications include cracks and crevices along walls or floors, behind or under equipment, under tables, within framing of tables or furniture, around sinks, within or near garbage collection areas, attics, crawl spaces, and cracks and crevices in cabinets.
- Inspect Advion Cockroach Gel Bait placements periodically for consumption or attractiveness and reapply as needed.

Applications in Nonfood/Nonfeed Handling Areas

Advion Cockroach Gel Bait is designed for use as a spot or crack and crevice treatment to control cockroaches in residential structures and the non-food/non-feed areas of commercial, industrial, public and institutional buildings/structures, including restaurants, warehouses, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, schools, laboratories, computer facilities, aircraft, buses, boats/ships, trains, pet shops and zoos.

Nonfood/non-feed areas include areas such as garbage rooms, lavatories, floor drains (to sewers), entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after bottling or canning). Refer to section below for use directions and restrictions when making applications in food/feed handling areas of Food/Feed Handling Establishments.

Applications in Food/Feed Handling Areas

Application to food/feed areas of food/feed handling establishments may only be made as a crack and crevice treatment only. Apply Advion Cockroach Gel Bait directly into cracks and crevices by placing the bait dispenser tip approximately 1/2 inch into cracks, crevices, holes and similar openings where cockroaches are found or are entering.

Food and feed handling areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation.

- Do not apply Advion Cockroach Gel Bait to areas where food/feed, food utensils or food processing surfaces may come into contact and become contaminated.
- Do not apply Advion Cockroach Gel Bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation and prepared food holding surfaces as bait could be removed by the washing.
- Do not apply Advion Cockroach Gel Bait to surfaces where the temperature exceeds 120-130°F because this could cause the bait to liquefy and lose effectiveness.
- Examples include, but are not limited to, parts of stoves, ovens, grills, fume hoods, saunas, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers, and hot water pipes.
- Use good application practice to avoid movement of Advion Cockroach Gel Bait to exposed surfaces used for food preparation.
- If Advion Cockroach Gel Bait contacts an exposed surface, first remove as much of the gel bait as possible, then wash with soap and water.

Outdoor Use Directions:

A thorough inspection around a structure may reveal areas where cockroaches are gaining access. Attempts should be made to remove the access. Advion Cockroach Gel Bait may also be applied as a spot or thin bead (see above) to pest entry sites such as along windows, doors, between construction elements, sewer areas, adjacent trees which harbor cockroaches, along walls, garbage holding areas, or suitable placement site acting as a cockroach harborage. Place baits in such a way as to reduce the potential of subsequent human contact. Inspect Advion Cockroach Gel Bait placements periodically and apply fresh bait as needed.

Advion Cockroach Gel Bait

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage

Store in cool, dry place inaccessible to children and pets.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal

Nonrefillable Container: Do not reuse or refill this container. Offer for recycling if available. Place empty bait dispensers in trash. If partially filled, wrap in newspaper and discard in trash or call your local solid waste authority for disposal instructions.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and of Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Advion®, the ALLIANCE FRAME 
the SYNGENTA Logo and the PURPOSE ICON 
are Trademarks of a Syngenta Group Company

©2020 Syngenta

For non-emergency (e.g., current product information),
call Syngenta Crop Protection at 1-800-334-9481.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1484A-L1C 0620

SPECIMEN Alpine® WSG

Water Soluble Granule Insecticide

SMART SOLUTIONS FACTS

Key attributes of this product include:

- Non-repellent active ingredient
- May be used in conjunction with non-repellent insecticides such as those containing dinotefuran (e.g., **Alpine® Cockroach Gel Bait Rotation 1 Reservoir** and **Alpine® Cockroach Gel Bait Rotation 2 Reservoir**) or similar products. Always follow label use directions and applicable laws.
- Ant control product
- Rotational partner for bed bug control
- For use on Stored Product Pests

KILLS: Ants (including Argentine, Carpenter, Harvester, Odorous House, Red Imported, and Southern Fire; excluding Pharaoh), Asian Lady Beetles, Bean Plataspid (Kudzu Bug), Bed Bugs, Beetles (including Confused Flour Beetle, Granary Weevil, Lesser Grain Borer, and Warehouse Beetle), Brown Marmorated Stink Bugs, Centipedes, Cockroaches (including American, German and Turkestan), Collembola, Crickets, Earwigs, Fleas, House Flies, Millipedes, Mosquitoes (*Anopheles*), Paper Wasps, Pillbugs, Sowbugs, Spiders (excluding Brown Recluse), Stored Product Pests (Confused Flour Beetle, Granary Weevil, Lesser Grain Borer, and Warehouse Beetle) and Yellowjackets

FOR USE IN AND AROUND: Animal Housing (including Calf Hutches, Calving Pens and Parlors, Chicken Houses, Dairy Areas, Dog Kennels, Hog Barns, Horse Barns, Milk Rooms and Rabbit Hutches); Apartments; Campgrounds; Cemeteries; Clinics; Food Handling Establishments; Garbage or Refuse Bins and Receptacles; Grain Mills; Granaries; Green Belts; Homes; Hospitals; Hotels; Mausoleums; Mobile Homes; Motels; Food Processing Plants; Nursing Homes; Resorts; Schools; Storage Buildings; Trailers; Transportation Equipment (including Autos, Buses, Boats, Ships, Trains, Trucks, Planes - cargo areas only); Utilities; Warehouses; Commercial, Industrial, and Residential Buildings; Theme Parks; Parks and Recreational Areas and Zoos

Active Ingredient:

Dinotefuran: N-methyl-N-[(tetrahydro-3-furanyl)methyl]guanidine 40.0%

Other Ingredients: 60.0%

Total: 100.0%

EPA Reg. No. 499-561

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See back booklet for **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific pest and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Weight:

Produced For:
BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


We create chemistry

FIRST AID

If in eyes

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes; then continue rinsing.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves (made out of any waterproof material, such as Natural Rubber, selection Category A).

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to aquatic invertebrates. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. **DO NOT** dispose equipment washwaters or rinsate into a natural drain or water body.

This product is toxic to honey bees. The persistence of residues and potential residual toxicity of dinotefuran in nectar and pollen suggests the possibility of chronic toxic risk to honey bee larvae and the eventual instability of the hive.

- This product is toxic to bees exposed to residues for more than 38 hours following treatment.
- **DO NOT** apply this product to blooming, pollen-shedding or nectar-producing parts of plants if bees may forage on the plants during this time period, unless the application is made in response to a public health emergency declared by appropriate state or federal authorities.

Groundwater Advisory

Dinotefuran and its degradate, MNG, have the properties and characteristics associated with chemicals detected in groundwater. The high water solubility of dinotefuran and its degradate, MNG, coupled with its very high mobility and resistance to biodegradation indicates that this compound has a strong potential to leach to the subsurface under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PROTECTION OF POLLINATORS



APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.

Look for the bee hazard icon  in the **Directions For Use** for each application site for specific use restrictions and instructions to protect bees and other insect pollinators.

This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications.
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When using this product take steps to:

- Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
- Minimize drift of this product onto beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

(continued)

PROTECTION OF POLLINATORS *(continued)*

Information on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: <http://pesticidestewardship.org/PollinatorProtection/Pages/default.aspx>.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

Physical and Chemical Hazards

DO NOT apply water-based sprays of this product to conduits, motor housing, junction and switch boxes or other electrical equipment because of possible shock hazard.

DO NOT spray where electrical short circuits might result, such as wall outlets, conduits, etc.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.



DO NOT apply this product while bees are foraging. DO NOT apply this product to plants that are flowering. Only apply after all flower petals have fallen off.

Use Restrictions

- **DO NOT** apply within aircraft cabins.
- **DO NOT** use in outdoor residential misting systems.
- **DO NOT** apply to humans or their clothing or to mattresses. **DO NOT** allow children or pets to contact treated surfaces until spray has dried.
- **DO NOT** apply to household pets. Cover fish tanks prior to application near tanks.
- **DO NOT** apply this product in patient rooms while occupied.
- **DO NOT** apply to classrooms when in use.
- When applying in residential or commercial structures, **DO NOT** contaminate food, feed stuffs or water supply. **DO NOT** contaminate food preparation surfaces, kitchen utensils, dishes or food/feed storage containers. Cover any food/feed contact surfaces and cooking utensils in the treatment area before treatment or thoroughly clean after treatment and before using.
- Apply to the point just prior to runoff. **DO NOT** allow applications of product to drip or run from surfaces.

- When applying to structural areas in and around live-stock housing, remove animals or treat in areas where animals are not present. **DO NOT** apply to animals. **DO NOT** allow insecticide to drift onto animals. **DO NOT** contaminate milk, food or drinking water.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry area away from heat or open flame.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance.

Container Handling

Single-use Packets

Nonrefillable container. DO NOT reuse or refill this container. Offer empty packet for recycling, if available, or dispose of in the trash or by incineration.

Nonrefillable container. DO NOT reuse or refill this container.

Triple rinse container promptly after emptying; then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Triple rinse container as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Product Information

This product is a water-soluble granule containing the non-repellent active ingredient dinotefuran. Apply indoors and/or outdoors via crack and crevice, spot or surface application for the listed pests and use sites.

Mixing Instructions

Thoroughly clean spray equipment before mixing and application of this product.

Single-use Packets: 0.35 ounce (10 grams)

Add approximately 1/2 of the water to the spray tank. Tear open the packet and add contents to water, then complete the filling of the spray tank. Agitate the tank until the product is completely dissolved. Reagitate finished dilution if application is interrupted for an extended period of time.

Resealable Bottle:

Add approximately 1/2 of the water to the spray tank. Add 10 to 30 grams of product to water per 1,000 square foot area being treated (1 Tablespoon = 10 grams), then complete the filling of the spray tank. Agitate the tank until the product is completely dissolved. Reagitate finished dilution if application is interrupted for an extended period of time.

SPRAY DILUTION CHART

Desired Finished Dilution Concentration	Grams of product to make 1 gallon of finished dilution to treat up to 1,000 square feet	Grams of product to make 5 gallons of finished dilution to treat up to 5,000 square feet	Grams of product to make 25 gallons of finished dilution to treat up to 25,000 square feet	Grams of product to make 50 gallons of finished dilution to treat up to 50,000 square feet
0.1%	10	50	250	500
0.2%	20	100	500	1,000
0.3%	30	150	750	1,500

Refer to specific pest categories below for the proper use rate. **One gallon of finished dilution treats 1,000 square feet.** Depending on the surface, more or less water may be used to properly treat areas with a different application apparatus, but the proper amount of active ingredient should be used per square footage.

Tank Mixing

This product may be tank mixed with other water based insecticides including, but not limited to, insect growth regulators (IGR), quick knockdown products and residuals. Before mixing, test to see if products are compatible. When tank mixing, follow the more restrictive label use directions of the mixed products.

Void/Crack & Crevice® Application

Apply into cracks, crevices and voids with equipment designed for such applications. Follow the manufacturer's recommendations for proper equipment set-up and operating pressures. Place the tip of the applicator at or into the crack, crevice or void to be treated. Apply product in short bursts while keeping tip in position. Allow the air stream to push the finished dilution into voids and/or cracks and crevices. Apply in such a manner to avoid dripping and runoff.

Indoor Applications

Dilute to the proper concentration according to targeted pest infestation. Follow the mixing directions found in the **SPRAY DILUTION CHART**. Use one gallon of finished dilution per 1,000 square feet. Refer to **INDOOR APPLICATIONS** table for use rates and pests.

INDOOR APPLICATIONS

Pests	Finished Dilution (grams per gallon)*	Application Instructions
Ants (excluding Pharaoh)	10	Make spot, Crack & Crevice ® and/or void applications where ants may be foraging and/or nesting, such as, around doors, window frames, under sinks, around pipes and in pipe voids, attic venting and other areas where ants may find access or nest.
Bed Bugs	10 to 30	Make spot, Crack & Crevice and/or void applications where evidence of bed bug infestation occurs or bed bugs are suspected, or likely to occur, such as bed frames; box springs; inside empty dressers, clothes closets and luggage; carpets; draperies; furniture; headboards; high and low wall moldings and wallpaper edges. DO NOT apply to mattresses.
Crawling Insects/Pests including: Pillbugs, Spiders (excluding: Brown Recluse), and Turkestan Cockroaches	30	Make spot and/or Crack & Crevice applications into hiding places and entry points such as cracks and crevices, void areas, moist areas, openings around pipes and sinks, under refrigerators or vending machines, interiorscapes, across lower doorway frames and window openings.
Crickets, Cockroaches (American and German), and Millipedes	10 to 30	
Fleas	10	Apply to infested areas or potentially infested areas, such as rugs, floors, carpets, upholstered furniture, pet beds and pet resting areas. When applying to upholstered furniture, treat under cushions and areas where flea development can occur. DO NOT treat pets with this product.
House Flies	10 to 20	Make spot applications on surfaces where flies frequently roost such as door and window frames, screens, trash cans, plants, interiorscapes, ceilings and upper wall surfaces.
When a rate range is indicated, use the lower rate for maintenance applications and the higher rates for heavy infestations. *Refer to SPRAY DILUTION CHART for concentration of active ingredient in finished product.		

Food/Feed Handling Establishments

Food/Feed handling establishments are places other than private residences in which food is held, processed, prepared or served, including those operating under the Federal meat, poultry, shell egg grading and egg products inspection programs. **APPLICATIONS OF THIS PRODUCT IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS MAY BE MADE AS A CRACK & CREVICE, VOID, SPOT OR GENERAL TREATMENT.** General surface application may be used only when the facility is not in operation provided exposed food/feed has been covered or removed from the area prior to application.

Food/Feed areas include areas for receiving, serving, storing (dry, cold, frozen, raw), packing (canning, bottling, wrapping, boxing), preparing (cleaning, slicing, cooking, grinding), edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups).

Non-food/feed areas include areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after packaging, canning or bottling).

Avoid contamination of food/feed or food/feed contact surfaces. Remove or cover food/feed, dishes, utensils, food processing equipment and food preparation surfaces in the treatment area or wash them before use. Apply as a **Crack & Crevice**, void, spot or general treatment to surfaces such as baseboards, under elements of construction, stainless steel equipment, shelving, machinery, storage areas, pallets, tables, chairs and other areas where listed pests may be harboring, traveling, breeding or entering the structure. Maximum use rate is 0.05 grams of active ingredient per linear foot. Reapplications may be made at 3 day intervals when applying as a **Crack & Crevice**, void or spot treatment, or at 6 day intervals when applying as a general surface treatment. This product can also be used as a preventative treatment for areas likely to have future infestations.

Indoor and Outdoor Applications

Dilute to the proper concentration according to targeted pest infestation. Follow the mixing directions found in the **SPRAY DILUTION CHART**. Use one gallon of finished dilution per 1,000 square feet. Refer to **INDOOR/OUTDOOR APPLICATIONS** table for use rates and pests.

In the state of New York, **DO NOT** apply outdoors except for **Crack & Crevice®** or spot treatments to structural components, stumps, utility poles, and fences.

INDOOR/OUTDOOR APPLICATIONS		
Pests	Finished Dilution (grams per gallon)*	Application Instructions
Asian Lady Beetles and Centipedes	10 to 30	Make spot, Crack & Crevice and/or void applications where these pests may harbor or hibernate, such as cracks and crevices, in weep holes, wall voids, around window and door frames, attics and behind siding. Apply to exterior wall surfaces around entry points and resting areas where insects congregate.
Bean Plataspid (Kudzu Bug), Brown Marmorated Stink Bugs	30	
Carpenter Ants	10 to 20	Make spot, Crack & Crevice and/or void applications to any trails, inside or outside, around doors and windows and other places where ants may enter premises (such as atriums, attic venting, crawl spaces, ceilings, fences, exterior building surfaces, landscape timbers, garages, hollow doors, soffits, wall voids, weep holes and other void areas). Where possible, directly apply to ant nests or infested wood or trees.
Paper Wasps and Yellowjackets	30	To treat a yellowjacket or wasp nest, apply liberally to the nest and the surrounding area. For nests inside wall voids, inject an amount sufficient to treat the entire nest. Most applications should require about 0.5 gallon of diluted material. Pay attention to surfaces or items that the product may contact (i.e. electrical wiring). DO NOT use where there is an electrical shock hazard. THIS PRODUCT DOES NOT PROVIDE INSTANT KNOCKDOWN.
Stored Product Pests (Confused Flour Beetle, Granary Weevil, Lesser Grain Borer, and Warehouse Beetle)	10	Before storing products, treat warehouses, production facilities, storage areas, rail cars, truck beds, and other areas where products are stored. Apply as a spot and/or Crack & Crevice treatment to cracks, crevices, and surfaces where pests have been seen or may have harborage. This product may be used to treat empty grain storage facilities and other listed areas for stored product pest control. Cleaning areas prior to use of this product will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.
When a rate range is indicated, use the lower rate for maintenance applications and the higher rates for heavy infestations. *Refer to SPRAY DILUTION CHART for concentration of active ingredient in finished product.		

PREVENTATIVE PROGRAMS: This product may be used as part of a preventative program or in anticipation of a pest problem associated with a structure or an event.

Outdoor Applications

Dilute to the proper concentration according to targeted pest infestation. Follow the mixing directions found in the **SPRAY DILUTION CHART**. Use one gallon of finished dilution per 1,000 square feet. Refer to **OUTDOOR APPLICATIONS** table for use rates and pests.

DO NOT apply this product, by any application method, to linden, basswood or other *Tilia* species in the State of Oregon.

In the state of New York, **DO NOT** apply outdoors except for **Crack & Crevice**® or spot treatments to structural components, stumps, utility poles, and fences.

OUTDOOR APPLICATIONS		
Pests	Finished Dilution (grams per gallon)*	Application Instructions
Ants (excluding Pharaoh)	10	Make spot and/or Crack & Crevice applications to any foraging trails, nests or mounds where ants are present. Make spot and/or Crack & Crevice applications to exterior areas where ants may enter a structure. Where possible, directly apply to ant nests or infested wood or trees. Curative broadcast applications can be made to yards, lawns, fields, parks, cemeteries, landscaping and structures. To help prevent infestations, make exterior structural and surrounding area applications according to this label.
Exterior Structural and Surrounding Area Applications for Collembola (Springtails), Crickets and Millipedes	10 to 30	Directly contacting pests, apply where pests rest, trail and harbor, such as structural and other voids (including voids associated with and around doors, windows, utility entry points, behind siding, weep holes and tree hollows), eaves of structures, around lights or other insect attractant areas, foliage, bushes, landscape materials, garbage receptacles, under trees and other shaded areas.
Exterior Structural and Surrounding Area Applications for Crawling Insects/Pests including: Earwigs, Pillbugs, Sowbugs, Spiders (excluding Brown Recluse), and Turkestan Cockroaches	30	
Fleas	10	Apply to yards, porches, patios or other areas of habitation for fleas, especially areas that pets frequent, directly contacting fleas. Pay particular attention to shaded areas with low protective covering such as under bushes, under decks, lawn furniture, lawn ornaments and other landscape materials. Apply to pet entry areas such as steps and stoops and other areas where pets rest or sit for extended periods of time, directly contacting fleas. DO NOT treat pets with this product.
House Flies	10 to 20	Apply to outside surfaces of screens, doors, window frames, porches, in garages, on foliage or wherever flies enter or congregate around or on the building, directly contacting flies. Application sites include inside and outside surfaces of dumpsters, other trash holding containers and surrounding surfaces, directly contacting flies.

(continued)

OUTDOOR APPLICATIONS *(continued)*

Pests	Finished Dilution (grams per gallon)*	Application Instructions
Mosquitoes (<i>Anopheles</i>)	20	<p>Apply using handheld, back-pack or truck mounted sprayers only. Not for use with truck mounted or handheld cold aerosol ULV sprayers, or thermal fogging devices. Use up to 50 gallons of diluted spray per 50,000 square feet. Apply to shrubbery, vegetation and other areas where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas, ponds and shorelines may be treated if applied as a wet spray. Direct application of this product to any body of water is prohibited.</p> <p>DO NOT apply by aircraft. Not approved for use on mosquitoes in the state of New York.</p>
<p>When a rate range is indicated, use the lower rate for maintenance applications and the higher rates for heavy infestations.</p> <p>*Refer to SPRAY DILUTION CHART for concentration of active ingredient in finished product.</p>		

USE IN AND AROUND LIVESTOCK HOUSING

For structural use indoors and outdoors of animal housing. Remove animals or treat in areas where animals are not present. **DO NOT** apply to animals. **DO NOT** allow insecticide to drift onto animals. **DO NOT** contaminate milk, food or drinking water.

To kill house flies on outdoor surfaces and in animal housing including calf hutches, calving pens and parlors, chicken houses, dairy areas, dog kennels, hog barns, horse barns, milk rooms and rabbit hutches: Treat normal fly resting areas such as ceilings, screens, upper corners of the area, around windows and doors, light fixtures and other surfaces known to attract flies. Apply as a coarse, wet spray. Using a mixture of 10 to 20 grams of product per gallon of water, apply at a rate of 1 gallon diluted product per 1,000 square feet of surface. For best results, initiate treatments at the beginning of the fly season and repeat during periods of heavy fly activity.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as environmental conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer. BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW: (A) BASF MAKES NO OTHER WARRANTIES EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY, (B) BUYER'S EXCLUSIVE REMEDY AND BASF'S AND SELLER'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT, AND (C) BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, INCIDENTAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** BASF and the Seller offer this product, and the Buyer and User accept it, subject to these **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. PCS915

Alpine and Crack & Crevice are registered trademarks of BASF.

© 2019 BASF Corporation
All rights reserved.

000499-00561.20190708.**NVA 2019-04-477-0122**
Based on: NVA 2019-04-477-0037
Supersedes: NVA 2018-04-477-0049

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**
We create chemistry

CONTROL OF BEDBUGS

To treat bedbugs, apply to harborage points mentioned above; in addition, under beds, bottom side of box springs, under and behind night stands, to points where carpet and wall meet, cracks in cove base, molding, and other areas where bed bugs harbor. The foam formulation reduces the chances of missed harborage areas. Retreat at 14 days for heavy infestations.

Restrictions

- Do not treat linens and mattresses

CONTROL OF FLEAS

The active ingredients in this product are (S)-Hydroprene, Novaluron and Pyriproxyfen, which are insect growth regulators. Insect growth hormones are naturally occurring in insects and this product prevents the adult pest from developing. These properties will gradually result in the reduction and eventual control of the pest infestation. Flea eggs and flea larvae that come into contact with this product will also not develop into adult fleas. This product works for up to 7 months on fleas by inhibiting the development of the immature stages of the flea and preventing them from reaching the biting adult stage. This product does not work on adult fleas and flea pupae that already exist.

Use Sites	Use Directions
Surface Application	For use on carpets, furniture, pet sleeping areas & throw rugs, etc. Repeat treatment every 7 months as necessary. If needed, repeat treatments may be made 21 days after previous treatment.

WAREHOUSES (NON-FOOD) APPLICATIONS

Apply as a surface, spot, or crack and crevice treatment in non-food storage warehouses. Apply to all areas that may harbor pests, including under and between pallets. Apply foam to undersides of shelving and storage areas.

FOOD HANDLING ESTABLISHMENT APPLICATIONS

Apply as a surface spray, spot, or crack and crevice treatment in both food and non-food areas of food handling establishments. These establishments are defined as: places other than private residences where exposed food is held, prepared, processed, or served. It also includes areas where food items are: received, prepared, packaged, and stored, as well as enclosed food processing systems (mills, dairies, etc.).

For control of drain flies and fruit flies: Apply to areas of fly breeding. For drains, apply foam into and around the downspout and catch basin area. Coat these areas thoroughly as fly larva develop in the scum buildup on the sides of drain pipes. Visible foam formulation ensures no areas are missed. Repeat applications at 2 to 4 week intervals depending upon the severity of infestation.

SPOT OR CRACK AND CREVICE APPLICATIONS

Prior to application, remove or cover all food items in area to be treated. Do not apply directly to food or food-handling surfaces, but applications may be made above these areas due to the no-drip formulation. Spot or crack and crevice applications can be made while facility is in operation.

SURFACE SPRAY APPLICATION RESTRICTION: Do not apply directly to food products.

AREA TREATMENTS: Prior to application, remove or cover all food items.

PARTIAL AREA TREATMENTS: For applications during facility operation, apply to partial areas of food handling establishments in cases of small or localized infestations.

PANTRY/CABINETS, SMALL FOOD STORAGE AREAS

Apply foam as a surface spray, crack and crevice spot, or spot treatment to pantry and cupboard shelves or other areas where food is stored. Be sure to treat the undersides of shelving.

Restrictions

- Prior to spraying, remove all food containers and foodstuffs
- Do not replace food containers or foodstuffs until spray has dried
- Dispose of any infested food packages or materials

WAREHOUSES (FOOD)

Apply as a surface spray, spot, or crack and crevice treatment in warehouses where packaged food items are stored in such a way that direct contact with the foodstuff is not anticipated. Contains no CFCs or other ozone depleting substances. Federal regulations prohibit CFC propellants in aerosols.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store upright at room temperature. Do not store near heat or open flame. Store in an area away from children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Do not puncture or incinerate! Non-refillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions.

LIMITATIONS OF LIABILITY

To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in control, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc.'s election, the replacement of the product.



Manufactured by:
Control Solutions Inc.
5903 Genoa-Ried Bluff Rd
Pasadena, Texas 77507
A Member of the ADAMA Group



HYDROPRENE	GROUP	7A	INSECTICIDE
NOVALURON	GROUP	15	INSECTICIDE
PYRIPROXYFEN	GROUP	7	INSECTICIDE



TEKKO® TRIO FOAM

Pressurized Solutions

- Aerosol with three Insect Growth Regulators
- Breaks the life cycle of drain flies
- Labeled for use in food areas of food handling establishments
- Inhibits reinfestation by cockroaches for 6 months

ACTIVE INGREDIENTS

(S)-Hydroprene:	0.36%
Novaluron:	0.02%
Pyriproxyfen:	0.02%
OTHER INGREDIENTS:	99.6%
TOTAL:	100.00%

EPA Reg. No. 53883-475

EPA Est. No. 53883-TX-002

SHAKE WELL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact SafetyCall® (866) 897-8050 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Restrictions

- For indoor use only. DO NOT use outdoors. READ ALL DIRECTIONS COMPLETELY BEFORE USE.

Resistance Management

For resistance management, this product contains Group 7, 7A and 15 insecticides. Any insect population may contain individuals naturally resistant to this product and other Group 7, 7A and 15 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Use take mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.

NET WEIGHT: .9375 lb (15 oz)

- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
- When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pests.
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact your local extension specialist or sales representative.

PRODUCT INFORMATION

TEKKO TRIO FOAM contains three Insect Growth Regulators in a dry foam formulation. The dry foam formulation reduces the chances of missed areas and ensures complete coverage of an area. The IGRs in TEKKO TRIO FOAM disrupt the normal development of immature cockroaches, drain flies, fruit flies, bed bugs, fleas, and stored product pests (examples of these pests include: Indian meal moths, rice moths, tobacco moths, red flour beetles, confused flour beetles, lesser grain borers, merchant grain beetles, saw-tooth grain beetles, warehouse beetles, cigarette beetles, and Dermestid beetles). TEKKO TRIO FOAM prevents future generations of bed bugs and cockroaches. TEKKO TRIO FOAM also disrupts the normal life cycle of drain flies, fruit flies, and stored product pests; thus, killing these insects during the pupal stage. In both cases, the cycle of an infestation ends.

Use in areas including: apartment buildings, bakeries, bottling facilities, breweries, boiler rooms, cafeterias, candy plants, grocery stores, day care centers, hospitals, residential homes, hotels/motels, office buildings, kitchens, laboratories, cereal processing facilities, manufacturing plants, mausoleums, meat and produce canneries, nursing homes, restaurants, schools, locker rooms, stores, taverns, warehouses, pet stores, as well as various modes of transportation, such as aircraft cargo areas and galleys, buses, trucks, trailers, rail cars, and marine vessels.

Use as a surface spray, spot, or crack and crevice application.

Prior to using in commercial or residential food areas, please refer to the Food Handling Establishment Applications section of the label. The preparations prior to treatment and the clean-up after application described are also to be followed for residential use.

TEKKO TRIO FOAM is non-repellent and should be considered an important component in a cockroach baiting program. Baits are an important part of a total cockroach control program but are not a complete solution. TEKKO TRIO FOAM can reach those nymphs that fail to forage to baits and the IGRs stimulate egg-bearing females to leave the harborage and feed.

TEKKO TRIO FOAM is a ready-to-use pressurized spray for application to a variety of locations and surfaces to control pests. Apply as a surface spray, spot, or crack and crevice application. Use the supplied fold up extension tube for crack and crevice applications.

To apply as a surface or spot application, hold can 10" to 12" away from surface to be treated. Use a sweeping motion over the surface to be treated for complete coverage. Product applies as a dry foam that stays where you spray and does not run.

For crack and crevice applications, use the supplied fold up extension tube and apply to wall voids, hollow table legs, tube iron, gondola frames, voids, supports and other areas where pests may harbor. TEKKO TRIO FOAM is generally nonstaining, but in questionable situations, apply a small amount in an inconspicuous location. Apply once a month for continuous protection, or more often if conditions warrant.

SURFACE APPLICATION

Apply foam on surfaces at a rate of 15 ounces per 125 sq ft of surface area. Apply to areas such as: walls, floors, ceilings, attic, basements, gondolas, shelving and other areas where pests may harbor. The dry foam formulation will adhere to the undersides of shelving, cabinets, drawers, and other areas pests accumulate. Treat these areas thoroughly.

SPOT OR CRACK AND CREVICE APPLICATIONS

Apply dry foam as a crack and crevice application using the supplied fold up extension tube. Apply directly to areas such as: wall voids, pipe chases, crevices, false ceilings and other areas where pests may harbor. Apply spot treatments to areas such as: storage areas, closets, around water pipes, doors, windows, and behind and under appliances/equipment, furniture and beds. The dry foam formulation allows penetration deep into cracks and crevices to reach infestations that are not observed.

**Cleans
Without Harsh
Chemicals or
Odors**

InVade
technology™
Bio Zap™



DIRECTIONS FOR USE

InVade™ Bio Zap™ contains premium, natural, scum-eating, odor eliminating microbes. Citrus oil provides a fresh and pleasant scent. Product may be diluted with 1 to 2 parts water for spraying in an ongoing maintenance program. Spray walls, floors, carpets and fabrics where pet, smoke, dead animal, skunk or excrement odors originate, just to the point of wetness. Always check a small area for color fastness before use. Rinse garbage receptacles with water and then spray product liberally onto surfaces. Spray gravel or soil surfaces in crawl spaces, concrete in basements or garages and other areas where sewage

or wildlife have left odors. Spray into cracks, crevices, on floors, in garbage receptacles, bathroom walls, voids and urinals to eliminate odors. InVade Bio Zap may also be fogged into crawlspaces. Do not apply to direct food contact surfaces. Precautions: Keep out of reach of children. If more than an incidental amount is swallowed, seek medical attention immediately. Give one glass of water and do not induce vomiting. Wash hands well after use. If in eyes, remove contacts and flush with water for 15 minutes. Seek medical attention for any persistent irritation. Store in a closed, appropriately labeled container between 35 and 95 °F.

**Eating the
Scum of the
Earth**™

Rockwell Labs Ltd
creating microbial miracles™

816-283-3167 • www.rockwelllabs.com
North Kansas City, MO 64116



Net Contents: 1 Quart, 1 US Gallon