

Control Mature Weeds with Manuscript, Because They Grow Up so Fast

Mature weeds have met their match. Using a unique class of chemistry for turf, Manuscript[®] herbicide can be effectively sprayed at any weed growth stage. It controls **tropical signalgrass, crabgrass** and other grassy weeds any time they are actively growing and contains a built-in safener for outstanding turf safety.



syngenta.



@SyngentaTurf | #Time4Manuscript

A Unique Solution for Tough Weeds

Representing a novel class of chemistry for turf, Manuscript is a highly active herbicide that controls mature grassy weeds in certain warm-season grasses on **golf courses, residential and commercial lawns, sod farms and sports turf.** With Manuscript, you can get the application flexibility you need to treat mature weeds. It can be used in the heat of summer when desired turfgrass is actively growing and will fill in rapidly, and is formulated with a built-in, proprietary safener to enhance performance and turf safety. Manuscript is packaged with Adigor[™] surfactant from Syngenta, which is custom-built to be used with Manuscript to maximize the quantity and rate of absorption of *pinoxaden*, as well as the degree of translocation once *pinoxaden* is in the plant.



WEEDS CONTROLLED		
Tropical signalgrass	Large and smooth crabgrass	
Bull/thin paspalum	Dallisgrass	
Bahiagrass	agrass Seashore paspalum (suppression only)	
Tropical carpetgrass	Torpedograss (suppression only)	

TOLERANT TURFGRASS SPECIES

Bermudagrass
Zoysiagrass
St. Augustinegrass (sod only)



IMPROVED POST-EMERGENCE CONTROL OF TOUGH WEEDS

Selective herbicide with an advanced form of ACCase mode of action that works fast on tough mixed grasses regardless of growth stage, including large, mature weeds



OUTSTANDING TURF SAFETY

Includes a built-in safener (*cloquintocet-mexyl*) that speeds metabolism of *pinoxaden* in desirable turf and improves weed control performance:

- Enables spot treatments
- Further improves selectivity against tough, actively growing weeds without temperature restrictions
- Additionally, the MOA allows applications near ornamental beds and shrubs



LABOR SAVINGS

Reduced hand-weeding means greater labor savings

Manuscript is the most active tropical signalgrass herbicide I've ever applied. It's more active during summer months and [with Monument® 75WG herbicide] is about as close to a perfect herbicide solution as you're going to get."

– Neil Young, Owner, Turfgrass Environmental Research Inc.



QUICKER RECOVERY

Desirable turf quickly fills in voids left behind after weeds are controlled, especially in the summer when warm-season turf is actively growing

UNIQUE CHEMISTRY CLASS

With a novel class of chemistry for turf (phenylpyrazolin - DEN, Group 1), *pinoxaden* is rapidly absorbed and translocated to the growing points of weed foliage, where its powerful phenylpyrazolin inhibitor halts ACCase enzyme activity on tough grassy weeds for improved safety and resistance management

Customized Surfactant Maximizes Efficacy


Manuscript's safety on bermudagrass is incredible and definitely has at least a 2X safety."

- Young

Using Manuscript with Adigor provides greater application reliability. Adigor is an advanced surfactant system custom-built to be used with Manuscript to:

- Speed and maximize foliar absorption
- Increase spray droplet deposition
- Enhance droplet spreading

- Soften plant cuticle to speed and maximize absorption
- Improve rainfastness in as quickly as one hour



Time to rainfastness depends on atmospheric conditions, application volume, spray droplet size, thickness of plant cuticle and other factors.

Source: Muehlebach et al., 2011.

Performance assessments are based upon results or analysis of public information, field observations and/or internal Syngenta evaluations. Trials reflect treatment rates commonly recommended in the marketplace.

Use Recommendations

With the turf safety of Manuscript, applications are recommended between the end of May and the beginning of September (best timing depends on the target weed). This will encourage desirable grass to fill in quickly while it is actively growing.

While it can be applied as a broadcast treatment for control of bull/thin paspalum, spot treatments are recommended for difficult-to-control weeds like tropical signalgrass. Spot treatments can also be used to help control crabgrass breakouts in lawns. Manuscript must be tank-mixed with Adigor at 0.5% v/v to optimize spray droplet deposition, spreading and foliar absorption.

Manuscript has performed very well against tough weeds like crabgrasses. We've consistently seen a level of control up to and above 90 percent with excellent turf safety."

– Lane Tredway, Technical Services Manager, Syngenta

Spot Treatment Applications are Recommended for Optimal Results

RECOMMENDATION FOR HAND-HELD SPRAY WAND APPLICATIONS:

- Mix 0.96 fl. oz. of Manuscript and 0.64 fl. oz. of Adigor in 1 gal. of water to treat 1,000 ft.²
- Make applications in 40 gal./A
- Make a second application 14-21 days later when weed re-growth is observed

RECOMMENDATION FOR BOOM APPLICATIONS:

- Mix 9.6 fl. oz. of Manuscript and 6.4 fl. oz. of Adigor in 10 gal. of water to treat 10,000 ft.²
- Make applications in 40 gal./A using flat fan nozzles (standard or air induction) that produce medium to very coarse droplets
- Make a second application 14-21 days later when weed re-growth is observed

DO NOT treat more than 10,000 ft.2/A

DO NOT apply more than 19.2 fl. oz./A of Manuscript per calendar year



syngenta.

Program Approach to Control Secondary Weeds

After removing patches of undesirable weeds with Manuscript, secondary weeds like sedges and broadleaves tend to fill in. A follow-up application of Monument 75WG controls a broad spectrum of secondary weeds and will reduce weed reestablishment, so desirable turf can grow without weed competition.

HERBICIDE	RATE PER ACRE	APPLICATIONS	INTERVAL
Manuscript	9.6 fl. oz./10,000 ft. ²	2	14-21 days
Monument 75WG	0.53 oz./A	1	14 days

Manuscript causes little-to-no injury or growth regulation to bermudagrass or zoysiagrass when applied at broadcast or spot treatment rates.

OR

Manuscript Broadcast Applications (Bull/Thin Paspalum Only)


Two applications at 9.6 fl. oz./A on a 14- to 21-day interval (28-day interval for St. Augustinegrass)

One application at 19.2 fl. oz./A

Make applications in 40 gal./A for optimal control, and mix with 0.64 fl. oz. of surfactant per gallon of water (0.5% v/v). Supplement available Adigor with an MSO surfactant as needed to reach a total of 0.5% v/v.

Proven Weed Control

Manuscript has been evaluated for weed control across a series of field trials. The following results show the consistent efficacy of the product.



WAIT = weeks after initial treatment

Treatments applied twice on a 21-day interval. Manuscript mixed with Adigor at 0.5% v/v. Tribute® Total mixed with MSO at 0.5% v/v. Source: Neil Young, Miami, Florida, 2017. Performance assessments are based upon results or analysis of public information, field observation's and/or internal Syngent'a evaluations. Trials reflect treatment rates commonly recommended in the marketplace.



Square plots of crabgrass were treated with two spot treatments of Manuscript at 0.96 fl. oz./gal. and Adigor at 0.64 fl. oz./gal., 13 days apart. Huntsville, Alabama, 2018. Performance assessments are based upon results or analysis of public information, field observations and/or internal Syngenta evaluations. Trials reflect treatment rates commonly recommended in the marketplace.

For more information and additional trial data, please visit GreenCastOnline.com/Manuscript





All photos are either the property of Syngenta or are used with permission.

©2018 Syngenta. Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties and/or may have statespecific use requirements. Please check with your local extension service to ensure registration and proper use. Adjor²¹⁰, GreenCast[®], Manuscript[®], Monument[®], the Alliance Frame, the Purpose Icon and the Syngenta logo are trademarks of a Syngenta Group Company. All other trademarks are property of their respective owners. GS 3116_1_15 LGC 8683A 09-2018